



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 30, 2021 – 02:07 PM EST

PDB ID : 3DLI  
Title : Crystal structure of a SAM dependent methyltransferase from *Archaeoglobus fulgidus*  
Authors : Agarwal, R.; Burley, S.K.; Swaminathan, S.; New York SGX Research Center for Structural Genomics (NYSGXRC)  
Deposited on : 2008-06-27  
Resolution : 2.46 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.16  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.16


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1544 (2.48-2.44)
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)
RSRZ outliers	127900	1523 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	240	
1	B	240	
1	C	240	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5647 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called methyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	221	1830	1189	294	338	9	0	0	0
1	B	221	1830	1189	294	338	9	0	0	0
1	C	221	1830	1189	294	338	9	0	0	0

There are 33 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	242	MET	-	insertion	UNP O30190
A	243	SER	-	insertion	UNP O30190
A	244	LEU	-	insertion	UNP O30190
A	474	GLU	-	insertion	UNP O30190
A	475	GLY	-	insertion	UNP O30190
A	476	HIS	-	insertion	UNP O30190
A	477	HIS	-	insertion	UNP O30190
A	478	HIS	-	insertion	UNP O30190
A	479	HIS	-	insertion	UNP O30190
A	480	HIS	-	insertion	UNP O30190
A	481	HIS	-	insertion	UNP O30190
B	242	MET	-	insertion	UNP O30190
B	243	SER	-	insertion	UNP O30190
B	244	LEU	-	insertion	UNP O30190
B	474	GLU	-	insertion	UNP O30190
B	475	GLY	-	insertion	UNP O30190
B	476	HIS	-	insertion	UNP O30190
B	477	HIS	-	insertion	UNP O30190
B	478	HIS	-	insertion	UNP O30190
B	479	HIS	-	insertion	UNP O30190
B	480	HIS	-	insertion	UNP O30190
B	481	HIS	-	insertion	UNP O30190
C	242	MET	-	insertion	UNP O30190

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	243	SER	-	insertion	UNP O30190
C	244	LEU	-	insertion	UNP O30190
C	474	GLU	-	insertion	UNP O30190
C	475	GLY	-	insertion	UNP O30190
C	476	HIS	-	insertion	UNP O30190
C	477	HIS	-	insertion	UNP O30190
C	478	HIS	-	insertion	UNP O30190
C	479	HIS	-	insertion	UNP O30190
C	480	HIS	-	insertion	UNP O30190
C	481	HIS	-	insertion	UNP O30190

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	49	Total O 49 49	0	0
2	B	48	Total O 48 48	0	0
2	C	60	Total O 60 60	0	0



D438	S439	N440	E444	R448	E452	R459	P464	Y467	A468	I469	I470	A471	K472	K473	GLU	GLY	HIS						
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.15Å 81.88Å 122.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.12 – 2.46 41.12 – 2.46	Depositor EDS
% Data completeness (in resolution range)	96.9 (41.12-2.46) 97.0 (41.12-2.46)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.78 (at 2.45Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.230 , 0.286 0.230 , 0.287	Depositor DCC
$R_{free}$ test set	789 reflections (2.81%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.5	Xtrriage
Anisotropy	0.299	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 40.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	5647	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.38% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.45	0/1870	0.64	0/2515
1	B	0.47	0/1870	0.63	0/2515
1	C	0.47	0/1870	0.67	1/2515 (0.0%)
All	All	0.46	0/5610	0.64	1/7545 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	C	390	LEU	CA-CB-CG	5.74	128.51	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1830	0	1839	79	0
1	B	1830	0	1839	54	0
1	C	1830	0	1839	71	0
2	A	49	0	0	3	0
2	B	48	0	0	3	0
2	C	60	0	0	4	0
All	All	5647	0	5517	199	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 199 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:418:ARG:HD3	1:C:419:ASP:H	1.32	0.93
1:C:418:ARG:CD	1:C:419:ASP:H	1.85	0.89
1:B:404:HIS:HD2	1:B:406:GLU:H	1.21	0.88
1:A:310:VAL:HG12	1:A:327:VAL:HG13	1.54	0.87
1:C:404:HIS:HD2	1:C:406:GLU:H	1.23	0.87

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	219/240 (91%)	203 (93%)	14 (6%)	2 (1%)	17	19
1	B	219/240 (91%)	212 (97%)	7 (3%)	0	100	100
1	C	219/240 (91%)	212 (97%)	6 (3%)	1 (0%)	29	34
All	All	657/720 (91%)	627 (95%)	27 (4%)	3 (0%)	29	34

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	321	GLU
1	A	343	LEU
1	A	426	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	205/222 (92%)	193 (94%)	12 (6%)	19	25
1	B	205/222 (92%)	193 (94%)	12 (6%)	19	25
1	C	205/222 (92%)	193 (94%)	12 (6%)	19	25
All	All	615/666 (92%)	579 (94%)	36 (6%)	19	25

5 of 36 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	344	ASP
1	B	438	ASP
1	C	440	ASN
1	B	419	ASP
1	B	440	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	399	HIS
1	C	404	HIS
1	B	404	HIS
1	A	399	HIS
1	C	399	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	221/240 (92%)	-0.04	1 (0%) 91 92	17, 33, 44, 50	0
1	B	221/240 (92%)	-0.11	1 (0%) 91 92	17, 28, 39, 47	0
1	C	221/240 (92%)	-0.06	0 100 100	14, 28, 42, 53	0
All	All	663/720 (92%)	-0.07	2 (0%) 94 94	14, 30, 42, 53	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	337	SER	2.5
1	B	418	ARG	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.