



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 15, 2020 – 10:48 am BST

PDB ID : 1R5J  
Title : Crystal Structure of a Phosphotransacetylase from Streptococcus pyogenes  
Authors : Xu, Q.S.; Shin, D.H.; Pufan, R.; Yokota, H.; Kim, R.; Kim, S.H.; Berkeley Structural Genomics Center (BSGC)  
Deposited on : 2003-10-10  
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	337	
1	B	337	

## 2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 5034 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called putative phosphotransacetylase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	329	2502	1582	424	485	1	10	0	0	0
1	B	329	2502	1582	424	485	1	10	0	0	0

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	GLY	-	CLONING ARTIFACT	UNP Q99ZQ5
A	-4	GLY	-	CLONING ARTIFACT	UNP Q99ZQ5
A	-3	GLY	-	CLONING ARTIFACT	UNP Q99ZQ5
A	-2	GLY	-	CLONING ARTIFACT	UNP Q99ZQ5
A	-1	GLY	-	CLONING ARTIFACT	UNP Q99ZQ5
A	0	GLY	-	CLONING ARTIFACT	UNP Q99ZQ5
A	1	MSE	MET	MODIFIED RESIDUE	UNP Q99ZQ5
A	19	MSE	MET	MODIFIED RESIDUE	UNP Q99ZQ5
A	83	MSE	MET	MODIFIED RESIDUE	UNP Q99ZQ5
A	103	MSE	MET	MODIFIED RESIDUE	UNP Q99ZQ5
A	113	MSE	MET	MODIFIED RESIDUE	UNP Q99ZQ5
A	117	MSE	MET	MODIFIED RESIDUE	UNP Q99ZQ5
A	123	MSE	MET	MODIFIED RESIDUE	UNP Q99ZQ5
A	157	MSE	MET	MODIFIED RESIDUE	UNP Q99ZQ5
A	204	MSE	MET	MODIFIED RESIDUE	UNP Q99ZQ5
A	288	MSE	MET	MODIFIED RESIDUE	UNP Q99ZQ5
B	-5	GLY	-	CLONING ARTIFACT	UNP Q99ZQ5
B	-4	GLY	-	CLONING ARTIFACT	UNP Q99ZQ5
B	-3	GLY	-	CLONING ARTIFACT	UNP Q99ZQ5
B	-2	GLY	-	CLONING ARTIFACT	UNP Q99ZQ5
B	-1	GLY	-	CLONING ARTIFACT	UNP Q99ZQ5
B	0	GLY	-	CLONING ARTIFACT	UNP Q99ZQ5
B	1	MSE	MET	MODIFIED RESIDUE	UNP Q99ZQ5
B	19	MSE	MET	MODIFIED RESIDUE	UNP Q99ZQ5
B	83	MSE	MET	MODIFIED RESIDUE	UNP Q99ZQ5

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	103	MSE	MET	MODIFIED RESIDUE	UNP Q99ZQ5
B	113	MSE	MET	MODIFIED RESIDUE	UNP Q99ZQ5
B	117	MSE	MET	MODIFIED RESIDUE	UNP Q99ZQ5
B	123	MSE	MET	MODIFIED RESIDUE	UNP Q99ZQ5
B	157	MSE	MET	MODIFIED RESIDUE	UNP Q99ZQ5
B	204	MSE	MET	MODIFIED RESIDUE	UNP Q99ZQ5
B	288	MSE	MET	MODIFIED RESIDUE	UNP Q99ZQ5


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	12	Total O 12 12	0	0
2	B	18	Total O 18 18	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: putative phosphotransacetylase



I327	E328	S329	GLN	GLY
------	------	------	-----	-----

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	173.72Å 173.72Å 173.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.93 – 2.70 19.93 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (19.93-2.70) 100.0 (19.93-2.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.12	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.57 (at 2.71Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.229 , 0.281 0.227 , 0.278	Depositor DCC
$R_{free}$ test set	2537 reflections (10.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	67.7	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 56.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5034	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	72.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.28% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.62	5/2529 (0.2%)	0.78	4/3400 (0.1%)
1	B	0.63	3/2529 (0.1%)	0.79	3/3400 (0.1%)
All	All	0.63	8/5058 (0.2%)	0.78	7/6800 (0.1%)

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	92	LYS	CE-NZ	7.80	1.68	1.49
1	A	96	THR	CB-CG2	6.92	1.75	1.52
1	A	94	LYS	CB-CG	6.36	1.69	1.52
1	B	92	LYS	CE-NZ	6.35	1.65	1.49
1	B	94	LYS	CD-CE	5.98	1.66	1.51

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	169	ALA	N-CA-C	5.92	126.98	111.00
1	A	239	GLY	N-CA-C	5.75	127.49	113.10
1	B	169	ALA	N-CA-C	5.65	126.25	111.00
1	A	201	LYS	N-CA-C	-5.36	96.54	111.00
1	B	201	LYS	N-CA-C	-5.33	96.62	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2502	0	2542	216	0
1	B	2502	0	2542	180	0
2	A	12	0	0	1	0
2	B	18	0	0	1	0
All	All	5034	0	5084	393	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 39.

The worst 5 of 393 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:96:THR:CG2	1:A:96:THR:CB	1.75	1.63
1:A:92:LYS:NZ	1:A:92:LYS:CE	1.68	1.56
1:A:169:ALA:CB	1:A:269:VAL:HA	1.71	1.19
1:B:169:ALA:CB	1:B:269:VAL:HA	1.74	1.18
1:A:92:LYS:N	1:A:92:LYS:HD2	1.58	1.15

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	327/337 (97%)	284 (87%)	32 (10%)	11 (3%)	<b>3</b> <b>8</b>
1	B	327/337 (97%)	284 (87%)	32 (10%)	11 (3%)	<b>3</b> <b>8</b>
All	All	654/674 (97%)	568 (87%)	64 (10%)	22 (3%)	<b>3</b> <b>8</b>

5 of 22 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	92	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	293	GLY
1	B	293	GLY
1	B	78	ALA
1	B	92	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	265/256 (104%)	239 (90%)	26 (10%)	8 18
1	B	265/256 (104%)	239 (90%)	26 (10%)	8 18
All	All	530/512 (104%)	478 (90%)	52 (10%)	8 18

5 of 52 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	285	ARG
1	B	58	LEU
1	B	285	ARG
1	A	308	ARG
1	B	1	MSE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	266	ASN
1	B	57	ASN
1	B	264	GLN
1	A	264	GLN
1	B	232	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

#### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

#### 5.7 Other polymers [i](#)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	319/337 (94%)	0.17	18 (5%) 24 23	42, 74, 121, 138	0
1	B	319/337 (94%)	-0.14	6 (1%) 66 69	40, 61, 106, 131	0
All	All	638/674 (94%)	0.02	24 (3%) 40 39	40, 67, 116, 138	0

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	95	ALA	5.6
1	A	105	ARG	5.1
1	A	93	GLY	3.8
1	A	146	PRO	3.5
1	A	51	GLN	3.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.