



# wwPDB X-ray Structure Validation Summary Report

Jan 27, 2024 – 11:50 AM EST

PDB ID : 1AFC  
Title : STRUCTURAL STUDIES OF THE BINDING OF THE ANTI-ULCER  
DRUG SUCROSE OCTASULFATE TO ACIDIC FIBROBLAST GROWTH  
FACTOR  
Authors : Zhu, X.; Hsu, B.T.; Rees, D.C.  
Deposited on : 1993-07-13  
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	140	10% 39% 29% 13% 9%
1	B	140	11% 39% 31% 10% 9%
1	C	140	13% 34% 33% 11% 9%
1	D	140	16% 33% 30% 11% 9%
1	E	140	16% 29% 34% 12% 9%
1	F	140	13% 34% 30% 14% 9%
1	G	140	13% 36% 31% 11% 9%
1	H	140	11% 33% 28% 19% 9%

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	I	2	 100%
2	J	2	 100%
2	K	2	 100%
2	L	2	 100%
2	M	2	 100%
2	N	2	 100%
2	O	2	 100%
2	P	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GU4	I	1	-	-	X	-
2	YYJ	I	2	-	-	X	-
2	GU4	K	1	-	-	X	-
2	YYJ	K	2	-	-	X	-
2	YYJ	L	2	-	-	X	-
2	GU4	M	1	-	-	X	-
2	YYJ	M	2	-	-	X	-
2	YYJ	N	2	-	-	X	-
2	GU4	O	1	-	-	X	-
2	YYJ	O	2	-	-	X	-
2	GU4	P	1	-	-	X	-
2	YYJ	P	2	-	-	X	-

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 8304 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ACIDIC FIBROBLAST GROWTH FACTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	127	983	630	168	182	3	0	0	0
1	B	127	983	630	168	182	3	0	0	0
1	C	127	983	630	168	182	3	0	0	0
1	D	127	983	630	168	182	3	0	0	0
1	E	127	983	630	168	182	3	0	0	0
1	F	127	983	630	168	182	3	0	0	0
1	G	127	983	630	168	182	3	0	0	0
1	H	127	983	630	168	182	3	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	47	ALA	CYS	conflict	UNP P03968
B	47	ALA	CYS	conflict	UNP P03968
C	47	ALA	CYS	conflict	UNP P03968
D	47	ALA	CYS	conflict	UNP P03968
E	47	ALA	CYS	conflict	UNP P03968
F	47	ALA	CYS	conflict	UNP P03968
G	47	ALA	CYS	conflict	UNP P03968
H	47	ALA	CYS	conflict	UNP P03968

- Molecule 2 is an oligosaccharide called 1,3,4,6-tetra-O-sulfo-beta-D-fructofuranose-(2-1)-2,3,4,6-tetra-O-sulfonato-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	I	2	Total	C	O	S	0	0	0
			55	12	35	8			
2	J	2	Total	C	O	S	0	0	0
			55	12	35	8			
2	K	2	Total	C	O	S	0	0	0
			55	12	35	8			
2	L	2	Total	C	O	S	0	0	0
			55	12	35	8			
2	M	2	Total	C	O	S	0	0	0
			55	12	35	8			
2	N	2	Total	C	O	S	0	0	0
			55	12	35	8			
2	O	2	Total	C	O	S	0	0	0
			55	12	35	8			
2	P	2	Total	C	O	S	0	0	0
			55	12	35	8			


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: ACIDIC FIBROBLAST GROWTH FACTOR

Chain A: 




- Molecule 1: ACIDIC FIBROBLAST GROWTH FACTOR

Chain B: 




- Molecule 1: ACIDIC FIBROBLAST GROWTH FACTOR

Chain C: 




- Molecule 1: ACIDIC FIBROBLAST GROWTH FACTOR

Chain D:  16% 33% 30% 11% 9%




• Molecule 1: ACIDIC FIBROBLAST GROWTH FACTOR

Chain E:  16% 29% 34% 12% 9%




• Molecule 1: ACIDIC FIBROBLAST GROWTH FACTOR

Chain F:  13% 34% 30% 14% 9%




• Molecule 1: ACIDIC FIBROBLAST GROWTH FACTOR

Chain G:  13% 36% 31% 11% 9%



- Molecule 1: ACIDIC FIBROBLAST GROWTH FACTOR

Chain H: 



- Molecule 2: 1,3,4,6-tetra-O-sulfo-beta-D-fructofuranose-(2-1)-2,3,4,6-tetra-O-sulfonato-alpha-D-glucopyranose

Chain I: 

G101  
Y122

- Molecule 2: 1,3,4,6-tetra-O-sulfo-beta-D-fructofuranose-(2-1)-2,3,4,6-tetra-O-sulfonato-alpha-D-glucopyranose

Chain J: 

G101  
Y122

- Molecule 2: 1,3,4,6-tetra-O-sulfo-beta-D-fructofuranose-(2-1)-2,3,4,6-tetra-O-sulfonato-alpha-D-glucopyranose

Chain K: 

G101  
Y122

- Molecule 2: 1,3,4,6-tetra-O-sulfo-beta-D-fructofuranose-(2-1)-2,3,4,6-tetra-O-sulfonato-alpha-D-glucopyranose

Chain L: 

G101  
Y122

- Molecule 2: 1,3,4,6-tetra-O-sulfo-beta-D-fructofuranose-(2-1)-2,3,4,6-tetra-O-sulfonato-alpha-D-glucopyranose

Chain M: 

G101  
Y122

- Molecule 2: 1,3,4,6-tetra-O-sulfo-beta-D-fructofuranose-(2-1)-2,3,4,6-tetra-O-sulfonato-alpha-D-glucopyranose

Chain N:  100%

CU01  
YJ2

- Molecule 2: 1,3,4,6-tetra-O-sulfo-beta-D-fructofuranose-(2-1)-2,3,4,6-tetra-O-sulfonato-alpha-D-glucopyranose

Chain O:  100%

CU01  
YJ2

- Molecule 2: 1,3,4,6-tetra-O-sulfo-beta-D-fructofuranose-(2-1)-2,3,4,6-tetra-O-sulfonato-alpha-D-glucopyranose

Chain P:  100%

CU01  
YJ2

## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	110.60Å 110.60Å 172.20Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	(Not available) – 2.70	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-2.70)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, $R_{free}$	0.204 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	8304	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	13.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: YYJ, GU4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.87	0/1007	1.54	8/1363 (0.6%)
1	B	0.91	0/1007	1.48	8/1363 (0.6%)
1	C	0.89	0/1007	1.50	10/1363 (0.7%)
1	D	0.89	0/1007	1.49	11/1363 (0.8%)
1	E	0.91	0/1007	1.49	9/1363 (0.7%)
1	F	0.90	0/1007	1.52	8/1363 (0.6%)
1	G	0.89	0/1007	1.46	4/1363 (0.3%)
1	H	0.90	0/1007	1.51	8/1363 (0.6%)
All	All	0.89	0/8056	1.50	66/10904 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	89
1	B	0	77
1	C	0	76
1	D	0	76
1	E	0	74
1	F	0	78
1	G	0	67
1	H	0	83
All	All	0	620

There are no bond length outliers.

The worst 5 of 66 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	107	TRP	CD1-CG-CD2	11.01	115.11	106.30
1	A	24	ARG	NE-CZ-NH2	-10.65	114.97	120.30
1	D	37	ARG	NE-CZ-NH2	-10.06	115.27	120.30
1	E	24	ARG	NE-CZ-NH2	-9.73	115.43	120.30
1	D	24	ARG	NE-CZ-NH2	-9.52	115.54	120.30

There are no chirality outliers.

5 of 620 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	10	LYS	Mainchain
1	A	11	PRO	Mainchain
1	A	12	LYS	Mainchain
1	A	14	LEU	Mainchain
1	A	15	TYR	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	983	0	944	109	0
1	B	983	0	944	95	0
1	C	983	0	944	112	0
1	D	983	0	944	96	0
1	E	983	0	944	86	0
1	F	983	0	944	84	0
1	G	983	0	944	122	0
1	H	983	0	944	121	0
2	I	55	0	6	24	0
2	J	55	0	6	7	0
2	K	55	0	6	13	0
2	L	55	0	6	21	0
2	M	55	0	6	19	0
2	N	55	0	6	14	0
2	O	55	0	6	27	0
2	P	55	0	6	18	0
All	All	8304	0	7600	899	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 57.

The worst 5 of 899 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:114:ASN:HD21	1:A:116:ARG:NH1	1.43	1.16
1:G:24:ARG:HH11	1:G:26:LEU:HD11	1.01	1.14
1:H:27:PRO:HB3	1:H:61:THR:HG21	1.31	1.12
1:D:86:LEU:HD21	1:D:100:LYS:HG3	1.30	1.11
1:F:24:ARG:HD2	1:F:26:LEU:HD11	1.30	1.11

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	125/140 (89%)	104 (83%)	17 (14%)	4 (3%)	4	9
1	B	125/140 (89%)	112 (90%)	9 (7%)	4 (3%)	4	9
1	C	125/140 (89%)	103 (82%)	18 (14%)	4 (3%)	4	9
1	D	125/140 (89%)	104 (83%)	16 (13%)	5 (4%)	3	6
1	E	125/140 (89%)	102 (82%)	16 (13%)	7 (6%)	2	3
1	F	125/140 (89%)	105 (84%)	11 (9%)	9 (7%)	1	1
1	G	125/140 (89%)	105 (84%)	15 (12%)	5 (4%)	3	6
1	H	125/140 (89%)	92 (74%)	19 (15%)	14 (11%)	0	0
All	All	1000/1120 (89%)	827 (83%)	121 (12%)	52 (5%)	2	3

5 of 52 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	18	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	81	GLU
1	B	18	ASN
1	B	68	ASP
1	D	18	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	101/122 (83%)	69 (68%)	32 (32%)	0	0
1	B	101/122 (83%)	68 (67%)	33 (33%)	0	0
1	C	101/122 (83%)	67 (66%)	34 (34%)	0	0
1	D	101/122 (83%)	74 (73%)	27 (27%)	0	1
1	E	101/122 (83%)	69 (68%)	32 (32%)	0	0
1	F	101/122 (83%)	71 (70%)	30 (30%)	0	1
1	G	101/122 (83%)	64 (63%)	37 (37%)	0	0
1	H	101/122 (83%)	71 (70%)	30 (30%)	0	1
All	All	808/976 (83%)	553 (68%)	255 (32%)	0	0

5 of 255 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	84	LEU
1	G	122	ARG
1	E	80	ASN
1	G	116	ARG
1	H	58	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 37 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	95	ASN

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type
1	H	95	ASN
1	G	106	HIS
1	H	43	GLN
1	C	93	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

16 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GU4	I	1	2	27,27,28	1.49	4 (14%)	29,43,45	2.28	7 (24%)
2	YYJ	I	2	2	27,28,28	2.09	6 (22%)	28,46,46	1.59	3 (10%)
2	GU4	J	1	2	27,27,28	1.47	5 (18%)	29,43,45	2.07	7 (24%)
2	YYJ	J	2	2	27,28,28	1.78	5 (18%)	28,46,46	0.95	1 (3%)
2	GU4	K	1	2	27,27,28	1.51	4 (14%)	29,43,45	2.24	8 (27%)
2	YYJ	K	2	2	27,28,28	1.80	4 (14%)	28,46,46	1.53	5 (17%)
2	GU4	L	1	2	27,27,28	1.93	5 (18%)	29,43,45	1.55	2 (6%)
2	YYJ	L	2	2	27,28,28	1.45	4 (14%)	28,46,46	1.23	3 (10%)
2	GU4	M	1	2	27,27,28	1.66	5 (18%)	29,43,45	2.70	10 (34%)
2	YYJ	M	2	2	27,28,28	1.77	4 (14%)	28,46,46	0.99	1 (3%)
2	GU4	N	1	2	27,27,28	1.54	5 (18%)	29,43,45	1.47	3 (10%)
2	YYJ	N	2	2	27,28,28	1.54	4 (14%)	28,46,46	1.39	5 (17%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GU4	O	1	2	27,27,28	1.47	4 (14%)	29,43,45	2.28	8 (27%)
2	YYJ	O	2	2	27,28,28	1.43	4 (14%)	28,46,46	1.34	2 (7%)
2	GU4	P	1	2	27,27,28	1.87	5 (18%)	29,43,45	1.56	4 (13%)
2	YYJ	P	2	2	27,28,28	1.74	5 (18%)	28,46,46	1.52	3 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GU4	I	1	2	-	9/21/38/41	0/1/1/1
2	YYJ	I	2	2	-	6/23/42/42	0/1/1/1
2	GU4	J	1	2	-	10/21/38/41	0/1/1/1
2	YYJ	J	2	2	-	14/23/42/42	0/1/1/1
2	GU4	K	1	2	-	10/21/38/41	0/1/1/1
2	YYJ	K	2	2	-	6/23/42/42	0/1/1/1
2	GU4	L	1	2	-	13/21/38/41	0/1/1/1
2	YYJ	L	2	2	-	7/23/42/42	0/1/1/1
2	GU4	M	1	2	-	13/21/38/41	0/1/1/1
2	YYJ	M	2	2	-	15/23/42/42	0/1/1/1
2	GU4	N	1	2	-	9/21/38/41	0/1/1/1
2	YYJ	N	2	2	-	12/23/42/42	0/1/1/1
2	GU4	O	1	2	-	12/21/38/41	0/1/1/1
2	YYJ	O	2	2	-	13/23/42/42	0/1/1/1
2	GU4	P	1	2	-	9/21/38/41	0/1/1/1
2	YYJ	P	2	2	-	8/23/42/42	0/1/1/1

The worst 5 of 73 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	I	2	YYJ	O3-S3	-6.04	1.39	1.57
2	J	2	YYJ	O3-S3	-4.96	1.42	1.57
2	P	1	GU4	O4-S4	-4.92	1.42	1.57
2	L	1	GU4	O6-S6	-4.90	1.43	1.56
2	P	1	GU4	O2-S2	-4.77	1.43	1.57

The worst 5 of 72 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	K	1	GU4	C1-O5-C5	-8.06	101.27	112.19
2	M	1	GU4	C2-O2-S2	7.14	127.22	117.91
2	I	1	GU4	O2-C2-C3	6.87	114.26	106.65
2	J	1	GU4	C2-O2-S2	6.38	126.23	117.91
2	O	1	GU4	C4-O4-S4	6.30	131.05	118.88

There are no chirality outliers.

5 of 166 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	I	1	GU4	C4-C5-C6-O6
2	I	1	GU4	C3-O3-S3-O29
2	I	1	GU4	C2-O2-S2-O10
2	I	2	YYJ	C2-C1-O1-S1
2	I	2	YYJ	C6-O6-S6-O1S6


There are no ring outliers.


16 monomers are involved in 143 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	L	1	GU4	8	0
2	I	1	GU4	15	0
2	K	2	YYJ	9	0
2	P	1	GU4	13	0
2	O	2	YYJ	22	0
2	K	1	GU4	9	0
2	N	1	GU4	4	0
2	O	1	GU4	13	0
2	P	2	YYJ	11	0
2	N	2	YYJ	13	0
2	J	1	GU4	2	0
2	M	2	YYJ	14	0
2	L	2	YYJ	15	0
2	J	2	YYJ	6	0
2	I	2	YYJ	21	0
2	M	1	GU4	9	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.




















## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.