Full wwPDB X-ray Structure Validation Report (i)

May 22, 2020 – 12:39 am BST

PDB ID	:	4AFL
Title	:	The crystal structure of the ING4 dimerization domain reveals the functional organization of the ING family of chromatin binding proteins.
Authors	:	Culurgioni, S.; Munoz, I.G.; Moreno, A.; Palacios, A.; Villate, M.; Palmero, I.; Montoya, G.; Blanco, F.J.
Deposited on	:	2012-01-19
Resolution	:	2.27 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6980 (2.30-2.26)
Clashscore	141614	7711 (2.30-2.26)
Ramachandran outliers	138981	7597 (2.30-2.26)
Sidechain outliers	138945	7598 (2.30-2.26)
RSRZ outliers	127900	6849 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	104	88% 9% • •
1	B	104	80% 14% 6%
1	C	104	% 85% 13% • •
1	D	104	% 78% 13% • 9%
1	E	104	87% 9% • •
1	F	104	2% 74% 12% • 13%

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 4887 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called INHIBITOR OF GROWTH PROTEIN 4.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	102	816	511	135	164	1	5	0	0	0
1	B	98	777	490	131	150	1	5	0	0	0
1	C	102	837	522	142	167	1	5	0	1	0
1	D	95	754	474	127	147	1	5	0	0	0
1	E	101	800	503	135	156	1	5	0	0	0
1	F	91	717	446	120	146	1	4	0	0	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	28	28	28	0	0
2	B	30	30	30	0	0
2	C	43	43	43	0	0
2	D	30	30	30	0	0
2	E	32	32	32	0	0
2	F	23	23	23	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: INHIBITOR OF GROWTH PROTEIN 4

Chain A:



- Molecule 1: INHIBITOR OF GROWTH PROTEIN 4

Chain B:



- Molecule 1: INHIBITOR OF GROWTH PROTEIN 4

Chain C:



- Molecule 1: INHIBITOR OF GROWTH PROTEIN 4

Chain D:



- Molecule 1: INHIBITOR OF GROWTH PROTEIN 4

Chain E:



- Molecule 1: INHIBITOR OF GROWTH PROTEIN 4

Chain F:


ALA
ALA
GLY
MSE
TYR
LEU
GLU
HIS
Y10
L11
Q22
Q26
D32
E36
A46
M50
S51
S52
ALA
ARG
SER
L56
S57
E60
L64
I68
C75
M87
D103
LEU
ALA

4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants a, b, c, α, β, γ	128.45Å 186.60Å 62.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.69 – 2.27 46.65 – 2.28	Depositor EDS
% Data completeness (in resolution range)	98.5 (44.69-2.27) 98.5 (46.65-2.28)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.21 (at 2.27Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.205 , 0.256 0.206 , 0.251	Depositor DCC
R_{free} test set	1735 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å^2)	35.9	Xtriage
Anisotropy	0.107	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.33 , 37.9	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	4887	wwPDB-VP
Average B, all atoms (Å^2)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.48% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.40	0/821	0.52	0/1095
1	B	0.39	0/782	0.47	0/1041
1	C	0.42	0/842	0.55	0/1119
1	D	0.42	0/759	0.50	0/1012
1	E	0.37	0/805	0.48	0/1073
1	F	0.39	0/719	0.52	0/957
All	All	0.40	0/4728	0.50	0/6297

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	816	0	781	6	0
1	B	777	0	749	8	0
1	C	837	0	815	9	0
1	D	754	0	708	7	0
1	E	800	0	760	10	0
1	F	717	0	673	10	0
2	A	28	0	0	1	0
2	B	30	0	0	0	0
2	C	43	0	0	1	0
2	D	30	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	32	0	0	0	0
2	F	23	0	0	1	0
All	All	4887	0	4486	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:18:PRO:HD3	1:C:49:TYR:HE2	1.55	0.72
1:F:75:CYS:SG	2:F:2020:HOH:O	2.50	0.69
1:B:18:PRO:HB3	1:F:50:MSE:HG2	1.75	0.67
1:A:94:ASP:OD1	1:C:69:GLN:NE2	2.29	0.66
1:D:48:GLU:HG2	1:D:64:LEU:HD21	1.84	0.59
1:A:76:LYS:NZ	2:A:2021:HOH:O	2.36	0.58
1:D:87:MSE:O	1:D:91:GLU:HG2	2.04	0.58
1:D:50:MSE:HE1	1:E:19:PHE:CD1	2.40	0.57
1:F:22:GLN:O	1:F:26:GLN:HG2	2.05	0.56
1:F:64:LEU:O	1:F:68:ILE:HG13	2.07	0.53
1:F:46:ALA:HB1	1:F:50:MSE:HE3	1.91	0.52
1:B:62:LEU:HD11	1:B:66:LYS:HE2	1.93	0.50
1:C:34:ARG:HD3	1:C:78:PHE:CZ	2.47	0.50
1:C:22:GLN:O	1:C:26:GLN:HG2	2.13	0.48
1:E:102:THR:HA	1:F:87:MSE:SE	2.64	0.47
1:D:69:GLN:NE2	1:E:94:ASP:OD1	2.36	0.47
1:C:49:TYR:HE1	1:C:61:LYS:HG3	1.80	0.47
1:F:103:ASP:OD1	1:F:103:ASP:N	2.45	0.47
1:B:13:SER:HB3	1:B:15:GLU:OE2	2.15	0.46
1:C:34:ARG:NH1	2:C:2013:HOH:O	2.47	0.46
1:F:32:ASP:O	1:F:36:GLU:HB2	2.15	0.46
1:B:56:LEU:HD22	1:B:60:GLU:HB3	1.99	0.45
1:D:18:PRO:HB3	1:E:50:MSE:HB3	1.98	0.45
1:E:13:SER:HB3	1:E:15:GLU:OE2	2.15	0.45
1:A:98:ARG:O	1:A:102:THR:HG23	2.17	0.45
1:D:49:TYR:CE2	1:E:18:PRO:HG3	2.51	0.44
1:A:89:THR:HG21	1:C:72:TYR:CZ	2.52	0.44
1:B:5:MSE:HB2	1:E:5:MSE:HG3	2.00	0.43
1:B:52:SER:O	1:B:54:ARG:N	2.49	0.43
1:A:49:TYR:OH	1:A:61:LYS:HE3	2.20	0.42
1:B:20:GLU:HG2	1:B:92:MSE:SE	2.70	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:5:MSE:HE3	1:E:5:MSE:HB3	1.85	0.42
1:F:64:LEU:HD12	1:F:64:LEU:HA	1.90	0.41
1:B:22:GLN:O	1:B:26:GLN:HG2	2.21	0.41
1:E:64:LEU:HA	1:E:64:LEU:HD22	1.82	0.41
1:C:88[B]:GLN:HG3	1:C:89:THR:N	2.36	0.41
1:E:15:GLU:H	1:E:15:GLU:CD	2.25	0.40
1:D:22:GLN:O	1:D:26:GLN:HG2	2.20	0.40
1:F:56:LEU:O	1:F:57:SER:HB3	2.21	0.40
1:C:56:LEU:HD12	1:C:56:LEU:HA	1.96	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	100/104 (96%)	100 (100%)	0	0	100	100
1	B	96/104 (92%)	93 (97%)	3 (3%)	0	100	100
1	C	101/104 (97%)	100 (99%)	1 (1%)	0	100	100
1	D	91/104 (88%)	89 (98%)	2 (2%)	0	100	100
1	E	99/104 (95%)	97 (98%)	2 (2%)	0	100	100
1	F	87/104 (84%)	84 (97%)	2 (2%)	1 (1%)	14	14
All	All	574/624 (92%)	563 (98%)	10 (2%)	1 (0%)	47	57

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	57	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	86/87 (99%)	83 (96%)	3 (4%)	36	48
1	B	80/87 (92%)	79 (99%)	1 (1%)	69	80
1	C	90/87 (103%)	86 (96%)	4 (4%)	28	37
1	D	77/87 (88%)	72 (94%)	5 (6%)	17	21
1	E	81/87 (93%)	77 (95%)	4 (5%)	25	33
1	F	74/87 (85%)	72 (97%)	2 (3%)	44	59
All	All	488/522 (94%)	469 (96%)	19 (4%)	32	43

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	64	LEU
1	A	76	LYS
1	A	92	MSE
1	B	34	ARG
1	C	27	LEU
1	C	56	LEU
1	C	76	LYS
1	C	98	ARG
1	D	5	MSE
1	D	12	ASP
1	D	76	LYS
1	D	91	GLU
1	D	101	ASP
1	E	5	MSE
1	E	64	LEU
1	E	76	LYS
1	E	88	GLN
1	F	11	LEU
1	F	103	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	96	HIS
1	C	24	ASN
1	C	69	GLN
1	F	69	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	97/104 (93%)	-0.07	0	100	100	27, 41, 59, 68	0
1	B	93/104 (89%)	0.06	0	100	100	28, 47, 79, 88	0
1	C	97/104 (93%)	-0.02	1 (1%)	82	86	24, 38, 58, 74	0
1	D	90/104 (86%)	0.00	1 (1%)	80	84	25, 46, 89, 101	0
1	E	96/104 (92%)	-0.11	0	100	100	28, 44, 77, 90	0
1	F	87/104 (83%)	-0.03	2 (2%)	60	66	30, 45, 76, 85	0
All	All	560/624 (89%)	-0.03	4 (0%)	87	90	24, 43, 75, 101	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	56	LEU	4.2
1	F	11	LEU	3.0
1	C	105	ALA	2.7
1	F	60	GLU	2.6

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.