



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 20, 2024 – 09:51 pm GMT

PDB ID : 7ASX
Title : Fixed-target serial femtosecond crystallography using in cellulo grown *Neurospora crassa* HEX-1 microcrystals. (Chip 1)
Authors : Lahey-Rudolph, J.M.; Schoenherr, R.; Barthelmess, M.; Fischer, P.; Seuring, C.; Wagner, A.; Meents, A.; Redecke, L.
Deposited on : 2020-10-28
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	177	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2375 atoms, of which 1136 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called eIF-5a domain-containing protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	141	2247	698	1136	200	206	7	30	3	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	177	ALA	-	cloning artifact	UNP A0A0B0EDT5


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	128	128	128	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: eIF-5a domain-containing protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	58.75Å 58.75Å 192.83Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.73 – 1.80 30.73 – 1.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (30.73-1.80) 100.0 (30.73-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.42 (at 1.80Å)	Xtrriage
Refinement program	PHENIX v1.16-3546	Depositor
R, R_{free}	0.226 , 0.273 0.226 , 0.273	Depositor DCC
R_{free} test set	1541 reflections (8.02%)	wwPDB-VP
Wilson B-factor (Å ²)	17.3	Xtrriage
Anisotropy	0.065	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 60.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.31$, $\langle L^2 \rangle = 0.15$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	2375	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/1132	0.48	0/1535

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1111	1136	1137	2	0
2	A	128	0	0	1	0
All	All	1239	1136	1137	2	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (2) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:86:VAL:HG22	1:A:96:GLN:HG2	1.94	0.49
1:A:104:LYS:NZ	2:A:205:HOH:O	2.47	0.47

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	142/177 (80%)	138 (97%)	4 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	128/148 (86%)	128 (100%)	0	100 100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	141/177 (79%)	3.36	118 (83%) 0 0	4, 16, 44, 61	3 (2%)

All (118) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	170	VAL	8.2
1	A	160	HIS	8.0
1	A	171	VAL	7.8
1	A	113	ASP	7.6
1	A	157[A]	VAL	6.6
1	A	146	GLU	6.3
1	A	114	GLY	6.1
1	A	89	PRO	5.8
1	A	147	SER	5.7
1	A	138	TRP	5.6
1	A	91	PRO	5.3
1	A	112	GLN	5.2
1	A	142	GLN	5.2
1	A	161	GLY	5.0
1	A	159	ASP	4.9
1	A	145	PHE	4.8
1	A	139	ASN	4.7
1	A	53	CYS	4.7
1	A	165	ALA	4.6
1	A	144	ALA	4.5
1	A	111	MET	4.5
1	A	155	LEU	4.4
1	A	47	ILE	4.4
1	A	137	LEU	4.4
1	A	93	VAL	4.4
1	A	33	VAL	4.3
1	A	55	VAL	4.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	130	PRO	4.3
1	A	158[A]	SER	4.2
1	A	154	VAL	4.2
1	A	95	VAL	4.2
1	A	31	GLN	4.1
1	A	152	VAL	4.1
1	A	162	ARG	4.1
1	A	156	VAL	4.1
1	A	108	VAL	4.0
1	A	166	VAL	4.0
1	A	37	CYS	4.0
1	A	52	PRO	3.9
1	A	90	ALA	3.9
1	A	116	ILE	3.9
1	A	64	THR	3.9
1	A	117	VAL	3.9
1	A	85	PHE	3.8
1	A	79	LEU	3.8
1	A	62	ALA	3.8
1	A	136	SER	3.8
1	A	36	PRO	3.8
1	A	101	PRO	3.8
1	A	148	GLY	3.8
1	A	84	SER	3.7
1	A	72	VAL	3.7
1	A	143	LYS	3.7
1	A	106	TYR	3.7
1	A	88	ASN	3.6
1	A	122	THR	3.6
1	A	103	PHE	3.6
1	A	115	SER	3.6
1	A	35[A]	ILE	3.6
1	A	63	ALA	3.6
1	A	56	ILE	3.5
1	A	69	TYR	3.4
1	A	70	LEU	3.3
1	A	125	VAL	3.3
1	A	132	ILE	3.3
1	A	102	VAL	3.3
1	A	151	SER	3.3
1	A	131	VAL	3.3
1	A	48	LEU	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	120	THR	3.2
1	A	99	LEU	3.2
1	A	32	THR	3.2
1	A	87	SER	3.1
1	A	149	ARG	3.1
1	A	140	ARG	3.1
1	A	94	VAL	3.1
1	A	141	LEU	3.1
1	A	168	MET	3.1
1	A	128	ASN	3.1
1	A	107	ARG	3.0
1	A	75	PHE	3.0
1	A	42	LEU	3.0
1	A	74	LEU	3.0
1	A	118	ALA	3.0
1	A	40	ILE	2.9
1	A	34	THR	2.9
1	A	76	THR	2.9
1	A	46	LEU	2.9
1	A	92	SER	2.9
1	A	59	SER	2.8
1	A	121	GLU	2.8
1	A	58	ILE	2.8
1	A	98	MET	2.8
1	A	129	LEU	2.8
1	A	97	THR	2.7
1	A	45	ILE	2.7
1	A	133	ASP	2.7
1	A	86	VAL	2.7
1	A	109	LEU	2.6
1	A	51	ARG	2.6
1	A	135	SER	2.6
1	A	54	GLN	2.5
1	A	41	ARG	2.5
1	A	71	GLY	2.5
1	A	164	MET	2.5
1	A	134	GLN	2.4
1	A	83	SER	2.4
1	A	119	MET	2.4
1	A	169	LYS	2.3
1	A	60	THR	2.3
1	A	167	ASP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	78	GLN	2.3
1	A	67	HIS	2.3
1	A	65	GLY	2.2
1	A	153	ARG	2.2
1	A	73	ASP	2.2
1	A	61	SER	2.1
1	A	44	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.