



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 22, 2020 – 06:08 am BST

PDB ID : 5C0N  
Title : Development of a monoclonal antibody targeting secreted aP2 to treat diabetes and fatty liver disease  
Authors : Doyle, C.  
Deposited on : 2015-06-12  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	132	 77% 22%
1	B	132	 73% 23%
2	C	222	 74% 21%
2	H	222	 74% 20%
3	D	219	 75% 21%
3	L	219	 75% 21%

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8540 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fatty acid-binding protein, adipocyte.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	131	1017	639	174	198	6	0	0	0
1	B	131	1017	639	174	198	6	0	0	0

- Molecule 2 is a protein called Fab CA33 Heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	214	1585	1009	253	316	7	0	0	0
2	C	216	1599	1016	256	320	7	0	0	0

- Molecule 3 is a protein called Fab CA33 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	L	216	1661	1039	272	341	9	0	0	0
3	D	216	1661	1039	272	341	9	0	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fatty acid-binding protein, adipocyte




- Molecule 1: Fatty acid-binding protein, adipocyte



- Molecule 2: Fab CA33 Heavy chain



- Molecule 2: Fab CA33 Heavy chain



- Molecule 3: Fab CA33 light chain





- Molecule 3: Fab CA33 light chain

Chain D: 75% 21% ..



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	65.27Å 101.95Å 95.32Å 90.00° 90.03° 90.00°	Depositor
Resolution (Å)	54.97 – 3.00 54.97 – 2.95	Depositor EDS
% Data completeness (in resolution range)	98.9 (54.97-3.00) 98.8 (54.97-2.95)	Depositor EDS
$R_{merge}$	0.18	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.91 (at 2.96Å)	Xtrriage
Refinement program	REFMAC 5.8.0049	Depositor
R, $R_{free}$	0.186 , 0.265 0.196 , 0.197	Depositor DCC
$R_{free}$ test set	1333 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	57.5	Xtrriage
Anisotropy	0.445	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 41.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.039 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	8540	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	61.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.16% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.68	0/1029	0.78	0/1380
1	B	0.56	0/1029	0.79	0/1380
2	C	0.70	0/1642	0.84	1/2253 (0.0%)
2	H	0.68	0/1629	0.79	1/2236 (0.0%)
3	D	0.64	0/1700	0.81	0/2313
3	L	0.65	0/1700	0.79	0/2313
All	All	0.66	0/8729	0.80	2/11875 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	65	ARG	NE-CZ-NH2	-5.70	117.45	120.30
2	H	147	CYS	CA-CB-SG	-5.55	104.02	114.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1017	0	1040	10	0
1	B	1017	0	1040	16	0
2	C	1599	0	1568	27	0
2	H	1585	0	1555	23	0
3	D	1661	0	1580	25	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	L	1661	0	1580	28	0
All	All	8540	0	8363	119	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 119 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:194:HIS:O	3:L:216:ARG:NH1	1.87	1.06
2:C:98:ASP:OD1	3:D:50:LYS:NZ	2.17	0.77
3:L:61:ARG:NH2	3:L:82:ASP:OD2	2.20	0.74
3:L:31:ARG:NH1	3:L:66:GLY:O	2.23	0.72
3:D:54:LEU:HD21	3:D:58:VAL:O	1.90	0.71

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	129/132 (98%)	116 (90%)	13 (10%)	0	100	100
1	B	129/132 (98%)	116 (90%)	13 (10%)	0	100	100
2	C	212/222 (96%)	199 (94%)	13 (6%)	0	100	100
2	H	210/222 (95%)	195 (93%)	15 (7%)	0	100	100
3	D	214/219 (98%)	197 (92%)	17 (8%)	0	100	100
3	L	214/219 (98%)	198 (92%)	15 (7%)	1 (0%)	29	68
All	All	1108/1146 (97%)	1021 (92%)	86 (8%)	1 (0%)	51	85

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	L	159	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/114 (99%)	102 (90%)	11 (10%)	8	31
1	B	113/114 (99%)	101 (89%)	12 (11%)	6	26
2	C	184/189 (97%)	168 (91%)	16 (9%)	10	37
2	H	183/189 (97%)	163 (89%)	20 (11%)	6	25
3	D	189/192 (98%)	166 (88%)	23 (12%)	5	21
3	L	189/192 (98%)	167 (88%)	22 (12%)	5	23
All	All	971/990 (98%)	867 (89%)	104 (11%)	6	26

5 of 104 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	L	168	TRP
1	B	22	GLU
3	D	150	ASN
3	L	181	SER
3	L	204	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
3	L	166	ASN
1	B	93	GLN
3	D	38	GLN
3	L	150	ASN
2	C	162	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	131/132 (99%)	-0.47	1 (0%) 86 65	35, 55, 75, 98	0
1	B	131/132 (99%)	-0.18	1 (0%) 86 65	45, 83, 125, 170	0
2	C	216/222 (97%)	-0.41	1 (0%) 91 75	20, 50, 87, 123	0
2	H	214/222 (96%)	-0.51	0 100 100	27, 54, 82, 93	0
3	D	216/219 (98%)	-0.45	0 100 100	33, 61, 84, 98	0
3	L	216/219 (98%)	-0.46	0 100 100	34, 61, 98, 110	0
All	All	1124/1146 (98%)	-0.43	3 (0%) 94 84	20, 58, 99, 170	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	57	PHE	3.3
2	C	138	GLN	2.3
1	A	131	ALA	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.