



Full wwPDB X-ray Structure Validation Report i

Aug 22, 2023 – 05:54 PM EDT

PDB ID : 3C8J
Title : The crystal structure of natural killer cell receptor Ly49C
Authors : Deng, L.; Mariuzza, R.A.
Deposited on : 2008-02-12
Resolution : 2.60 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.35
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 4346 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Natural killer cell receptor Ly49C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	127	1054	687	175	181	11	0	0	0
1	B	126	1050	684	174	181	11	0	0	0
1	C	126	1050	684	174	181	11	0	0	0
1	D	130	1082	701	181	189	11	0	0	0

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	60	MET	-	initiating methionine	UNP Q61198
A	62	SER	VAL	SEE REMARK 999	UNP Q61198
A	74	THR	ILE	SEE REMARK 999	UNP Q61198
A	?	-	HIS	SEE REMARK 999	UNP Q61198
A	116	GLY	ARG	engineered mutation	UNP Q61198
A	119	HIS	-	SEE REMARK 999	UNP Q61198
A	171	GLY	SER	engineered mutation	UNP Q61198
A	193	GLY	GLU	engineered mutation	UNP Q61198
A	223	LYS	ARG	engineered mutation	UNP Q61198
B	60	MET	-	initiating methionine	UNP Q61198
B	62	SER	VAL	SEE REMARK 999	UNP Q61198
B	74	THR	ILE	SEE REMARK 999	UNP Q61198
B	?	-	HIS	variant	UNP Q61198
B	116	GLY	ARG	engineered mutation	UNP Q61198
B	119	HIS	-	SEE REMARK 999	UNP Q61198
B	171	GLY	SER	engineered mutation	UNP Q61198
B	193	GLY	GLU	engineered mutation	UNP Q61198
B	223	LYS	ARG	engineered mutation	UNP Q61198
C	60	MET	-	initiating methionine	UNP Q61198
C	62	SER	VAL	SEE REMARK 999	UNP Q61198
C	74	THR	ILE	SEE REMARK 999	UNP Q61198

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	?	-	HIS	SEE REMARK 999	UNP Q61198
C	116	GLY	ARG	engineered mutation	UNP Q61198
C	119	HIS	-	SEE REMARK 999	UNP Q61198
C	171	GLY	SER	engineered mutation	UNP Q61198
C	193	GLY	GLU	engineered mutation	UNP Q61198
C	223	LYS	ARG	engineered mutation	UNP Q61198
D	60	MET	-	initiating methionine	UNP Q61198
D	62	SER	VAL	SEE REMARK 999	UNP Q61198
D	74	THR	ILE	SEE REMARK 999	UNP Q61198
D	?	-	HIS	SEE REMARK 999	UNP Q61198
D	116	GLY	ARG	engineered mutation	UNP Q61198
D	119	HIS	-	SEE REMARK 999	UNP Q61198
D	171	GLY	SER	engineered mutation	UNP Q61198
D	193	GLY	GLU	engineered mutation	UNP Q61198
D	223	LYS	ARG	engineered mutation	UNP Q61198



- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	20	Total O 20 20	0	0
2	B	44	Total O 44 44	0	0
2	C	12	Total O 12 12	0	0
2	D	34	Total O 34 34	0	0



3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Natural killer cell receptor Ly49C



- Molecule 1: Natural killer cell receptor Ly49C




- Molecule 1: Natural killer cell receptor Ly49C



- Molecule 1: Natural killer cell receptor Ly49C

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.10 Å 94.89 Å 104.19 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.60 41.60 – 2.40	Depositor EDS
% Data completeness (in resolution range)	93.5 (30.00-2.60) 92.4 (41.60-2.40)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	2.13 (at 2.39 Å)	Xtriage
Refinement program	REFMAC 5.2.0003	Depositor
R , R_{free}	0.198 , 0.263 0.261 , 0.306	Depositor DCC
R_{free} test set	1276 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	47.0	Xtriage
Anisotropy	0.268	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 36.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	4346	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.58	0/1085	0.68	0/1457
1	B	0.70	0/1081	0.88	1/1450 (0.1%)
1	C	0.92	2/1081 (0.2%)	0.69	2/1450 (0.1%)
1	D	0.57	0/1113	0.75	2/1493 (0.1%)
All	All	0.71	2/4360 (0.0%)	0.75	5/5850 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	207	GLU	CD-OE2	19.24	1.46	1.25
1	C	207	GLU	CD-OE1	16.75	1.44	1.25

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	207	GLU	OE1-CD-OE2	6.67	131.30	123.30
1	D	148	THR	O-C-N	-6.26	112.68	122.70
1	D	146	TYR	CB-CG-CD2	5.75	124.45	121.00
1	C	148	THR	O-C-N	-5.54	113.84	122.70
1	B	147	SER	CB-CA-C	5.17	119.93	110.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1054	0	1038	14	0
1	B	1050	0	1036	19	0
1	C	1050	0	1036	20	1
1	D	1082	0	1065	29	1
2	A	20	0	0	1	0
2	B	44	0	0	2	0
2	C	12	0	0	1	0
2	D	34	0	0	6	0
All	All	4346	0	4175	78	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:135:ASP:HA	2:D:297:HOH:O	1.48	1.14
1:D:138:ARG:HB3	1:D:156:ASN:HD21	1.24	1.01
1:D:136:THR:OG1	1:D:137:GLY:N	1.96	0.91
1:D:190:VAL:O	1:D:237:LYS:HE2	1.74	0.88
1:C:139:GLY:H	1:C:156:ASN:HD21	1.33	0.77
1:B:205:LYS:HB2	2:B:275:HOH:O	1.86	0.76
1:D:138:ARG:HB3	1:D:156:ASN:ND2	2.01	0.74
1:D:136:THR:HG1	1:D:137:GLY:H	1.37	0.70
1:B:190:VAL:O	1:B:237:LYS:HE2	1.92	0.69
1:D:188:ARG:HD3	2:D:319:HOH:O	1.94	0.67
1:D:251:CYS:HB2	2:D:300:HOH:O	1.95	0.66
1:D:188:ARG:CD	2:D:319:HOH:O	2.45	0.64
1:B:158:THR:HG23	1:B:159:THR:O	1.98	0.63
1:C:172:VAL:HG13	1:C:255:LYS:HB2	1.81	0.62
1:D:167:CYS:HB3	1:D:172:VAL:O	2.00	0.61
1:C:167:CYS:HB3	1:C:172:VAL:O	2.00	0.61
1:B:138:ARG:HH21	1:D:184:LYS:HD2	1.65	0.60
1:C:139:GLY:H	1:C:156:ASN:ND2	2.01	0.58
1:C:153:PHE:CE1	1:C:189:HIS:HE1	2.22	0.58
1:B:228:LYS:O	1:B:229:SER:HB3	2.05	0.57
1:C:139:GLY:N	1:C:156:ASN:HD21	1.99	0.57
1:B:153:PHE:HB3	1:B:155:MET:CE	2.36	0.56
1:A:195:TYR:HB3	1:A:252:ILE:HG13	1.89	0.55
1:B:167:CYS:HB3	1:B:172:VAL:O	2.07	0.55
1:D:133:SER:O	1:D:134:ARG:HG3	2.07	0.54
1:D:166:ASN:O	1:D:169:HIS:HB3	2.08	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:141:LYS:HE2	1:A:170:TYR:CZ	2.44	0.53
1:C:201:TYR:CE2	1:C:230:ARG:HG2	2.45	0.52
1:B:153:PHE:HB3	1:B:155:MET:HE1	1.90	0.52
1:C:143:TRP:HB3	1:C:152:TYR:HD1	1.74	0.52
1:A:152:TYR:HB2	1:A:253:CYS:HB2	1.93	0.51
1:B:228:LYS:O	1:B:229:SER:CB	2.56	0.51
1:D:174:ILE:HG22	1:D:211:ILE:HD12	1.93	0.51
1:D:229:SER:OG	1:D:230:ARG:N	2.43	0.50
1:C:218:LEU:HB3	1:C:222:ILE:HD12	1.94	0.50
1:A:201:TYR:CE2	1:A:230:ARG:HG2	2.47	0.50
1:B:141:LYS:HE2	1:B:170:TYR:CZ	2.47	0.49
1:C:156:ASN:ND2	2:C:265:HOH:O	2.45	0.49
1:A:218:LEU:HB3	1:A:222:ILE:HD12	1.94	0.48
1:B:172:VAL:HG13	1:B:255:LYS:HB2	1.95	0.47
1:A:235:LEU:HD12	1:A:235:LEU:C	2.35	0.47
1:D:158:THR:O	1:D:248:PRO:HA	2.15	0.47
1:D:139:GLY:O	1:D:141:LYS:HE2	2.15	0.46
1:D:199:LEU:HD23	1:D:233:VAL:HG21	1.97	0.46
1:D:163:CYS:HB2	2:D:300:HOH:O	2.14	0.46
1:C:146:TYR:CD1	1:C:185:PHE:CE1	3.04	0.46
1:D:186:LEU:O	1:D:190:VAL:HB	2.16	0.46
1:C:175:LEU:HD21	1:C:186:LEU:CD1	2.46	0.46
1:D:194:ASN:HB3	1:D:234:PHE:CD1	2.52	0.45
1:B:138:ARG:NH2	1:D:184:LYS:HD2	2.31	0.45
1:D:145:CYS:HA	1:D:150:CYS:HA	1.99	0.45
1:B:211:ILE:HA	2:B:276:HOH:O	2.16	0.45
1:B:231:GLY:HA2	1:B:245:CYS:SG	2.57	0.44
1:C:155:MET:CE	1:C:195:TYR:HE2	2.31	0.44
1:A:187:GLN:HE22	1:A:239:ARG:HA	1.81	0.44
1:C:143:TRP:HB3	1:C:152:TYR:CD1	2.52	0.44
1:B:146:TYR:O	1:B:147:SER:C	2.52	0.44
1:B:191:ILE:O	1:B:192:PRO:C	2.56	0.44
1:A:157:LYS:HA	1:A:249:TYR:O	2.17	0.44
1:D:153:PHE:CE2	1:D:190:VAL:HG23	2.53	0.44
1:A:232:CYS:O	1:A:242:ASP:HA	2.18	0.43
1:A:243:ILE:HG12	1:A:247:ILE:HD12	1.99	0.43
1:B:195:TYR:HB3	1:B:252:ILE:HG13	1.99	0.43
1:D:226:ASN:O	1:D:228:LYS:HG2	2.18	0.43
1:A:172:VAL:HG13	1:A:255:LYS:HB3	2.00	0.43
1:B:146:TYR:CD1	1:B:185:PHE:CE1	3.06	0.43
1:A:181:ASP:CG	1:C:138:ARG:HD2	2.39	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:175:LEU:HD21	1:C:186:LEU:HD13	2.01	0.42
1:C:261:PRO:HG2	1:D:150:CYS:SG	2.60	0.42
1:A:140:VAL:HA	2:A:263:HOH:O	2.19	0.42
1:A:190:VAL:HG13	1:A:191:ILE:O	2.20	0.42
1:D:133:SER:C	1:D:134:ARG:HG3	2.40	0.42
1:B:160:TRP:O	1:B:163:CYS:HB3	2.20	0.42
1:C:195:TYR:HB3	1:C:252:ILE:HG13	2.01	0.41
1:C:157:LYS:HB3	1:C:248:PRO:HB2	2.03	0.41
1:C:191:ILE:O	1:C:192:PRO:C	2.57	0.41
1:D:195:TYR:HB3	1:D:252:ILE:HG13	2.03	0.41
1:D:184:LYS:HE2	2:D:294:HOH:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:239:ARG:NH2	1:D:228:LYS:O[2_875]	1.85	0.35

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	125/203 (62%)	114 (91%)	10 (8%)	1 (1%)	19 39
1	B	124/203 (61%)	117 (94%)	5 (4%)	2 (2%)	9 19
1	C	124/203 (61%)	111 (90%)	12 (10%)	1 (1%)	19 39
1	D	128/203 (63%)	116 (91%)	11 (9%)	1 (1%)	19 39
All	All	501/812 (62%)	458 (91%)	38 (8%)	5 (1%)	15 32

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	138	ARG
1	B	229	SER
1	B	230	ARG
1	D	134	ARG
1	C	261	PRO

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	114/188 (61%)	106 (93%)	8 (7%)	15 30
1	B	114/188 (61%)	106 (93%)	8 (7%)	15 30
1	C	114/188 (61%)	104 (91%)	10 (9%)	10 19
1	D	118/188 (63%)	104 (88%)	14 (12%)	5 9
All	All	460/752 (61%)	420 (91%)	40 (9%)	10 20

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	138	ARG
1	A	161	SER
1	A	190	VAL
1	A	218	LEU
1	A	220	MET
1	A	235	LEU
1	A	255	LYS
1	A	259	LYS
1	B	138	ARG
1	B	141	LYS
1	B	156	ASN
1	B	190	VAL
1	B	205	LYS
1	B	211	ILE
1	B	229	SER
1	B	230	ARG
1	C	141	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	180	GLU
1	C	190	VAL
1	C	204	LYS
1	C	218	LEU
1	C	220	MET
1	C	235	LEU
1	C	246	ASN
1	C	257	LEU
1	C	258	ASP
1	D	136	THR
1	D	141	LYS
1	D	161	SER
1	D	163	CYS
1	D	184	LYS
1	D	190	VAL
1	D	211	ILE
1	D	221	LYS
1	D	226	ASN
1	D	230	ARG
1	D	235	LEU
1	D	246	ASN
1	D	258	ASP
1	D	262	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	187	GLN
1	A	194	ASN
1	B	166	ASN
1	B	168	GLN
1	B	194	ASN
1	B	213	ASN
1	C	156	ASN
1	C	166	ASN
1	C	168	GLN
1	C	187	GLN
1	C	189	HIS
1	C	194	ASN
1	C	246	ASN
1	D	156	ASN
1	D	189	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	194	ASN
1	D	246	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.