



wwPDB X-ray Structure Validation Summary Report i

Sep 26, 2023 – 08:52 PM EDT

PDB ID : 6CH9
Title : Crystal structure of a natively-glycosylated B41 SOSIP.664 HIV-1 Envelope Trimer in complex with the broadly-neutralizing antibodies BG18 and 35O22
Authors : Barnes, C.O.; Bjorkman, P.J.
Deposited on : 2018-02-22
Resolution : 4.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 4.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




| Metric | Whole archive (#Entries) | Similar resolution (#Entries, resolution range(Å)) |
|-----------------------|--------------------------|--|
| R_{free} | 130704 | 1098 (5.82-3.80) |
| Clashscore | 141614 | 1172 (5.82-3.80) |
| Ramachandran outliers | 138981 | 1107 (5.82-3.80) |
| Sidechain outliers | 138945 | 1087 (5.82-3.80) |
| RSRZ outliers | 127900 | 1128 (5.98-3.70) |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



Continued on next page...

Continued from previous page...



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

| Mol | Type | Chain | Res | Chirality | Geometry | Clashes | Electron density |
|-----|------|-------|-----|-----------|----------|---------|------------------|
| 10 | NAG | L | 1 | - | - | X | - |
| 13 | NAG | U | 1 | - | - | X | - |
| 13 | MAN | U | 5 | - | - | - | X |
| 7 | NAG | A | 1 | - | - | - | X |
| 7 | NAG | A | 2 | - | - | - | X |
| 7 | BMA | A | 3 | - | - | - | X |
| 7 | BMA | H | 3 | - | - | - | X |
| 8 | MAN | F | 4 | - | - | X | - |
| 9 | NAG | K | 1 | - | - | X | - |

2 Entry composition i

There are 14 unique types of molecules in this entry. The entry contains 11974 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Envelope glycoprotein gp41.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----------|----------|----------|----------|---------|---------|-------|
| 1 | B | 136 | Total | C 1094 | N 692 | O 188 | S 206 | 8 | 0 | 0 |

There are 2 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|---------------------|------------|
| B | 559 | PRO | ILE | conflict | UNP B3UEZ6 |
| B | 605 | CYS | THR | engineered mutation | UNP B3UEZ6 |

- Molecule 2 is a protein called 35O22 Heavy Chain.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----------|-----------|----------|----------|---------|---------|-------|
| 2 | D | 236 | Total | C 1777 | N 1129 | O 296 | S 344 | 8 | 0 | 0 |

- Molecule 3 is a protein called 35O22 Light Chain.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----------|-----------|----------|----------|---------|---------|-------|
| 3 | E | 213 | Total | C 1615 | N 1012 | O 267 | S 328 | 8 | 0 | 0 |

- Molecule 4 is a protein called Envelope glycoprotein gp120.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----------|-----------|----------|----------|---------|---------|-------|
| 4 | G | 448 | Total | C 3499 | N 2198 | O 617 | S 658 | 26 | 0 | 0 |

There are 36 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|-----------------------|------------|
| G | -4 | MET | - | initiating methionine | UNP B3UES2 |
| G | -3 | ASP | - | expression tag | UNP B3UES2 |

Continued on next page...

Continued from previous page...

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|---------------------|------------|
| G | -2 | ALA | - | expression tag | UNP B3UES2 |
| G | -1 | MET | - | expression tag | UNP B3UES2 |
| G | 0 | LYS | - | expression tag | UNP B3UES2 |
| G | 1 | ARG | - | expression tag | UNP B3UES2 |
| G | 2 | GLY | - | expression tag | UNP B3UES2 |
| G | 3 | LEU | - | expression tag | UNP B3UES2 |
| G | 4 | CYS | - | expression tag | UNP B3UES2 |
| G | 5 | CYS | - | expression tag | UNP B3UES2 |
| G | 6 | VAL | - | expression tag | UNP B3UES2 |
| G | 7 | LEU | - | expression tag | UNP B3UES2 |
| G | 8 | LEU | - | expression tag | UNP B3UES2 |
| G | 9 | LEU | - | expression tag | UNP B3UES2 |
| G | 10 | CYS | - | expression tag | UNP B3UES2 |
| G | 11 | GLY | - | expression tag | UNP B3UES2 |
| G | 12 | ALA | - | expression tag | UNP B3UES2 |
| G | 13 | VAL | - | expression tag | UNP B3UES2 |
| G | 14 | PHE | - | expression tag | UNP B3UES2 |
| G | 15 | VAL | - | expression tag | UNP B3UES2 |
| G | 16 | SER | - | expression tag | UNP B3UES2 |
| G | 17 | PRO | - | expression tag | UNP B3UES2 |
| G | 18 | SER | - | expression tag | UNP B3UES2 |
| G | 19 | GLN | - | expression tag | UNP B3UES2 |
| G | 20 | GLU | - | expression tag | UNP B3UES2 |
| G | 21 | ILE | - | expression tag | UNP B3UES2 |
| G | 22 | HIS | - | expression tag | UNP B3UES2 |
| G | 23 | ALA | - | expression tag | UNP B3UES2 |
| G | 24 | ARG | - | expression tag | UNP B3UES2 |
| G | 25 | PHE | - | expression tag | UNP B3UES2 |
| G | 26 | ARG | - | expression tag | UNP B3UES2 |
| G | 27 | ARG | - | expression tag | UNP B3UES2 |
| G | 28 | GLY | - | expression tag | UNP B3UES2 |
| G | 29 | ALA | - | expression tag | UNP B3UES2 |
| G | 30 | ARG | - | expression tag | UNP B3UES2 |
| G | 501 | CYS | ALA | engineered mutation | UNP B3UES2 |


- Molecule 5 is a protein called BG18 Heavy Chain.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|---------|-------|
| 5 | Q | 231 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1731 | 1089 | 298 | 336 | 8 | | | |

- Molecule 6 is a protein called BG18 Light Chain.

| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----------|----------|----------|----------|---------|-------|
| 6 | R | 205 | Total | C 1540 | N 963 | O 257 | S 314 | 0 | 0 |

- Molecule 7 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|---------|---------|--------|---------|---------|-------|
| 7 | A | 3 | Total | C 39 | N 22 | O 2 | S 15 | 0 | 0 |
| 7 | C | 3 | Total | C 39 | N 22 | O 2 | S 15 | 0 | 0 |
| 7 | H | 3 | Total | C 39 | N 22 | O 2 | S 15 | 0 | 0 |
| 7 | J | 3 | Total | C 39 | N 22 | O 2 | S 15 | 0 | 0 |
| 7 | O | 3 | Total | C 39 | N 22 | O 2 | S 15 | 0 | 0 |
| 7 | P | 3 | Total | C 39 | N 22 | O 2 | S 15 | 0 | 0 |
| 7 | T | 3 | Total | C 39 | N 22 | O 2 | S 15 | 0 | 0 |

- Molecule 8 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|---------|---------|--------|---------|---------|-------|
| 8 | F | 4 | Total | C 50 | N 28 | O 2 | S 20 | 0 | 0 |

- Molecule 9 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




| Mol | Chain | Residues | Atoms | ZeroOcc | AltConf | Trace |
|-----|-------|----------|---------------------------|---------|---------|-------|
| 9 | I | 2 | Total C N O 28 16 2 10 | 0 | 0 | 0 |
| 9 | K | 2 | Total C N O 28 16 2 10 | 0 | 0 | 0 |
| 9 | N | 2 | Total C N O 28 16 2 10 | 0 | 0 | 0 |

- Molecule 10 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




| Mol | Chain | Residues | Atoms | ZeroOcc | AltConf | Trace |
|-----|-------|----------|---------------------------|---------|---------|-------|
| 10 | L | 6 | Total C N O 72 40 2 30 | 0 | 0 | 0 |

- Molecule 11 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




| Mol | Chain | Residues | Atoms | ZeroOcc | AltConf | Trace |
|-----|-------|----------|---------------------------|---------|---------|-------|
| 11 | M | 4 | Total C N O 50 28 2 20 | 0 | 0 | 0 |

- Molecule 12 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|----|---|----|---------|---------|-------|
| 12 | S | 5 | Total | C | N | O | 0 | 0 | 0 |
| | | | 61 | 34 | 2 | 25 | | | |

- Molecule 13 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|----|---|----|---------|---------|-------|
| 13 | U | 6 | Total | C | N | O | 0 | 0 | 0 |
| | | | 72 | 40 | 2 | 30 | | | |

- Molecule 14 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).




| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf |
|-----|-------|----------|-------|---|---|---|---------|---------|
| 14 | B | 1 | Total | C | N | O | 0 | 0 |
| | | | 14 | 8 | 1 | 5 | | |
| 14 | G | 1 | Total | C | N | O | 0 | 0 |
| | | | 14 | 8 | 1 | 5 | | |
| 14 | G | 1 | Total | C | N | O | 0 | 0 |
| | | | 14 | 8 | 1 | 5 | | |
| 14 | G | 1 | Total | C | N | O | 0 | 0 |
| | | | 14 | 8 | 1 | 5 | | |


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Envelope glycoprotein gp41




- Molecule 2: 35O22 Heavy Chain




- Molecule 3: 35O22 Light Chain




- Molecule 4: Envelope glycoprotein gp120








- Molecule 5: BG18 Heavy Chain



- Molecule 6: BG18 Light Chain



- Molecule 7: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain A:  100%




- Molecule 7: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose


Chain C: 67%



- Molecule 7: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 7: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 7: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 7: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 7: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 8: alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 9: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose





- Molecule 9: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K: 50% 50%



- Molecule 9: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain N: 50% 50%



- Molecule 10: alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L: 50% 50%



- Molecule 11: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain M: 75% 25%



- Molecule 12: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain S: 60% 40%



- Molecule 13: alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain U: 33% 67%

| |
|------|
| MAG1 |
| MA2 |
| BM43 |
| MA44 |
| MA45 |
| MA46 |

4 Data and refinement statistics i

| Property | Value | Source |
|---|--|------------------|
| Space group | H 3 2 | Depositor |
| Cell constants a, b, c, α , β , γ | 241.08Å 241.08Å 345.47Å 90.00° 90.00° 120.00° | Depositor |
| Resolution (Å) | 39.90 – 4.85 39.90 – 4.85 | Depositor EDS |
| % Data completeness (in resolution range) | 95.3 (39.90-4.85) 95.3 (39.90-4.85) | Depositor EDS |
| R_{merge} | 0.26 | Depositor |
| R_{sym} | (Not available) | Depositor |
| $< I/\sigma(I) >$ ¹ | 1.42 (at 4.84Å) | Xtriage |
| Refinement program | BUSTER 2.10.3 | Depositor |
| R , R_{free} | 0.285 , 0.292 0.298 , 0.328 | Depositor DCC |
| R_{free} test set | 838 reflections (4.75%) | wwPDB-VP |
| Wilson B-factor (Å ²) | 275.7 | Xtriage |
| Anisotropy | 0.141 | Xtriage |
| Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²) | 0.27 , 291.9 | EDS |
| L-test for twinning ² | $< L > = 0.35$, $< L^2 > = 0.18$ | Xtriage |
| Estimated twinning fraction | No twinning to report. | Xtriage |
| F_o, F_c correlation | 0.89 | EDS |
| Total number of atoms | 11974 | wwPDB-VP |
| Average B, all atoms (Å ²) | 300.0 | wwPDB-VP |

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, BMA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|----------------|-------------|-----------------|
| | | RMSZ | # $ Z > 5$ | RMSZ | # $ Z > 5$ |
| 1 | B | 0.41 | 0/1113 | 0.61 | 1/1509 (0.1%) |
| 2 | D | 0.41 | 0/1822 | 0.63 | 2/2481 (0.1%) |
| 3 | E | 0.38 | 0/1659 | 0.58 | 0/2269 |
| 4 | G | 0.45 | 1/3572 (0.0%) | 0.91 | 18/4857 (0.4%) |
| 5 | Q | 0.38 | 0/1776 | 0.65 | 1/2423 (0.0%) |
| 6 | R | 0.44 | 0/1577 | 0.65 | 0/2153 |
| All | All | 0.42 | 1/11519 (0.0%) | 0.72 | 22/15692 (0.1%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 4 | G | 0 | 1 |

All (1) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|------|-------------|----------|
| 4 | G | 392 | ASN | C-N | 6.72 | 1.49 | 1.34 |

The worst 5 of 22 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|--------|--------|-------------|----------|
| 4 | G | 234 | ASN | O-C-N | -16.57 | 95.02 | 123.20 |
| 4 | G | 233 | PHE | O-C-N | 13.15 | 143.75 | 122.70 |
| 4 | G | 275 | GLU | O-C-N | 11.18 | 140.59 | 122.70 |
| 4 | G | 295 | ASN | O-C-N | -10.77 | 105.46 | 122.70 |
| 4 | G | 234 | ASN | CA-C-N | 10.41 | 137.02 | 116.20 |

There are no chirality outliers.

All (1) planarity outliers are listed below:

| Mol | Chain | Res | Type | Group |
|-----|-------|-----|------|-----------|
| 4 | G | 234 | ASN | Mainchain |

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | B | 1094 | 0 | 1075 | 34 | 0 |
| 2 | D | 1777 | 0 | 1755 | 20 | 0 |
| 3 | E | 1615 | 0 | 1550 | 10 | 0 |
| 4 | G | 3499 | 0 | 3440 | 197 | 0 |
| 5 | Q | 1731 | 0 | 1685 | 45 | 0 |
| 6 | R | 1540 | 0 | 1496 | 15 | 0 |
| 7 | A | 39 | 0 | 34 | 1 | 0 |
| 7 | C | 39 | 0 | 34 | 1 | 0 |
| 7 | H | 39 | 0 | 34 | 5 | 0 |
| 7 | J | 39 | 0 | 34 | 0 | 0 |
| 7 | O | 39 | 0 | 34 | 1 | 0 |
| 7 | P | 39 | 0 | 34 | 1 | 0 |
| 7 | T | 39 | 0 | 34 | 0 | 0 |
| 8 | F | 50 | 0 | 43 | 11 | 0 |
| 9 | I | 28 | 0 | 25 | 0 | 0 |
| 9 | K | 28 | 0 | 25 | 10 | 0 |
| 9 | N | 28 | 0 | 25 | 1 | 0 |
| 10 | L | 72 | 0 | 61 | 11 | 0 |
| 11 | M | 50 | 0 | 43 | 5 | 0 |
| 12 | S | 61 | 0 | 52 | 4 | 0 |
| 13 | U | 72 | 0 | 61 | 18 | 0 |
| 14 | B | 14 | 0 | 13 | 4 | 0 |
| 14 | G | 42 | 0 | 39 | 0 | 0 |
| All | All | 11974 | 0 | 11626 | 311 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 311 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|-----------------|------------------|--------------------------|-------------------|
| 4:G:333:ILE:CD1 | 4:G:414:ILE:HB | 1.30 | 1.60 |
| 4:G:338:TRP:CZ3 | 4:G:390:LEU:HD22 | 1.37 | 1.57 |
| 4:G:292:VAL:HB | 4:G:449:ILE:CD1 | 1.30 | 1.55 |
| 4:G:292:VAL:CB | 4:G:449:ILE:HD12 | 1.46 | 1.44 |
| 4:G:292:VAL:O | 4:G:449:ILE:CD1 | 1.68 | 1.41 |

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles |
|-----|-------|-----------------|------------|-----------|----------|-------------|
| 1 | B | 132/153 (86%) | 119 (90%) | 13 (10%) | 0 | 100 100 |
| 2 | D | 232/243 (96%) | 205 (88%) | 26 (11%) | 1 (0%) | 34 72 |
| 3 | E | 211/216 (98%) | 192 (91%) | 17 (8%) | 2 (1%) | 17 56 |
| 4 | G | 440/518 (85%) | 366 (83%) | 69 (16%) | 5 (1%) | 14 51 |
| 5 | Q | 229/240 (95%) | 178 (78%) | 44 (19%) | 7 (3%) | 4 30 |
| 6 | R | 201/215 (94%) | 140 (70%) | 54 (27%) | 7 (4%) | 3 28 |
| All | All | 1445/1585 (91%) | 1200 (83%) | 223 (15%) | 22 (2%) | 10 46 |

5 of 22 Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 3 | E | 53 | ASP |
| 4 | G | 197 | ASN |
| 5 | Q | 51 | ASP |
| 5 | Q | 159 | VAL |
| 6 | R | 29 | LEU |

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles |
|-----|-------|-----------------|------------|----------|-------------|
| 1 | B | 119/130 (92%) | 118 (99%) | 1 (1%) | 81 89 |
| 2 | D | 199/206 (97%) | 192 (96%) | 7 (4%) | 36 60 |
| 3 | E | 186/189 (98%) | 181 (97%) | 5 (3%) | 44 66 |
| 4 | G | 393/455 (86%) | 361 (92%) | 32 (8%) | 11 37 |
| 5 | Q | 193/207 (93%) | 188 (97%) | 5 (3%) | 46 67 |
| 6 | R | 174/182 (96%) | 165 (95%) | 9 (5%) | 23 49 |
| All | All | 1264/1369 (92%) | 1205 (95%) | 59 (5%) | 26 52 |

5 of 59 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 4 | G | 295 | ASN |
| 6 | R | 152 | TRP |
| 4 | G | 364 | SER |
| 6 | R | 141 | SER |
| 5 | Q | 205 | SER |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 4 | G | 363 | HIS |
| 4 | G | 463 | ASN |
| 6 | R | 192 | HIS |
| 4 | G | 478 | ASN |
| 4 | G | 130 | ASN |

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

52 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 7 | NAG | A | 1 | 1,7 | 14,14,15 | 0.92 | 0 | 17,19,21 | 2.23 | 5 (29%) |
| 7 | NAG | A | 2 | 7 | 14,14,15 | 0.81 | 1 (7%) | 17,19,21 | 1.21 | 2 (11%) |
| 7 | BMA | A | 3 | 7 | 11,11,12 | 0.40 | 0 | 15,15,17 | 0.90 | 0 |
| 7 | NAG | C | 1 | 1,7 | 14,14,15 | 0.79 | 0 | 17,19,21 | 1.88 | 3 (17%) |
| 7 | NAG | C | 2 | 7 | 14,14,15 | 0.67 | 0 | 17,19,21 | 2.17 | 4 (23%) |
| 7 | BMA | C | 3 | 7 | 11,11,12 | 0.44 | 0 | 15,15,17 | 0.98 | 1 (6%) |
| 8 | NAG | F | 1 | 8 | 14,14,15 | 0.95 | 0 | 17,19,21 | 2.21 | 6 (35%) |
| 8 | NAG | F | 2 | 8 | 14,14,15 | 1.12 | 2 (14%) | 17,19,21 | 2.75 | 7 (41%) |
| 8 | BMA | F | 3 | 8 | 11,11,12 | 1.53 | 2 (18%) | 15,15,17 | 2.28 | 6 (40%) |
| 8 | MAN | F | 4 | 8 | 11,11,12 | 2.18 | 2 (18%) | 15,15,17 | 2.22 | 8 (53%) |
| 7 | NAG | H | 1 | 4,7 | 14,14,15 | 0.95 | 1 (7%) | 17,19,21 | 1.89 | 7 (41%) |
| 7 | NAG | H | 2 | 7 | 14,14,15 | 1.21 | 2 (14%) | 17,19,21 | 2.30 | 5 (29%) |
| 7 | BMA | H | 3 | 7 | 11,11,12 | 0.84 | 0 | 15,15,17 | 1.72 | 3 (20%) |
| 9 | NAG | I | 1 | 9,4 | 14,14,15 | 1.02 | 0 | 17,19,21 | 1.23 | 1 (5%) |
| 9 | NAG | I | 2 | 9 | 14,14,15 | 1.14 | 1 (7%) | 17,19,21 | 1.64 | 4 (23%) |
| 7 | NAG | J | 1 | 4,7 | 14,14,15 | 0.81 | 0 | 17,19,21 | 1.71 | 4 (23%) |
| 7 | NAG | J | 2 | 7 | 14,14,15 | 1.35 | 1 (7%) | 17,19,21 | 1.47 | 3 (17%) |
| 7 | BMA | J | 3 | 7 | 11,11,12 | 0.76 | 0 | 15,15,17 | 1.27 | 2 (13%) |
| 9 | NAG | K | 1 | 9,4 | 14,14,15 | 0.89 | 1 (7%) | 17,19,21 | 2.26 | 6 (35%) |
| 9 | NAG | K | 2 | 9 | 14,14,15 | 0.65 | 1 (7%) | 17,19,21 | 0.98 | 0 |
| 10 | NAG | L | 1 | 10,4 | 14,14,15 | 1.13 | 1 (7%) | 17,19,21 | 2.94 | 8 (47%) |
| 10 | NAG | L | 2 | 10 | 14,14,15 | 0.73 | 0 | 17,19,21 | 2.42 | 7 (41%) |

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 10 | BMA | L | 3 | 10 | 11,11,12 | 0.74 | 0 | 15,15,17 | 2.73 | 7 (46%) |
| 10 | MAN | L | 4 | 10 | 11,11,12 | 0.42 | 0 | 15,15,17 | 2.10 | 3 (20%) |
| 10 | MAN | L | 5 | 10 | 11,11,12 | 0.77 | 0 | 15,15,17 | 1.57 | 1 (6%) |
| 10 | MAN | L | 6 | 10 | 11,11,12 | 1.02 | 1 (9%) | 15,15,17 | 1.58 | 2 (13%) |
| 11 | NAG | M | 1 | 4,11 | 14,14,15 | 1.36 | 2 (14%) | 17,19,21 | 2.47 | 5 (29%) |
| 11 | NAG | M | 2 | 11 | 14,14,15 | 0.58 | 0 | 17,19,21 | 1.49 | 2 (11%) |
| 11 | BMA | M | 3 | 11 | 11,11,12 | 0.91 | 1 (9%) | 15,15,17 | 1.89 | 5 (33%) |
| 11 | MAN | M | 4 | 11 | 11,11,12 | 0.67 | 0 | 15,15,17 | 1.68 | 3 (20%) |
| 9 | NAG | N | 1 | 9,4 | 14,14,15 | 0.81 | 1 (7%) | 17,19,21 | 2.23 | 3 (17%) |
| 9 | NAG | N | 2 | 9 | 14,14,15 | 0.47 | 0 | 17,19,21 | 1.28 | 3 (17%) |
| 7 | NAG | O | 1 | 4,7 | 14,14,15 | 0.37 | 0 | 17,19,21 | 1.28 | 3 (17%) |
| 7 | NAG | O | 2 | 7 | 14,14,15 | 0.72 | 0 | 17,19,21 | 1.64 | 3 (17%) |
| 7 | BMA | O | 3 | 7 | 11,11,12 | 0.39 | 0 | 15,15,17 | 1.19 | 1 (6%) |
| 7 | NAG | P | 1 | 4,7 | 14,14,15 | 0.64 | 0 | 17,19,21 | 1.49 | 2 (11%) |
| 7 | NAG | P | 2 | 7 | 14,14,15 | 0.81 | 1 (7%) | 17,19,21 | 1.24 | 2 (11%) |
| 7 | BMA | P | 3 | 7 | 11,11,12 | 0.46 | 0 | 15,15,17 | 1.01 | 1 (6%) |
| 12 | NAG | S | 1 | 4,12 | 14,14,15 | 1.28 | 1 (7%) | 17,19,21 | 1.66 | 4 (23%) |
| 12 | NAG | S | 2 | 12 | 14,14,15 | 1.36 | 2 (14%) | 17,19,21 | 2.61 | 6 (35%) |
| 12 | BMA | S | 3 | 12 | 11,11,12 | 1.34 | 1 (9%) | 15,15,17 | 2.64 | 5 (33%) |
| 12 | MAN | S | 4 | 12 | 11,11,12 | 0.86 | 1 (9%) | 15,15,17 | 2.75 | 8 (53%) |
| 12 | MAN | S | 5 | 12 | 11,11,12 | 0.73 | 0 | 15,15,17 | 1.28 | 2 (13%) |
| 7 | NAG | T | 1 | 4,7 | 14,14,15 | 0.68 | 0 | 17,19,21 | 1.64 | 4 (23%) |
| 7 | NAG | T | 2 | 7 | 14,14,15 | 0.82 | 0 | 17,19,21 | 1.45 | 4 (23%) |
| 7 | BMA | T | 3 | 7 | 11,11,12 | 0.66 | 0 | 15,15,17 | 1.72 | 3 (20%) |
| 13 | NAG | U | 1 | 13 | 14,14,15 | 1.48 | 4 (28%) | 17,19,21 | 2.16 | 5 (29%) |
| 13 | NAG | U | 2 | 13 | 14,14,15 | 0.86 | 0 | 17,19,21 | 2.91 | 7 (41%) |
| 13 | BMA | U | 3 | 13 | 11,11,12 | 0.78 | 0 | 15,15,17 | 2.42 | 5 (33%) |
| 13 | MAN | U | 4 | 13 | 11,11,12 | 0.65 | 0 | 15,15,17 | 1.94 | 5 (33%) |
| 13 | MAN | U | 5 | 13 | 11,11,12 | 0.53 | 0 | 15,15,17 | 1.35 | 1 (6%) |
| 13 | MAN | U | 6 | 13 | 11,11,12 | 0.87 | 1 (9%) | 15,15,17 | 3.02 | 9 (60%) |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|-----------|---------|
| 7 | NAG | A | 1 | 1,7 | - | 4/6/23/26 | 0/1/1/1 |
| 7 | NAG | A | 2 | 7 | - | 4/6/23/26 | 0/1/1/1 |
| 7 | BMA | A | 3 | 7 | - | 0/2/19/22 | 0/1/1/1 |
| 7 | NAG | C | 1 | 1,7 | - | 6/6/23/26 | 0/1/1/1 |
| 7 | NAG | C | 2 | 7 | - | 1/6/23/26 | 0/1/1/1 |
| 7 | BMA | C | 3 | 7 | - | 1/2/19/22 | 0/1/1/1 |
| 8 | NAG | F | 1 | 8 | - | 4/6/23/26 | 0/1/1/1 |
| 8 | NAG | F | 2 | 8 | - | 4/6/23/26 | 0/1/1/1 |
| 8 | BMA | F | 3 | 8 | - | 2/2/19/22 | 0/1/1/1 |
| 8 | MAN | F | 4 | 8 | - | 2/2/19/22 | 0/1/1/1 |
| 7 | NAG | H | 1 | 4,7 | - | 5/6/23/26 | 0/1/1/1 |
| 7 | NAG | H | 2 | 7 | - | 4/6/23/26 | 0/1/1/1 |
| 7 | BMA | H | 3 | 7 | - | 2/2/19/22 | 0/1/1/1 |
| 9 | NAG | I | 1 | 9,4 | - | 3/6/23/26 | 0/1/1/1 |
| 9 | NAG | I | 2 | 9 | - | 3/6/23/26 | 0/1/1/1 |
| 7 | NAG | J | 1 | 4,7 | - | 2/6/23/26 | 0/1/1/1 |
| 7 | NAG | J | 2 | 7 | - | 4/6/23/26 | 0/1/1/1 |
| 7 | BMA | J | 3 | 7 | - | 1/2/19/22 | 0/1/1/1 |
| 9 | NAG | K | 1 | 9,4 | - | 5/6/23/26 | 0/1/1/1 |
| 9 | NAG | K | 2 | 9 | - | 3/6/23/26 | 0/1/1/1 |
| 10 | NAG | L | 1 | 10,4 | - | 5/6/23/26 | 0/1/1/1 |
| 10 | NAG | L | 2 | 10 | - | 1/6/23/26 | 0/1/1/1 |
| 10 | BMA | L | 3 | 10 | - | 2/2/19/22 | 0/1/1/1 |
| 10 | MAN | L | 4 | 10 | - | 2/2/19/22 | 0/1/1/1 |
| 10 | MAN | L | 5 | 10 | - | 1/2/19/22 | 0/1/1/1 |
| 10 | MAN | L | 6 | 10 | - | 1/2/19/22 | 0/1/1/1 |
| 11 | NAG | M | 1 | 4,11 | - | 4/6/23/26 | 0/1/1/1 |
| 11 | NAG | M | 2 | 11 | - | 3/6/23/26 | 0/1/1/1 |
| 11 | BMA | M | 3 | 11 | - | 0/2/19/22 | 0/1/1/1 |
| 11 | MAN | M | 4 | 11 | - | 0/2/19/22 | 0/1/1/1 |
| 9 | NAG | N | 1 | 9,4 | - | 5/6/23/26 | 0/1/1/1 |
| 9 | NAG | N | 2 | 9 | - | 4/6/23/26 | 0/1/1/1 |
| 7 | NAG | O | 1 | 4,7 | - | 4/6/23/26 | 0/1/1/1 |
| 7 | NAG | O | 2 | 7 | - | 5/6/23/26 | 0/1/1/1 |
| 7 | BMA | O | 3 | 7 | - | 1/2/19/22 | 0/1/1/1 |
| 7 | NAG | P | 1 | 4,7 | - | 3/6/23/26 | 0/1/1/1 |
| 7 | NAG | P | 2 | 7 | - | 2/6/23/26 | 0/1/1/1 |

Continued on next page...

Continued from previous page...

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|-----------|---------|
| 7 | BMA | P | 3 | 7 | - | 1/2/19/22 | 0/1/1/1 |
| 12 | NAG | S | 1 | 4,12 | - | 2/6/23/26 | 0/1/1/1 |
| 12 | NAG | S | 2 | 12 | - | 2/6/23/26 | 0/1/1/1 |
| 12 | BMA | S | 3 | 12 | - | 2/2/19/22 | 0/1/1/1 |
| 12 | MAN | S | 4 | 12 | - | 2/2/19/22 | 0/1/1/1 |
| 12 | MAN | S | 5 | 12 | - | 1/2/19/22 | 0/1/1/1 |
| 7 | NAG | T | 1 | 4,7 | - | 4/6/23/26 | 0/1/1/1 |
| 7 | NAG | T | 2 | 7 | - | 2/6/23/26 | 0/1/1/1 |
| 7 | BMA | T | 3 | 7 | - | 1/2/19/22 | 0/1/1/1 |
| 13 | NAG | U | 1 | 13 | - | 4/6/23/26 | 0/1/1/1 |
| 13 | NAG | U | 2 | 13 | - | 2/6/23/26 | 0/1/1/1 |
| 13 | BMA | U | 3 | 13 | - | 0/2/19/22 | 0/1/1/1 |
| 13 | MAN | U | 4 | 13 | - | 1/2/19/22 | 0/1/1/1 |
| 13 | MAN | U | 5 | 13 | - | 0/2/19/22 | 0/1/1/1 |
| 13 | MAN | U | 6 | 13 | - | 2/2/19/22 | 0/1/1/1 |

The worst 5 of 31 bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|------|-------------|----------|
| 8 | F | 4 | MAN | O5-C1 | 5.83 | 1.53 | 1.43 |
| 8 | F | 4 | MAN | C1-C2 | 3.43 | 1.60 | 1.52 |
| 11 | M | 1 | NAG | C1-C2 | 3.42 | 1.57 | 1.52 |
| 12 | S | 3 | BMA | C2-C3 | 3.39 | 1.57 | 1.52 |
| 7 | J | 2 | NAG | C1-C2 | 3.33 | 1.57 | 1.52 |

The worst 5 of 206 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|----------|-------|-------------|----------|
| 10 | L | 1 | NAG | C2-N2-C7 | 7.59 | 133.71 | 122.90 |
| 9 | N | 1 | NAG | C2-N2-C7 | 7.43 | 133.49 | 122.90 |
| 13 | U | 6 | MAN | C1-C2-C3 | 7.37 | 118.73 | 109.67 |
| 12 | S | 2 | NAG | C2-N2-C7 | 6.73 | 132.49 | 122.90 |
| 13 | U | 2 | NAG | O4-C4-C5 | -6.68 | 92.71 | 109.30 |

There are no chirality outliers.

5 of 129 torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|-------------|
| 7 | A | 1 | NAG | C3-C2-N2-C7 |

Continued on next page...

Continued from previous page...


| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|-------------|
| 7 | A | 1 | NAG | C8-C7-N2-C2 |
| 7 | A | 1 | NAG | O7-C7-N2-C2 |
| 7 | A | 2 | NAG | C8-C7-N2-C2 |
| 7 | A | 2 | NAG | O7-C7-N2-C2 |



There are no ring outliers.



21 monomers are involved in 69 short contacts:



| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|-----|-------|-----|------|---------|--------------|
| 12 | S | 4 | MAN | 2 | 0 |
| 10 | L | 1 | NAG | 9 | 0 |
| 10 | L | 6 | MAN | 1 | 0 |
| 8 | F | 1 | NAG | 2 | 0 |
| 9 | N | 1 | NAG | 1 | 0 |
| 13 | U | 3 | BMA | 2 | 0 |
| 13 | U | 2 | NAG | 5 | 0 |
| 7 | O | 1 | NAG | 1 | 0 |
| 8 | F | 4 | MAN | 7 | 0 |
| 7 | P | 1 | NAG | 1 | 0 |
| 9 | K | 1 | NAG | 10 | 0 |
| 10 | L | 2 | NAG | 1 | 0 |
| 11 | M | 1 | NAG | 5 | 0 |
| 7 | O | 2 | NAG | 1 | 0 |
| 7 | H | 1 | NAG | 5 | 0 |
| 8 | F | 2 | NAG | 2 | 0 |
| 13 | U | 1 | NAG | 10 | 0 |
| 13 | U | 6 | MAN | 1 | 0 |
| 12 | S | 1 | NAG | 2 | 0 |
| 7 | C | 1 | NAG | 1 | 0 |
| 7 | A | 3 | BMA | 1 | 0 |


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.


























5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 14 | NAG | G | 606 | 4 | 14,14,15 | 1.20 | 1 (7%) | 17,19,21 | 1.93 | 5 (29%) |

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 14 | NAG | G | 622 | 4 | 14,14,15 | 0.90 | 1 (7%) | 17,19,21 | 1.92 | 6 (35%) |
| 14 | NAG | G | 631 | 4 | 14,14,15 | 0.60 | 0 | 17,19,21 | 1.77 | 4 (23%) |
| 14 | NAG | B | 704 | 1 | 14,14,15 | 0.30 | 0 | 17,19,21 | 1.11 | 2 (11%) |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|-----------|---------|
| 14 | NAG | G | 606 | 4 | - | 5/6/23/26 | 0/1/1/1 |
| 14 | NAG | G | 622 | 4 | - | 5/6/23/26 | 0/1/1/1 |
| 14 | NAG | G | 631 | 4 | - | 1/6/23/26 | 0/1/1/1 |
| 14 | NAG | B | 704 | 1 | - | 4/6/23/26 | 0/1/1/1 |

All (2) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|------|-------------|----------|
| 14 | G | 606 | NAG | C1-C2 | 3.53 | 1.57 | 1.52 |
| 14 | G | 622 | NAG | C1-C2 | 2.36 | 1.55 | 1.52 |

The worst 5 of 17 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|----------|-------|-------------|----------|
| 14 | G | 606 | NAG | C2-N2-C7 | 5.43 | 130.64 | 122.90 |
| 14 | G | 631 | NAG | C2-N2-C7 | 4.16 | 128.83 | 122.90 |
| 14 | G | 631 | NAG | O5-C5-C6 | 3.74 | 113.07 | 107.20 |
| 14 | G | 622 | NAG | C2-N2-C7 | 3.10 | 127.31 | 122.90 |
| 14 | G | 622 | NAG | C6-C5-C4 | -3.07 | 105.82 | 113.00 |

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|-------------|
| 14 | B | 704 | NAG | C8-C7-N2-C2 |
| 14 | B | 704 | NAG | O7-C7-N2-C2 |
| 14 | G | 606 | NAG | C3-C2-N2-C7 |
| 14 | G | 606 | NAG | C8-C7-N2-C2 |
| 14 | G | 606 | NAG | O7-C7-N2-C2 |

There are no ring outliers.

1 monomer is involved in 4 short contacts:

| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|-----|-------|-----|------|---------|--------------|
| 14 | B | 704 | NAG | 4 | 0 |

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

| Mol | Chain | Analysed | <RSRZ> | #RSRZ>2 | OWAB(Å ²) | Q<0.9 |
|-----|-------|-----------------|--------|---------------|-----------------------|-------|
| 1 | B | 136/153 (88%) | 0.13 | 6 (4%) 34 29 | 183, 285, 325, 365 | 0 |
| 2 | D | 236/243 (97%) | 0.22 | 9 (3%) 40 33 | 207, 303, 325, 325 | 0 |
| 3 | E | 213/216 (98%) | 0.02 | 8 (3%) 40 33 | 196, 297, 325, 325 | 0 |
| 4 | G | 448/518 (86%) | -0.00 | 7 (1%) 72 63 | 186, 292, 325, 375 | 0 |
| 5 | Q | 231/240 (96%) | 0.24 | 13 (5%) 24 21 | 210, 298, 325, 325 | 0 |
| 6 | R | 205/215 (95%) | 0.25 | 9 (4%) 34 29 | 219, 309, 325, 325 | 0 |
| All | All | 1469/1585 (92%) | 0.12 | 52 (3%) 44 36 | 183, 298, 325, 375 | 0 |

The worst 5 of 52 RSRZ outliers are listed below:

| Mol | Chain | Res | Type | RSRZ |
|-----|-------|-----|------|------|
| 2 | D | 216 | CYS | 5.0 |
| 4 | G | 73 | ALA | 4.9 |
| 5 | Q | 150 | GLY | 4.8 |
| 5 | Q | 149 | SER | 4.7 |
| 1 | B | 600 | GLY | 4.4 |

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


| Mol | Type | Chain | Res | Atoms | RSCC | RSR | B-factors(Å ²) | Q<0.9 |
|-----|------|-------|-----|-------|------|------|----------------------------|-------|
| 7 | BMA | A | 3 | 11/12 | 0.42 | 0.41 | 420,420,420,420 | 0 |
| 7 | BMA | O | 3 | 11/12 | 0.47 | 0.22 | 435,435,435,435 | 0 |
| 13 | MAN | U | 5 | 11/12 | 0.48 | 0.73 | 366,366,366,366 | 0 |
| 7 | NAG | A | 2 | 14/15 | 0.52 | 0.69 | 430,430,430,430 | 0 |
| 7 | BMA | H | 3 | 11/12 | 0.52 | 0.42 | 478,478,478,478 | 0 |
| 12 | MAN | S | 4 | 11/12 | 0.56 | 0.35 | 327,327,327,327 | 0 |
| 7 | BMA | J | 3 | 11/12 | 0.57 | 0.34 | 441,441,441,441 | 0 |
| 9 | NAG | I | 2 | 14/15 | 0.60 | 0.33 | 433,433,433,433 | 0 |
| 8 | MAN | F | 4 | 11/12 | 0.62 | 0.26 | 360,360,360,360 | 0 |
| 8 | NAG | F | 2 | 14/15 | 0.64 | 0.29 | 337,337,337,337 | 0 |
| 7 | BMA | C | 3 | 11/12 | 0.66 | 0.29 | 450,450,450,450 | 0 |
| 11 | MAN | M | 4 | 11/12 | 0.67 | 0.25 | 456,456,456,456 | 0 |
| 8 | NAG | F | 1 | 14/15 | 0.67 | 0.31 | 343,343,343,343 | 0 |
| 7 | NAG | H | 2 | 14/15 | 0.67 | 0.40 | 449,449,449,449 | 0 |
| 12 | BMA | S | 3 | 11/12 | 0.70 | 0.23 | 415,415,415,415 | 0 |
| 9 | NAG | I | 1 | 14/15 | 0.73 | 0.30 | 396,396,396,396 | 0 |
| 7 | BMA | T | 3 | 11/12 | 0.74 | 0.18 | 415,415,415,415 | 0 |
| 7 | NAG | A | 1 | 14/15 | 0.76 | 0.40 | 411,411,411,411 | 0 |
| 7 | NAG | C | 2 | 14/15 | 0.76 | 0.38 | 449,449,449,449 | 0 |
| 8 | BMA | F | 3 | 11/12 | 0.76 | 0.25 | 349,349,349,349 | 0 |
| 9 | NAG | N | 1 | 14/15 | 0.76 | 0.36 | 391,391,391,391 | 0 |
| 10 | NAG | L | 1 | 14/15 | 0.78 | 0.28 | 396,396,396,396 | 0 |
| 11 | BMA | M | 3 | 11/12 | 0.78 | 0.21 | 432,432,432,432 | 0 |
| 9 | NAG | K | 1 | 14/15 | 0.78 | 0.17 | 311,311,311,311 | 0 |
| 10 | MAN | L | 4 | 11/12 | 0.79 | 0.12 | 387,387,387,387 | 0 |
| 12 | NAG | S | 2 | 14/15 | 0.80 | 0.16 | 398,398,398,398 | 0 |
| 12 | MAN | S | 5 | 11/12 | 0.81 | 0.30 | 397,397,397,397 | 0 |
| 7 | NAG | H | 1 | 14/15 | 0.81 | 0.21 | 383,383,383,383 | 0 |
| 10 | MAN | L | 6 | 11/12 | 0.82 | 0.19 | 369,369,369,369 | 0 |
| 10 | MAN | L | 5 | 11/12 | 0.85 | 0.13 | 416,416,416,416 | 0 |
| 7 | BMA | P | 3 | 11/12 | 0.86 | 0.18 | 417,417,417,417 | 0 |
| 7 | NAG | O | 2 | 14/15 | 0.86 | 0.32 | 428,428,428,428 | 0 |
| 7 | NAG | O | 1 | 14/15 | 0.86 | 0.40 | 394,394,394,394 | 0 |
| 12 | NAG | S | 1 | 14/15 | 0.87 | 0.17 | 343,343,343,343 | 0 |
| 9 | NAG | K | 2 | 14/15 | 0.88 | 0.19 | 333,333,333,333 | 0 |
| 13 | NAG | U | 1 | 14/15 | 0.88 | 0.17 | 352,352,352,352 | 0 |
| 13 | BMA | U | 3 | 11/12 | 0.88 | 0.13 | 334,334,334,334 | 0 |
| 7 | NAG | J | 1 | 14/15 | 0.88 | 0.22 | 384,384,384,384 | 0 |
| 10 | NAG | L | 2 | 14/15 | 0.89 | 0.24 | 340,340,340,340 | 0 |
| 9 | NAG | N | 2 | 14/15 | 0.90 | 0.28 | 422,422,422,422 | 0 |
| 7 | NAG | P | 1 | 14/15 | 0.90 | 0.36 | 386,386,386,386 | 0 |
| 7 | NAG | C | 1 | 14/15 | 0.90 | 0.14 | 428,428,428,428 | 0 |
| 13 | MAN | U | 4 | 11/12 | 0.90 | 0.17 | 373,373,373,373 | 0 |

Continued on next page...

Continued from previous page...




| Mol | Type | Chain | Res | Atoms | RSCC | RSR | B-factors(Å ²) | Q<0.9 |
|-----|------|-------|-----|-------|------|------|----------------------------|-------|
| 7 | NAG | J | 2 | 14/15 | 0.90 | 0.29 | 418,418,418,418 | 0 |
| 13 | MAN | U | 6 | 11/12 | 0.90 | 0.12 | 331,331,331,331 | 0 |
| 7 | NAG | T | 2 | 14/15 | 0.91 | 0.24 | 425,425,425,425 | 0 |
| 7 | NAG | T | 1 | 14/15 | 0.93 | 0.16 | 380,380,380,380 | 0 |
| 13 | NAG | U | 2 | 14/15 | 0.93 | 0.26 | 359,359,359,359 | 0 |
| 11 | NAG | M | 2 | 14/15 | 0.94 | 0.12 | 405,405,405,405 | 0 |
| 11 | NAG | M | 1 | 14/15 | 0.94 | 0.21 | 390,390,390,390 | 0 |
| 7 | NAG | P | 2 | 14/15 | 0.95 | 0.23 | 409,409,409,409 | 0 |
| 10 | BMA | L | 3 | 11/12 | 0.95 | 0.07 | 386,386,386,386 | 0 |

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain C:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain H:**

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain J:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain O:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain P:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain T:**

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain F:


$2mF_o\text{-}DF_c$ (at 0.7 rmsd) in gray
 $mF_o\text{-}DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain I:**

$2mF_o\text{-}DF_c$ (at 0.7 rmsd) in gray
 $mF_o\text{-}DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)














Electron density around Chain L:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain M:**

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

| Mol | Type | Chain | Res | Atoms | RSCC | RSR | B-factors(Å ²) | Q<0.9 |
|-----|------|-------|-----|-------|------|------|----------------------------|-------|
| 14 | NAG | B | 704 | 14/15 | 0.78 | 0.40 | 326,326,326,326 | 0 |
| 14 | NAG | G | 631 | 14/15 | 0.84 | 0.40 | 365,365,365,365 | 0 |
| 14 | NAG | G | 622 | 14/15 | 0.89 | 0.23 | 367,367,367,367 | 0 |
| 14 | NAG | G | 606 | 14/15 | 0.92 | 0.17 | 368,368,368,368 | 0 |

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.