wwPDB NMR Structure Validation Summary Report (i)

Feb 26, 2022 – 09:27 AM EST

PDB ID	:	2CQ0
Title	:	solution structure of RNA binding domain in eukaryotic translation initiation factor 3 subunit 4
Authors	:	Tsuda, K.; Muto, Y.; Inoue, M.; Kigawa, T.; Terada, T.; Shirouzu, M.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on	:	2005-05-19

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	2.27
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.27

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	103	50% 22% • 25%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 7 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:239-A:315 (77)	0.22	7

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 5 single-model clusters were found.

Cluster number	Models
1	1, 4, 6, 8, 9, 13, 14
2	3, 5, 7, 15
3	11, 17, 19, 20
Single-model clusters	2; 10; 12; 16; 18

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1538 atoms, of which 753 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Eukaryotic translation initiation factor 3 subunit 4.

Mol	Chain	Residues	Atoms					Trace
			Total	C	H	N	O	
1	A	103	1538	482	753	146	157	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	224	GLY	-	cloning artifact	UNP O75821
A	225	SER	-	cloning artifact	UNP O75821
A	226	SER	-	cloning artifact	UNP O75821
A	227	GLY	-	cloning artifact	UNP O75821
A	228	SER	-	cloning artifact	UNP O75821
A	229	SER	-	cloning artifact	UNP O75821
A	230	GLY	-	cloning artifact	UNP O75821
A	321	SER	-	cloning artifact	UNP O75821
A	322	GLY	-	cloning artifact	UNP O75821
A	323	PRO	-	cloning artifact	UNP O75821
A	324	SER	-	cloning artifact	UNP O75821
A	325	SER	-	cloning artifact	UNP O75821
A	326	GLY	-	cloning artifact	UNP O75821

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: Eukaryotic translation initiation factor 3 subunit 4

Chain A: 50% 22% • 25%

G224 S225 S226 G227 S228 S229 G230 P231 N232 R233 R234 A235 D236 D237 N238 V243 F259 R260 P261 F262 G263 S264 I265 S266 Y269 L270 A271 G281 F282 I285 H288 A296 I297 A298 G299 V300 Y305 D306 H307 L308 I309 V312 E313 W314 A315 K316 P317 S318 T319 N320 S321

G322 P323 S324 S325 G326

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 7. Colouring as in section 4.1 above.

• Molecule 1: Eukaryotic translation initiation factor 3 subunit 4

Chain A: 50% 22% • 25%

G224 S225 S226 G227 S228 S229 G230 P231 N232 R233 R234 A235 D236 D237 N238 V243 T244 N245 F259 R260 P261 F262 G263 S264 I265 S266 Y269 L270 K274 I285 H288 A296 I297 A298 G299 V300 S301 Y305 D306 H307 L308 I309 L310 N311 V312 E313 W314 A315 K316 P317 S318 T319

N320 S321 G322 P323 S324 S325 G326

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *torsion angle dynamics, restrainted molecular dynamics.*

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *target function, structures with the lowest energy, structures with the least restraint violations.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	2.0.17
CYANA	refinement	2.0.17

No chemical shift data was provided.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	613	600	600	30±4
All	All	12260	12000	12000	601

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

5 of 110 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:243:VAL:HG22	1:A:312:VAL:HG22	0.88	1.45	15	20
1:A:246:LEU:HD22	1:A:250:THR:HG21	0.85	1.46	1	6
1:A:241:ILE:HG22	1:A:297:ILE:HD11	0.80	1.53	13	3
1:A:296:ALA:O	1:A:300:VAL:HG22	0.79	1.76	9	10
1:A:287:PHE:HB2	1:A:293:ALA:HB2	0.75	1.59	4	9

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	77/103 (75%)	67±2 (87±3%)	8±2 (11±2%)	2±1 (2±1%)	11 52
All	All	1540/2060 (75%)	1340 (87%)	169 (11%)	31 (2%)	11 52

All 5 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	281	GLY	14
1	A	297	ILE	11
1	A	307	HIS	3
1	A	280	LYS	2
1	A	273	ASP	1

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	63/83 (76%)	60±1 (94±2%)	4±1 (6±2%)	25 74
All	All	1260/1660 (76%)	1190 (94%)	70 (6%)	25 74

5 of 19 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	305	TYR	18
1	A	309	ILE	10
1	A	260	ARG	8
1	A	251	ARG	6
1	A	274	LYS	3

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

No chemical shift data were provided