



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 24, 2022 – 12:10 pm GMT

PDB ID : 5CZ2  
Title : Crystal structure of a two-domain fragment of MMTV integrase  
Authors : Cook, N.; Ballandras-Colas, A.; Engelman, A.; Cherepanov, P.  
Deposited on : 2015-07-31  
Resolution : 2.72 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.27  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.27


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3359 (2.74-2.70)
Clashscore	141614	3686 (2.74-2.70)
Ramachandran outliers	138981	3622 (2.74-2.70)
Sidechain outliers	138945	3623 (2.74-2.70)
RSRZ outliers	127900	3276 (2.74-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	210	<p>3% 61% 11% 28%</p>
1	B	210	<p>3% 59% 9% 32%</p>
1	C	210	<p>3% 62% 10% 28%</p>
1	D	210	<p>6% 59% 7% 34%</p>
1	E	210	<p>5% 67% 6% 27%</p>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	210	<p>10% 49% 6% 45%</p>
1	G	210	<p>20% 80%</p>
1	H	210	<p>2% 19% 81%</p>
1	I	210	<p>1% 19% 81%</p>
1	J	210	<p>3% 19% 80%</p>
1	K	210	<p>3% 17% 83%</p>
1	L	210	<p>1% 19% 80%</p>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8591 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pol polyprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	151	1214	775	225	211	3	0	0	0
1	B	142	1150	737	209	201	3	0	0	0
1	C	152	1207	770	224	210	3	0	0	0
1	D	139	1098	701	200	194	3	0	0	0
1	E	153	1215	774	227	211	3	0	0	0
1	F	115	895	574	163	155	3	0	0	0
1	G	41	323	200	65	56	2	0	0	0
1	H	40	291	181	57	51	2	0	0	0
1	I	39	282	176	56	48	2	0	0	0
1	J	41	315	195	64	54	2	0	0	0
1	K	36	270	167	56	45	2	0	0	0
1	L	41	319	198	65	54	2	0	0	0

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		
2	B	1	Total	Mg	0	0
			1	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	1	Total 1	Mg 1	0	0
2	D	1	Total 1	Mg 1	0	0
2	E	1	Total 1	Mg 1	0	0
2	F	1	Total 1	Mg 1	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	G	1	Total 1	Zn 1	0	0
3	H	1	Total 1	Zn 1	0	0
3	I	1	Total 1	Zn 1	0	0
3	J	1	Total 1	Zn 1	0	0
3	K	1	Total 1	Zn 1	0	0
3	L	1	Total 1	Zn 1	0	0







THR  
ALA  
ALA  
GLU  
ARG  
HIS  
TRP  
GLY  
PRO

● Molecule 1: Pol polyprotein

Chain L: 19% 80%

**A1** **R20** **T26** **L36** **P41** ASP TRP GLY HIS ALA PRO GLN LEU GLY VAL ASN PRO ARG GLY LEU LYS PRO ARG VAL TRP TRP TRP GLN MET ASP VAL THR HIS VAL SER VAL SER GLU PHE GLY LYS LEU LYS TYR VAL HIS VAL THR VAL ASP THR TYR SER SER HIS ILE ILE PHE LEU LYS LEU LYS

ALA ARG THR GLY GLU ALA THR LYS ASP VAL GLN HIS LEU ALA GLN SER PHE ALA TYR MET GLY ILE LYS PRO GLN LYS ILE THR ASP ASN ALA PRO TYR VAL SER ARG SER ILE GLN PHE LEU ALA ARG TRP LYS ILE SER HIS VAL THR TYR ILE HIS VAL THR TYR ASN PRO GLN

GLY GLN ALA ILE VAL GLU ARG THR LYS HIS GLN ASN ILE LYS ALA GLN LEU ASN LYS LEU GLN LYS ALA GLN ALA HIS LEU PHE VAL LEU ASN MET ASP ASN GLN GLY HIS THR ALA ALA ARG THR THR ALA ARG GLU ILE TRP GLY TRP GLY PRO

## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	54.37Å 83.15Å 141.14Å 90.00° 90.19° 90.00°	Depositor
Resolution (Å)	70.57 – 2.72 70.57 – 2.72	Depositor EDS
% Data completeness (in resolution range)	99.3 (70.57-2.72) 98.3 (70.57-2.72)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.14 (at 2.73Å)	Xtrriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, $R_{free}$	0.241 , 0.280 0.239 , 0.275	Depositor DCC
$R_{free}$ test set	1655 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	54.4	Xtrriage
Anisotropy	0.082	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.37$ , $\langle L^2 \rangle = 0.20$	Xtrriage
Estimated twinning fraction	0.116 for h,-k,-l	Xtrriage
Reported twinning fraction	0.865 for H, K, L 0.135 for -h,-k,l	Depositor
Outliers	0 of 33771 reflections	Xtrriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	8591	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	67.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.86 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.2301e-03.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.23	0/1247	0.39	0/1692
1	B	0.23	0/1181	0.39	0/1602
1	C	0.23	0/1239	0.39	0/1682
1	D	0.23	0/1128	0.38	0/1535
1	E	0.23	0/1247	0.39	0/1693
1	F	0.22	0/916	0.38	0/1245
1	G	0.22	0/330	0.37	0/446
1	H	0.23	0/298	0.36	0/407
1	I	0.23	0/289	0.36	0/395
1	J	0.22	0/322	0.37	0/437
1	K	0.23	0/276	0.36	0/374
1	L	0.22	0/326	0.37	0/441
All	All	0.23	0/8799	0.38	0/11949

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1214	0	1189	12	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1150	0	1117	10	0
1	C	1207	0	1183	12	0
1	D	1098	0	1032	9	0
1	E	1215	0	1187	6	0
1	F	895	0	847	6	0
1	G	323	0	313	0	0
1	H	291	0	255	0	0
1	I	282	0	249	0	0
1	J	315	0	298	2	0
1	K	270	0	235	0	0
1	L	319	0	309	2	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	G	1	0	0	0	0
3	H	1	0	0	0	0
3	I	1	0	0	0	0
3	J	1	0	0	0	0
3	K	1	0	0	0	0
3	L	1	0	0	0	0
All	All	8591	0	8214	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 53 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:115:ILE:HD12	1:E:141:ILE:HD11	1.79	0.65
1:A:115:ILE:HD12	1:A:141:ILE:HD11	1.82	0.61
1:C:115:ILE:HD12	1:C:141:ILE:HD11	1.83	0.61
1:E:54:ARG:NH2	1:E:85:TYR:O	2.40	0.55
1:A:143:HIS:O	1:L:20:ARG:NH2	2.39	0.54

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	147/210 (70%)	142 (97%)	5 (3%)	0	100	100
1	B	136/210 (65%)	134 (98%)	2 (2%)	0	100	100
1	C	148/210 (70%)	144 (97%)	4 (3%)	0	100	100
1	D	133/210 (63%)	131 (98%)	2 (2%)	0	100	100
1	E	149/210 (71%)	144 (97%)	5 (3%)	0	100	100
1	F	105/210 (50%)	104 (99%)	1 (1%)	0	100	100
1	G	39/210 (19%)	38 (97%)	1 (3%)	0	100	100
1	H	38/210 (18%)	37 (97%)	1 (3%)	0	100	100
1	I	37/210 (18%)	36 (97%)	1 (3%)	0	100	100
1	J	39/210 (19%)	38 (97%)	1 (3%)	0	100	100
1	K	32/210 (15%)	31 (97%)	1 (3%)	0	100	100
1	L	39/210 (19%)	38 (97%)	1 (3%)	0	100	100
All	All	1042/2520 (41%)	1017 (98%)	25 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	126/176 (72%)	122 (97%)	4 (3%)	39	67
1	B	120/176 (68%)	118 (98%)	2 (2%)	60	83

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	125/176 (71%)	120 (96%)	5 (4%)	31	58
1	D	111/176 (63%)	109 (98%)	2 (2%)	59	82
1	E	125/176 (71%)	122 (98%)	3 (2%)	49	76
1	F	90/176 (51%)	87 (97%)	3 (3%)	38	66
1	G	33/176 (19%)	33 (100%)	0	100	100
1	H	27/176 (15%)	27 (100%)	0	100	100
1	I	26/176 (15%)	26 (100%)	0	100	100
1	J	31/176 (18%)	31 (100%)	0	100	100
1	K	26/176 (15%)	26 (100%)	0	100	100
1	L	32/176 (18%)	32 (100%)	0	100	100
All	All	872/2112 (41%)	853 (98%)	19 (2%)	52	78

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	172	GLN
1	F	173	LYS
1	F	183	LEU
1	F	172	GLN
1	C	172	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	104	GLN
1	I	39	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 12 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	151/210 (71%)	0.44	7 (4%) 32 31	30, 53, 100, 129	0
1	B	142/210 (67%)	0.55	7 (4%) 29 28	32, 57, 87, 102	0
1	C	152/210 (72%)	0.48	6 (3%) 39 39	33, 57, 91, 106	0
1	D	139/210 (66%)	0.74	13 (9%) 8 7	41, 72, 112, 155	0
1	E	153/210 (72%)	0.59	10 (6%) 18 18	35, 61, 105, 137	0
1	F	115/210 (54%)	1.06	20 (17%) 1 1	52, 74, 116, 150	0
1	G	41/210 (19%)	0.37	1 (2%) 59 60	44, 53, 81, 120	0
1	H	40/210 (19%)	1.02	4 (10%) 7 5	52, 83, 129, 166	0
1	I	39/210 (18%)	0.83	2 (5%) 28 27	50, 70, 115, 133	0
1	J	41/210 (19%)	0.96	6 (14%) 2 1	65, 79, 107, 133	0
1	K	36/210 (17%)	0.92	6 (16%) 1 1	72, 96, 113, 131	0
1	L	41/210 (19%)	0.17	1 (2%) 59 60	40, 57, 81, 91	0
All	All	1090/2520 (43%)	0.64	83 (7%) 13 12	30, 64, 108, 166	0

The worst 5 of 83 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	174	ALA	5.9
1	I	34	VAL	5.5
1	F	179	THR	5.3
1	K	34	VAL	5.1
1	H	16	ALA	5.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MG	B	301	1/1	0.78	0.11	52,52,52,52	0
2	MG	D	301	1/1	0.80	0.16	81,81,81,81	0
2	MG	C	301	1/1	0.81	0.25	65,65,65,65	0
2	MG	A	301	1/1	0.87	0.17	52,52,52,52	0
2	MG	F	301	1/1	0.88	0.05	53,53,53,53	0
3	ZN	G	301	1/1	0.88	0.13	55,55,55,55	0
3	ZN	I	301	1/1	0.89	0.10	83,83,83,83	0
2	MG	E	301	1/1	0.90	0.20	46,46,46,46	0
3	ZN	H	301	1/1	0.93	0.11	86,86,86,86	0
3	ZN	L	301	1/1	0.93	0.17	55,55,55,55	0
3	ZN	J	301	1/1	0.94	0.08	57,57,57,57	0
3	ZN	K	301	1/1	0.97	0.07	64,64,64,64	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.