Full wwPDB X-ray Structure Validation Report (i)

Sep 26, 2023 – 11:38 PM EDT

PDB ID	:	6DBF
Title	:	Crystal Structure of VHH R303 in complex with InlB-LRR
Authors	:	Brooks, C.L.; Toride King, M.; Huh, I.
Deposited on	:	2018-05-03
Resolution	:	1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1483 (1.56-1.56)
Clashscore	141614	1529 (1.56-1.56)
Ramachandran outliers	138981	1498 (1.56-1.56)
Sidechain outliers	138945	1495 (1.56-1.56)
RSRZ outliers	127900	1465 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	229	82% 8% • 9%
2	B	142	% 79% 5% • 15%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 5565 atoms, of which 2580 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Listeria virulence factor InlB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
1	A	208	3373	1068	1704	270	331	0	11	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	21	MET	-	initiating methionine	UNP Q45GD4
A	22	GLY	-	expression tag	UNP Q45GD4
A	23	SER	-	expression tag	UNP Q45GD4
A	24	HIS	-	expression tag	UNP Q45GD4
A	25	HIS	-	expression tag	UNP Q45GD4
A	26	HIS	-	expression tag	UNP Q45GD4
A	27	HIS	-	expression tag	UNP Q45GD4
A	28	HIS	-	expression tag	UNP Q45GD4
A	29	HIS	-	expression tag	UNP Q45GD4
A	30	GLU	-	expression tag	UNP Q45GD4
A	31	ASN	-	expression tag	UNP Q45GD4
A	32	LEU	-	expression tag	UNP Q45GD4
A	33	TYR	-	expression tag	UNP Q45GD4
A	34	PHE	-	expression tag	UNP Q45GD4
A	35	GLN	-	expression tag	UNP Q45GD4
A	36	GLY	-	expression tag	UNP Q45GD4
A	118	VAL	THR	conflict	UNP Q45GD4
A	243	ALA	CYS	conflict	UNP Q45GD4

- Molecule 2 is a protein called InlB specific VHH R303.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	B	121	1810	580	876	164	184	6	0	5	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	255	255	255	0	0
			Total	O		
3	B	127	127	127	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Listeria virulence factor InlB

Chain A:



- Molecule 2: InlB specific VHH R303

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α, β, γ	82.96Å 82.96Å 64.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	58.66 – 1.55 58.66 – 1.55	Depositor EDS
% Data completeness (in resolution range)	98.2 (58.66-1.55) 98.2 (58.66-1.55)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	7.77 (at 1.55Å)	Xtriage
Refinement program	PHENIX 1.12_2829	Depositor
R, R_{free}	0.138 , 0.163 0.138 , 0.163	Depositor DCC
R_{free} test set	3156 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å^2)	17.7	Xtriage
Anisotropy	0.124	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.44 , 59.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.028 for h,-k,-l	Xtriage
F_o,F_c correlation	0.98	EDS
Total number of atoms	5565	wwPDB-VP
Average B, all atoms (Å^2)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.53% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.76	3/1723 (0.2%)	0.82	0/2343
2	B	0.77	1/963 (0.1%)	0.85	2/1306 (0.2%)
All	All	0.76	4/2686 (0.1%)	0.83	2/3649 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	195	GLU	CB-CG	-9.59	1.33	1.52
2	B	102	CYS	CB-SG	7.88	1.95	1.82
1	A	195	GLU	CG-CD	6.40	1.61	1.51
1	A	118	VAL	CB-CG2	-5.57	1.41	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	68	PHE	CB-CG-CD1	6.19	125.13	120.80
2	B	68	PHE	CB-CG-CD2	-5.86	116.70	120.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1669	1704	1716	14	0
2	B	934	876	881	4	2

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	255	0	0	7	2
3	B	127	0	0	1	2
All	All	2985	2580	2597	18	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:174[B]:ASN:ND2	1:A:196:ASP:OD2	2.13	0.81
1:A:115:LYS:O	1:A:118:VAL:HG23	1.80	0.81
1:A:37:GLU:N	3:A:303:HOH:O	2.15	0.78
1:A:174[B]:ASN:OD1	3:A:301:HOH:O	2.02	0.77
2:B:115:GLN:NE2	3:B:202:HOH:O	2.24	0.68
1:A:80[B]:ASP:OD1	3:A:302:HOH:O	2.14	0.65
1:A:96[B]:GLN:NE2	3:A:307:HOH:O	2.27	0.64
1:A:224:LEU:HD11	1:A:238:LEU:HD13	1.86	0.56
1:A:174[B]:ASN:ND2	3:A:308:HOH:O	2.28	0.54
2:B:33:CYS:CB	2:B:102:CYS:SG	2.97	0.51
1:A:159[A]:ASN:CG	3:A:306:HOH:O	2.50	0.50
2:B:12:VAL:HG11	2:B:86:LEU:HD13	1.95	0.48
1:A:56:GLU:OE1	3:A:304:HOH:O	2.21	0.45
1:A:98:LEU:HB3	1:A:101[B]:VAL:HG23	1.99	0.44
2:B:60:TYR:O	2:B:65:ARG:NH1	2.51	0.44
1:A:82:ILE:HD11	1:A:98:LEU:CD2	2.47	0.43
1:A:243:ALA:O	1:A:244:LEU:CB	2.66	0.43
1:A:173:GLY:HA2	1:A:195:GLU:O	2.21	0.41

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:84:ASN:HD21	2:B:115:GLN:OE1[4_565]	1.41	0.19
3:B:241:HOH:O	3:B:301:HOH:O[3_654]	2.05	0.15
3:A:412:HOH:O	3:A:491:HOH:O[3_664]	2.09	0.11
3:A:454:HOH:O	3:B:322:HOH:O[2_565]	2.14	0.06
2:B:84:ASN:ND2	2:B:115:GLN:OE1[4_565]	2.17	0.03

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	217/229 (95%)	208 (96%)	9 (4%)	0	100	100
2	B	122/142 (86%)	120 (98%)	2 (2%)	0	100	100
All	All	339/371 (91%)	328 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	195/211 (92%)	193 (99%)	2 (1%)	76	57
2	B	100/120 (83%)	99 (99%)	1 (1%)	76	57
All	All	295/331 (89%)	292 (99%)	3 (1%)	73	57

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	143	LYS
1	A	225	ARG
2	B	115	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	208/229 (90%)	-0.42	1 (0%)	91	93	12, 19, 37, 59	0
2	B	121/142 (85%)	-0.30	1 (0%)	86	89	13, 23, 44, 54	0
All	All	329/371 (88%)	-0.37	2 (0%)	89	92	12, 21, 40, 59	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	40	VAL	3.1
1	A	243	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.