



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 21, 2020 – 04:35 AM BST

PDB ID : 6EUP
Title : Crystal structure of Neisseria meningitidis NadA variant 3 double mutant A33I-I38L
Authors : Dello Iacono, L.; Liguori, A.; Malito, E.; Bottomley, M.J.
Deposited on : 2017-10-31
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	156	 85% 8% • 6%
1	B	156	 85% 8% • 6%
1	C	156	 86% 8% 6%

2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 3513 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adhesin A.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
1	A	147	1098	677	181	240	0	0	0
1	B	147	1098	677	181	240	0	0	0
1	C	147	1098	677	181	240	0	0	0

There are 33 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	33	ILE	ALA	engineered mutation	UNP Q8KH85
A	38	LEU	ILE	engineered mutation	UNP Q8KH85
A	171	ALA	-	expression tag	UNP Q8KH85
A	172	SER	-	expression tag	UNP Q8KH85
A	173	LYS	-	expression tag	UNP Q8KH85
A	174	HIS	-	expression tag	UNP Q8KH85
A	175	HIS	-	expression tag	UNP Q8KH85
A	176	HIS	-	expression tag	UNP Q8KH85
A	177	HIS	-	expression tag	UNP Q8KH85
A	178	HIS	-	expression tag	UNP Q8KH85
A	179	HIS	-	expression tag	UNP Q8KH85
B	33	ILE	ALA	engineered mutation	UNP Q8KH85
B	38	LEU	ILE	engineered mutation	UNP Q8KH85
B	171	ALA	-	expression tag	UNP Q8KH85
B	172	SER	-	expression tag	UNP Q8KH85
B	173	LYS	-	expression tag	UNP Q8KH85
B	174	HIS	-	expression tag	UNP Q8KH85
B	175	HIS	-	expression tag	UNP Q8KH85
B	176	HIS	-	expression tag	UNP Q8KH85
B	177	HIS	-	expression tag	UNP Q8KH85
B	178	HIS	-	expression tag	UNP Q8KH85
B	179	HIS	-	expression tag	UNP Q8KH85
C	33	ILE	ALA	engineered mutation	UNP Q8KH85

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	38	LEU	ILE	engineered mutation	UNP Q8KH85
C	171	ALA	-	expression tag	UNP Q8KH85
C	172	SER	-	expression tag	UNP Q8KH85
C	173	LYS	-	expression tag	UNP Q8KH85
C	174	HIS	-	expression tag	UNP Q8KH85
C	175	HIS	-	expression tag	UNP Q8KH85
C	176	HIS	-	expression tag	UNP Q8KH85
C	177	HIS	-	expression tag	UNP Q8KH85
C	178	HIS	-	expression tag	UNP Q8KH85
C	179	HIS	-	expression tag	UNP Q8KH85

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C₆H₁₄O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 8 6 2	0	0
4	A	1	Total C O 8 6 2	0	0
4	B	1	Total C O 8 6 2	0	0
4	C	1	Total C O 8 6 2	0	0
4	C	1	Total C O 8 6 2	0	0

- Molecule 5 is HEXAETHYLENE GLYCOL (three-letter code: P6G) (formula: C₁₂H₂₆O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			19	12	7		
5	B	1	Total	C	O	0	0
			19	12	7		

- Molecule 6 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	O	P	0	0
			5	4	1		
6	B	1	Total	O	P	0	0
			5	4	1		
6	C	1	Total	O	P	0	0
			5	4	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	22	Total	O	0	0
			22	22		
7	B	16	Total	O	0	0
			16	16		
7	C	15	Total	O	0	0
			15	15		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Adhesin A



- Molecule 1: Adhesin A



- Molecule 1: Adhesin A



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	68.92Å 39.75Å 255.63Å 90.00° 95.13° 90.00°	Depositor
Resolution (Å)	63.65 – 2.65 63.65 – 2.65	Depositor EDS
% Data completeness (in resolution range)	88.9 (63.65-2.65) 88.6 (63.65-2.65)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.40 (at 2.65Å)	Xtrriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.198 , 0.236 0.208 , 0.251	Depositor DCC
R_{free} test set	995 reflections (5.45%)	wwPDB-VP
Wilson B-factor (Å ²)	33.5	Xtrriage
Anisotropy	0.537	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 39.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.447 for $1/2^*h-3/2^*k,-1/2^*h-1/2^*k,-1/2^*h+1/2^*k-l$ 0.448 for $1/2^*h+3/2^*k,1/2^*h-1/2^*k,-1/2^*h-1/2^*k-l$	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3513	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, EDO, PO4, P6G, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/1103	0.61	0/1491
1	B	0.48	0/1103	0.60	0/1491
1	C	0.48	0/1103	0.60	0/1491
All	All	0.48	0/3309	0.60	0/4473

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1098	0	1100	10	0
1	B	1098	0	1100	12	0
1	C	1098	0	1100	7	0
2	A	1	0	0	0	0
3	A	24	0	36	2	0
3	B	20	0	30	0	0
3	C	28	0	42	0	0
4	A	16	0	28	2	0
4	B	8	0	14	1	0
4	C	16	0	28	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	38	0	52	6	0
6	B	10	0	0	0	0
6	C	5	0	0	0	0
7	A	22	0	0	0	0
7	B	16	0	0	0	0
7	C	15	0	0	0	0
All	All	3513	0	3530	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:81:PHE:HB3	1:C:89:VAL:HG21	1.75	0.69
1:A:81:PHE:HB3	1:A:89:VAL:HG21	1.75	0.69
1:B:81:PHE:HB3	1:B:89:VAL:HG21	1.75	0.68
1:A:149:ASN:HB3	4:A:209:MPD:H12	1.78	0.66
1:A:151:THR:HB	3:A:207:EDO:H11	1.77	0.66
1:B:142:ALA:HB1	5:B:207:P6G:H112	1.78	0.65
1:B:146:LEU:HG	1:B:150:ILE:HD11	1.89	0.55
1:B:141:ASN:HD22	5:B:208:P6G:H142	1.72	0.54
1:B:148:GLU:HG3	4:B:206:MPD:H13	1.93	0.51
1:A:153:PHE:CZ	1:C:154:ALA:HA	2.46	0.49
1:A:122:LEU:HD21	1:B:121:LYS:HB3	1.93	0.49
1:B:141:ASN:HD22	5:B:208:P6G:H122	1.78	0.49
4:A:208:MPD:H32	1:C:158:LYS:HG3	1.95	0.48
1:A:154:ALA:HA	1:B:153:PHE:CZ	2.49	0.47
1:B:145:LYS:HB3	5:B:207:P6G:H92	1.96	0.47
1:B:142:ALA:HB2	5:B:208:P6G:H171	1.96	0.46
1:A:81:PHE:HE2	1:C:87:LYS:HG2	1.82	0.45
1:B:141:ASN:ND2	5:B:208:P6G:H122	2.33	0.44
1:C:149:ASN:HB3	4:C:208:MPD:H31	2.01	0.43
1:A:121:LYS:HB3	1:C:122:LEU:HD21	2.01	0.43
1:A:87:LYS:HG2	1:B:81:PHE:HE2	1.84	0.42
1:C:147:GLY:HA3	4:C:209:MPD:H4	2.03	0.40
1:A:152:THR:OG1	3:A:207:EDO:H12	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	145/156 (93%)	143 (99%)	2 (1%)	0	100	100
1	B	145/156 (93%)	144 (99%)	1 (1%)	0	100	100
1	C	145/156 (93%)	143 (99%)	2 (1%)	0	100	100
All	All	435/468 (93%)	430 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	116/124 (94%)	111 (96%)	5 (4%)	29	44
1	B	116/124 (94%)	111 (96%)	5 (4%)	29	44
1	C	116/124 (94%)	111 (96%)	5 (4%)	29	44
All	All	348/372 (94%)	333 (96%)	15 (4%)	29	44

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	ASP
1	A	67	THR
1	A	70	ASP
1	A	121	LYS
1	A	150	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	29	ASP
1	B	43	ASN
1	B	67	THR
1	B	116	GLU
1	B	121	LYS
1	C	28	ASP
1	C	29	ASP
1	C	67	THR
1	C	70	ASP
1	C	112	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	141	ASN
1	C	149	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 29 ligands modelled in this entry, 1 is monoatomic - leaving 28 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	A	203	-	3,3,3	0.54	0	2,2,2	0.41	0
4	MPD	C	208	-	7,7,7	0.63	0	9,10,10	0.47	0
4	MPD	A	208	-	7,7,7	0.84	1 (14%)	9,10,10	0.61	0
3	EDO	C	205	-	3,3,3	0.62	0	2,2,2	0.29	0
4	MPD	B	206	-	7,7,7	0.98	1 (14%)	9,10,10	0.76	0
3	EDO	A	202	-	3,3,3	0.67	0	2,2,2	0.29	0
3	EDO	B	202	-	3,3,3	0.63	0	2,2,2	0.34	0
3	EDO	C	203	-	3,3,3	0.57	0	2,2,2	0.47	0
3	EDO	A	204	-	3,3,3	0.60	0	2,2,2	0.34	0
3	EDO	C	201	-	3,3,3	0.68	0	2,2,2	0.15	0
5	P6G	B	208	-	18,18,18	0.57	0	17,17,17	0.41	0
5	P6G	B	207	-	18,18,18	0.57	0	17,17,17	0.44	0
3	EDO	C	202	-	3,3,3	0.70	0	2,2,2	0.11	0
4	MPD	A	209	-	7,7,7	1.02	1 (14%)	9,10,10	0.67	0
6	PO4	C	210	-	4,4,4	2.47	1 (25%)	6,6,6	0.53	0
6	PO4	B	210	-	4,4,4	2.48	1 (25%)	6,6,6	0.56	0
3	EDO	C	204	-	3,3,3	0.61	0	2,2,2	0.33	0
3	EDO	B	203	-	3,3,3	0.67	0	2,2,2	0.16	0
6	PO4	B	209	-	4,4,4	2.53	1 (25%)	6,6,6	0.64	0
3	EDO	C	206	-	3,3,3	0.89	0	2,2,2	0.13	0
3	EDO	A	206	-	3,3,3	0.61	0	2,2,2	0.22	0
3	EDO	A	207	-	3,3,3	0.91	0	2,2,2	0.64	0
3	EDO	B	201	-	3,3,3	0.55	0	2,2,2	0.37	0
3	EDO	B	205	-	3,3,3	0.57	0	2,2,2	0.33	0
3	EDO	C	207	-	3,3,3	0.68	0	2,2,2	0.24	0
3	EDO	B	204	-	3,3,3	0.81	0	2,2,2	0.08	0
4	MPD	C	209	-	7,7,7	0.65	0	9,10,10	0.51	0
3	EDO	A	205	-	3,3,3	0.56	0	2,2,2	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	203	-	-	0/1/1/1	-
4	MPD	C	208	-	-	3/5/5/5	-
4	MPD	A	208	-	-	3/5/5/5	-
3	EDO	C	205	-	-	0/1/1/1	-
4	MPD	B	206	-	-	3/5/5/5	-
3	EDO	A	202	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	202	-	-	1/1/1/1	-
3	EDO	C	203	-	-	1/1/1/1	-
3	EDO	A	204	-	-	0/1/1/1	-
3	EDO	C	201	-	-	0/1/1/1	-
5	P6G	B	208	-	-	12/16/16/16	-
5	P6G	B	207	-	-	10/16/16/16	-
3	EDO	C	202	-	-	0/1/1/1	-
4	MPD	A	209	-	-	0/5/5/5	-
3	EDO	C	204	-	-	0/1/1/1	-
3	EDO	B	203	-	-	0/1/1/1	-
3	EDO	C	206	-	-	0/1/1/1	-
3	EDO	A	206	-	-	0/1/1/1	-
3	EDO	A	207	-	-	1/1/1/1	-
3	EDO	B	201	-	-	0/1/1/1	-
3	EDO	B	205	-	-	0/1/1/1	-
3	EDO	C	207	-	-	1/1/1/1	-
3	EDO	B	204	-	-	0/1/1/1	-
4	MPD	C	209	-	-	1/5/5/5	-
3	EDO	A	205	-	-	1/1/1/1	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	209	PO4	P-O1	4.19	1.60	1.50
6	C	210	PO4	P-O1	4.07	1.60	1.50
6	B	210	PO4	P-O1	4.06	1.60	1.50
4	B	206	MPD	C3-C2	2.25	1.59	1.53
4	A	209	MPD	C3-C2	2.24	1.59	1.53
4	A	208	MPD	C3-C2	2.02	1.59	1.53

There are no bond angle outliers.

There are no chirality outliers.

All (37) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	208	MPD	C1-C2-C3-C4
4	A	208	MPD	O2-C2-C3-C4
5	B	207	P6G	O10-C11-C12-O13
5	B	208	P6G	O13-C14-C15-O16
5	B	207	P6G	O4-C5-C6-O7
5	B	208	P6G	O4-C5-C6-O7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	B	208	P6G	O7-C8-C9-O10
5	B	207	P6G	O1-C2-C3-O4
3	C	203	EDO	O1-C1-C2-O2
5	B	207	P6G	O7-C8-C9-O10
3	A	207	EDO	O1-C1-C2-O2
4	C	208	MPD	O2-C2-C3-C4
4	C	209	MPD	O2-C2-C3-C4
5	B	208	P6G	O1-C2-C3-O4
5	B	208	P6G	C11-C12-O13-C14
5	B	208	P6G	C14-C15-O16-C17
5	B	208	P6G	C5-C6-O7-C8
5	B	208	P6G	C9-C8-O7-C6
5	B	207	P6G	C8-C9-O10-C11
5	B	207	P6G	C18-C17-O16-C15
5	B	208	P6G	C2-C3-O4-C5
5	B	207	P6G	C6-C5-O4-C3
4	C	208	MPD	C2-C3-C4-C5
4	C	208	MPD	C2-C3-C4-O4
4	A	208	MPD	CM-C2-C3-C4
4	B	206	MPD	CM-C2-C3-C4
5	B	208	P6G	C8-C9-O10-C11
5	B	208	P6G	C12-C11-O10-C9
4	B	206	MPD	O2-C2-C3-C4
5	B	207	P6G	O16-C17-C18-O19
3	B	202	EDO	O1-C1-C2-O2
3	C	207	EDO	O1-C1-C2-O2
3	A	205	EDO	O1-C1-C2-O2
5	B	208	P6G	O16-C17-C18-O19
5	B	207	P6G	C9-C8-O7-C6
4	B	206	MPD	C2-C3-C4-C5
5	B	207	P6G	C14-C15-O16-C17

There are no ring outliers.

8 monomers are involved in 13 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	208	MPD	1	0
4	A	208	MPD	1	0
4	B	206	MPD	1	0
5	B	208	P6G	4	0
5	B	207	P6G	2	0
4	A	209	MPD	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	207	EDO	2	0
4	C	209	MPD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	147/156 (94%)	0.02	1 (0%) 87 87	11, 61, 112, 139	0
1	B	147/156 (94%)	0.15	4 (2%) 54 50	10, 62, 110, 136	0
1	C	147/156 (94%)	0.10	3 (2%) 65 60	13, 61, 118, 141	0
All	All	441/468 (94%)	0.09	8 (1%) 68 65	10, 62, 113, 141	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	108	VAL	4.4
1	C	63	ASP	2.9
1	B	61	ASP	2.9
1	C	58	TYR	2.7
1	B	67	THR	2.4
1	C	60	ILE	2.3
1	A	30	VAL	2.3
1	B	64	GLY	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	C	202	4/4	0.67	0.24	86,87,88,88	0
3	EDO	A	206	4/4	0.72	0.24	100,100,100,100	0
3	EDO	C	204	4/4	0.73	0.27	73,73,76,77	0
3	EDO	C	206	4/4	0.79	0.14	68,70,71,71	0
3	EDO	B	205	4/4	0.81	0.28	73,74,76,76	0
3	EDO	A	205	4/4	0.82	0.20	62,63,63,64	0
3	EDO	B	204	4/4	0.83	0.20	59,63,67,67	0
5	P6G	B	208	19/19	0.85	0.35	72,97,101,101	0
5	P6G	B	207	19/19	0.87	0.29	37,63,71,74	0
3	EDO	A	204	4/4	0.87	0.15	89,90,90,91	0
6	PO4	C	210	5/5	0.87	0.20	111,111,111,111	0
3	EDO	A	202	4/4	0.88	0.14	61,63,66,67	0
3	EDO	C	207	4/4	0.88	0.23	71,73,73,73	0
3	EDO	A	203	4/4	0.89	0.18	45,46,49,52	0
3	EDO	A	207	4/4	0.89	0.24	24,25,29,30	0
4	MPD	C	208	8/8	0.90	0.24	42,49,59,60	0
3	EDO	B	203	4/4	0.90	0.20	82,82,83,83	0
4	MPD	B	206	8/8	0.90	0.19	41,50,53,55	0
3	EDO	C	201	4/4	0.91	0.15	58,60,61,62	0
6	PO4	B	209	5/5	0.91	0.16	90,91,92,93	0
3	EDO	C	205	4/4	0.91	0.23	61,62,64,64	0
4	MPD	A	208	8/8	0.92	0.22	76,80,80,82	0
4	MPD	C	209	8/8	0.93	0.17	49,52,57,58	0
4	MPD	A	209	8/8	0.93	0.16	50,54,60,62	0
3	EDO	B	201	4/4	0.95	0.16	60,60,62,62	0
3	EDO	B	202	4/4	0.95	0.13	56,56,57,57	0
3	EDO	C	203	4/4	0.96	0.23	44,47,49,49	0
6	PO4	B	210	5/5	0.96	0.17	108,109,109,110	0
2	CL	A	201	1/1	0.99	0.09	44,44,44,44	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around P6G B 208:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.