



# Full wwPDB NMR Structure Validation Report ⓘ

Oct 23, 2021 – 05:15 PM EDT

PDB ID : 1F68  
Title : NMR SOLUTION STRUCTURE OF THE BROMODOMAIN FROM HUMAN GCN5  
Authors : Wright, P.E.; Hudson, B.P.; Dyson, H.J.  
Deposited on : 2000-06-20

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : 2.23.2  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	103	 93% 5%

## 2 Ensemble composition and analysis

This entry contains 30 models. The atoms present in the NMR models are not consistent. Some calculations may have failed as a result. All residues are included in the validation scores. Model 4 is the overall representative, medoid model (most similar to other models). The authors have identified model 21 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:732-A:829 (98)	0.32	4

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 6 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 5, 6, 7, 9, 10, 11, 12, 13, 15, 16, 18, 21, 22, 23, 25, 26, 28, 29
2	17, 30
Single-model clusters	8; 14; 19; 20; 24; 27

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1724 atoms, of which 861 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called HISTONE ACETYLTRANSFERASE.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	103	1724	558	861	145	155	5	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	730	GLY	PRO	engineered mutation	UNP Q92830

## 4 Residue-property plots

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  93% 5%



### 4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

#### 4.2.1 Score per residue for model 1

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  91% 5%



#### 4.2.2 Score per residue for model 2

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  92% 5%



### 4.2.3 Score per residue for model 3

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  92% • 5%



### 4.2.4 Score per residue for model 4 (medoid)

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  91% • 5%



### 4.2.5 Score per residue for model 5

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  91% • 5%



### 4.2.6 Score per residue for model 6

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  88% 7% 5%



### 4.2.7 Score per residue for model 7

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  90% 5% 5%



#### 4.2.8 Score per residue for model 8

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  91% 5%



#### 4.2.9 Score per residue for model 9

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  90% 5% 5%



#### 4.2.10 Score per residue for model 10

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  90% 5% 5%



#### 4.2.11 Score per residue for model 11

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  90% 5%



#### 4.2.12 Score per residue for model 12

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  89% 6% 5%



#### 4.2.13 Score per residue for model 13

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  90% . . 5%



#### 4.2.14 Score per residue for model 14

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  89% 6% 5%



#### 4.2.15 Score per residue for model 15

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  89% 6% 5%



#### 4.2.16 Score per residue for model 16

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  91% . 5%



#### 4.2.17 Score per residue for model 17

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  89% 5% . 5%



#### 4.2.18 Score per residue for model 18

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  87% 8% 5%



#### 4.2.19 Score per residue for model 19

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  88% 7% 5%



#### 4.2.20 Score per residue for model 20

- Molecule 1: HISTONE ACETYLTRANSFERASE


Chain A:  90% 5% 5%



#### 4.2.21 Score per residue for model 21

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  87% 8% 5%



#### 4.2.22 Score per residue for model 22

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  87% 8% 5%



#### 4.2.23 Score per residue for model 23

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  89% 6% 5%



#### 4.2.24 Score per residue for model 24

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  89% 6% 5%



#### 4.2.25 Score per residue for model 25

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  91% 5%



#### 4.2.26 Score per residue for model 26

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  87% 7% 5%



#### 4.2.27 Score per residue for model 27

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  88% 7% 5%



#### 4.2.28 Score per residue for model 28

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  91% • 5%



#### 4.2.29 Score per residue for model 29

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  88% 7% 5%



#### 4.2.30 Score per residue for model 30

- Molecule 1: HISTONE ACETYLTRANSFERASE

Chain A:  91% • 5%



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *Torsion angle dynamics (DYANA) for generation of initial structures; Structure-mediated assignment of ambiguous experimental restraints (SANE); Iteration between DYANA and SANE; Simulated Annealing and Energy Minimization in AMBER using full restraint set.*

Of the 185 calculated structures, 30 were deposited, based on the following criterion: *NO DISTANCE OR ANGLE VIOLATIONS GREATER THAN 0.15 A OR 5 DEGREES.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DYANA	refinement	1.5
SANE	refinement	1.0
Amber	structure solution	6.0

No chemical shift data was provided.

## 6 Model quality i

### 6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.63±0.00	0±0/852 ( 0.0± 0.0%)	0.83±0.02	0±1/1154 ( 0.0± 0.0%)
All	All	0.63	0/25560 ( 0.0%)	0.83	12/34620 ( 0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	2.7±1.2
All	All	0	80

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	770	ARG	NE-CZ-NH2	-5.83	117.39	120.30	17	4
1	A	826	TYR	CB-CG-CD2	-5.60	117.64	121.00	18	1
1	A	816	ARG	NE-CZ-NH2	-5.53	117.53	120.30	4	2
1	A	785	ARG	NE-CZ-NH1	5.42	123.01	120.30	21	2
1	A	799	ARG	NE-CZ-NH2	-5.25	117.67	120.30	24	2
1	A	785	ARG	NE-CZ-NH2	-5.02	117.79	120.30	29	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	766	TYR	Sidechain	18
1	A	787	TYR	Sidechain	16

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Group	Models (Total)
1	A	786	TYR	Sidechain	11
1	A	814	TYR	Sidechain	10
1	A	781	ARG	Sidechain	6
1	A	807	TYR	Sidechain	4
1	A	734	TYR	Sidechain	3
1	A	826	TYR	Sidechain	3
1	A	753	PHE	Sidechain	2
1	A	799	ARG	Sidechain	2
1	A	770	ARG	Sidechain	2
1	A	805	ARG	Sidechain	1
1	A	790	ARG	Sidechain	1
1	A	825	PHE	Sidechain	1

## 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
All	All	24840	24960	24960	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	98/103 (95%)	92±2 (93±2%)	6±2 (6±2%)	0±0 (0±0%)	54 85
All	All	2940/3090 (95%)	2745 (93%)	191 (6%)	4 (0%)	54 85

All 3 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	757	VAL	2
1	A	749	SER	1
1	A	770	ARG	1

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	91/94 (97%)	89±1 (97±1%)	2±1 (3±1%)	49	91
All	All	2730/2820 (97%)	2658 (97%)	72 (3%)	49	91

All 17 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	749	SER	11
1	A	770	ARG	7
1	A	776	LYS	7
1	A	785	ARG	6
1	A	791	LYS	6
1	A	828	LYS	5
1	A	774	ASP	5
1	A	796	ASP	4
1	A	823	LYS	4
1	A	777	THR	4
1	A	773	ILE	4
1	A	797	LEU	3
1	A	780	GLU	2
1	A	783	ARG	1
1	A	738	LYS	1
1	A	767	GLU	1
1	A	806	GLU	1

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 6.7 Other polymers [i](#)

There are no such molecules in this entry.

### 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

No chemical shift data were provided