



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 22, 2020 – 03:15 pm BST

PDB ID : 5GNR  
Title : the structure of mini-MFN1 K88A in complex with GDP  
Authors : Yan, L.; Yu, C.; Ming, Z.; Lou, Z.; Rao, Z.; Hu, J.  
Deposited on : 2016-07-24  
Resolution : 2.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	421	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3116 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitofusin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	387	3069	1944	527	584	14	0	0	0

There are 58 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	88	ALA	LYS	engineered mutation	UNP Q8IWA4
A	365	GLY	-	expression tag	UNP Q8IWA4
A	366	SER	-	expression tag	UNP Q8IWA4
A	367	GLY	-	expression tag	UNP Q8IWA4
A	368	SER	-	expression tag	UNP Q8IWA4
A	369	GLY	-	expression tag	UNP Q8IWA4
A	370	SER	-	expression tag	UNP Q8IWA4
A	371	GLY	-	expression tag	UNP Q8IWA4
A	372	GLY	-	expression tag	UNP Q8IWA4
A	373	SER	-	expression tag	UNP Q8IWA4
A	374	GLU	-	expression tag	UNP Q8IWA4
A	375	ILE	-	expression tag	UNP Q8IWA4
A	376	ALA	-	expression tag	UNP Q8IWA4
A	377	ARG	-	expression tag	UNP Q8IWA4
A	378	LEU	-	expression tag	UNP Q8IWA4
A	379	PRO	-	expression tag	UNP Q8IWA4
A	380	LYS	-	expression tag	UNP Q8IWA4
A	381	GLU	-	expression tag	UNP Q8IWA4
A	382	ILE	-	expression tag	UNP Q8IWA4
A	383	ASP	-	expression tag	UNP Q8IWA4
A	384	GLN	-	expression tag	UNP Q8IWA4
A	385	LEU	-	expression tag	UNP Q8IWA4
A	386	GLU	-	expression tag	UNP Q8IWA4
A	387	LYS	-	expression tag	UNP Q8IWA4
A	388	ILE	-	expression tag	UNP Q8IWA4
A	389	GLN	-	expression tag	UNP Q8IWA4
A	390	ASN	-	expression tag	UNP Q8IWA4

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	391	ASN	-	expression tag	UNP Q8IWA4
A	392	SER	-	expression tag	UNP Q8IWA4
A	393	LYS	-	expression tag	UNP Q8IWA4
A	394	LEU	-	expression tag	UNP Q8IWA4
A	395	LEU	-	expression tag	UNP Q8IWA4
A	396	ARG	-	expression tag	UNP Q8IWA4
A	397	ASN	-	expression tag	UNP Q8IWA4
A	398	LYS	-	expression tag	UNP Q8IWA4
A	399	ALA	-	expression tag	UNP Q8IWA4
A	400	VAL	-	expression tag	UNP Q8IWA4
A	401	GLN	-	expression tag	UNP Q8IWA4
A	402	LEU	-	expression tag	UNP Q8IWA4
A	403	GLU	-	expression tag	UNP Q8IWA4
A	404	ASN	-	expression tag	UNP Q8IWA4
A	405	GLU	-	expression tag	UNP Q8IWA4
A	406	LEU	-	expression tag	UNP Q8IWA4
A	407	GLU	-	expression tag	UNP Q8IWA4
A	408	ASN	-	expression tag	UNP Q8IWA4
A	409	PHE	-	expression tag	UNP Q8IWA4
A	410	THR	-	expression tag	UNP Q8IWA4
A	411	LYS	-	expression tag	UNP Q8IWA4
A	412	GLN	-	expression tag	UNP Q8IWA4
A	413	PHE	-	expression tag	UNP Q8IWA4
A	414	LEU	-	expression tag	UNP Q8IWA4
A	415	PRO	-	expression tag	UNP Q8IWA4
A	416	SER	-	expression tag	UNP Q8IWA4
A	417	SER	-	expression tag	UNP Q8IWA4
A	418	ASN	-	expression tag	UNP Q8IWA4
A	419	GLU	-	expression tag	UNP Q8IWA4
A	420	GLU	-	expression tag	UNP Q8IWA4
A	421	SER	-	expression tag	UNP Q8IWA4

- Molecule 2 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula:  $C_{10}H_{15}N_5O_{11}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	28	10	5	11	2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	19	Total	O	0	0
			19	19		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mitofusin-1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.34Å 74.97Å 94.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.96 – 2.65 39.96 – 2.65	Depositor EDS
% Data completeness (in resolution range)	51.4 (39.96-2.65) 89.9 (39.96-2.65)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.70 (at 2.65Å)	Xtrriage
Refinement program	PHENIX 1.8.4_1496	Depositor
R, $R_{free}$	0.216 , 0.302 0.224 , 0.298	Depositor DCC
$R_{free}$ test set	1429 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.6	Xtrriage
Anisotropy	0.060	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 42.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.028 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	3116	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.21% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.57	0/3119	0.88	18/4198 (0.4%)

There are no bond length outliers.

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	268	LEU	CB-CA-C	-11.42	88.51	110.20
1	A	191	TRP	CB-CA-C	-9.88	90.65	110.40
1	A	104	GLY	N-CA-C	9.50	136.86	113.10
1	A	210	SER	CB-CA-C	-8.76	93.45	110.10
1	A	193	ASP	CB-CA-C	8.11	126.62	110.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3069	0	3066	144	0
2	A	28	0	12	1	0
3	A	19	0	0	5	0
All	All	3116	0	3078	144	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 144 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:102:PRO:HG2	1:A:105:ILE:CD1	1.41	1.49
1:A:136:VAL:HG13	1:A:137:LYS:N	1.54	1.15
1:A:142:LEU:HD12	1:A:143:ALA:N	1.62	1.15
1:A:136:VAL:CG1	1:A:137:LYS:H	1.58	1.12
1:A:102:PRO:HG2	1:A:105:ILE:HD11	1.24	1.08

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	377/421 (90%)	344 (91%)	28 (7%)	5 (1%)	12 18

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	136	VAL
1	A	379	PRO
1	A	193	ASP
1	A	304	VAL
1	A	414	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	339/367 (92%)	320 (94%)	19 (6%)	21	33

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	139	VAL
1	A	142	LEU
1	A	269	LYS
1	A	138	THR
1	A	293	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	140	ASN
1	A	218	HIS
1	A	256	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GDP	A	1000	-	24,30,30	1.26	2 (8%)	31,47,47	2.09	8 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GDP	A	1000	-	-	3/12/32/32	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1000	GDP	C6-C5	4.23	1.48	1.41
2	A	1000	GDP	C5-C4	2.48	1.47	1.40

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1000	GDP	C2-N3-C4	4.94	121.00	115.36
2	A	1000	GDP	C6-C5-C4	-4.65	116.36	120.80
2	A	1000	GDP	C6-N1-C2	4.42	122.95	115.93
2	A	1000	GDP	C5-C6-N1	-4.00	117.96	123.43
2	A	1000	GDP	N3-C2-N1	-3.81	122.14	127.22

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1000	GDP	PA-O3A-PB-O2B
2	A	1000	GDP	PA-O3A-PB-O1B
2	A	1000	GDP	O4'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1000	GDP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	387/421 (91%)	0.10	19 (4%) 29 26	17, 36, 72, 128	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	191	TRP	8.9
1	A	154	ALA	7.9
1	A	155	GLY	6.7
1	A	297	GLN	4.3
1	A	190	SER	4.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.


### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GDP	A	1000	28/28	0.97	0.13	20,35,41,43	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.