



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 8, 2024 – 09:08 am GMT

PDB ID : 6H9B  
Title : 1,1-Diheterocyclic Ethylenes Derived from Quinaldine and Carbazole as New Tubulin Polymerization Inhibitors: Synthesis, Metabolism, and Biological Evaluation  
Authors : Varela, P.F.; Gigant, B.  
Deposited on : 2018-08-03  
Resolution : 2.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1235 (2.78-2.74)
Clashscore	141614	1277 (2.78-2.74)
Ramachandran outliers	138981	1257 (2.78-2.74)
Sidechain outliers	138945	1257 (2.78-2.74)
RSRZ outliers	127900	1207 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	433	 9% 89% 9%
1	C	433	 7% 86% 12%
2	B	431	 12% 82% 17%
2	D	431	 2% 83% 15%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	E	142	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SO4	A	502	-	-	-	X

## 2 Entry composition i

There are 9 unique types of molecules in this entry. The entry contains 15148 atoms, of which 40 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	431	3380	2141	574	643	22	0	0	0
1	C	433	3393	2149	576	646	22	0	0	0

- Molecule 2 is a protein called Tubulin beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	431	3393	2128	580	658	27	0	0	0
2	D	427	3356	2105	572	652	27	0	0	0

- Molecule 3 is a protein called Stathmin-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	E	136	1125	691	209	221	4	0	0	0

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
E	4	ALA	SER	conflict	UNP P63043
E	14	ALA	CYS	engineered mutation	UNP P63043
E	20	TRP	PHE	engineered mutation	UNP P63043

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	C	1	Total O S 5 4 1	0	0
4	C	1	Total O S 5 4 1	0	0
4	D	1	Total O S 5 4 1	0	0
4	D	1	Total O S 5 4 1	0	0
4	D	1	Total O S 5 4 1	0	0
4	E	1	Total O S 5 4 1	0	0

- Molecule 5 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula:  $C_{10}H_{16}N_5O_{14}P_3$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
5	A	1	32	10	5	14	3	0	0
5	C	1	32	10	5	14	3	0	0

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
6	A	1	1	1	0	0
6	C	1	1	1	0	0

- Molecule 7 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>11</sub>P<sub>2</sub>).



*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	N		
8	D	1	47	25	20	2	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	51	Total	O	0	0
			51	51		
9	B	43	Total	O	0	0
			43	43		
9	C	57	Total	O	0	0
			57	57		
9	D	64	Total	O	0	0
			64	64		
9	E	15	Total	O	0	0
			15	15		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tubulin alpha chain




- Molecule 1: Tubulin alpha chain





- Molecule 2: Tubulin beta chain







• Molecule 2: Tubulin beta chain



• Molecule 3: Stathmin-4



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	64.94Å 128.48Å 251.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.65 – 2.75 47.65 – 2.74	Depositor EDS
% Data completeness (in resolution range)	99.7 (47.65-2.75) 99.7 (47.65-2.74)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.04 (at 2.73Å)	Xtrriage
Refinement program	BUSTER	Depositor
R, $R_{free}$	0.172 , 0.216 0.189 , 0.235	Depositor DCC
$R_{free}$ test set	2789 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	80.7	Xtrriage
Anisotropy	0.521	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 93.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	15148	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	108.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.58% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, GDP, FWH, MG, GTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.48	0/3457	0.64	0/4693
1	C	0.52	0/3470	0.68	0/4711
2	B	0.54	0/3468	0.73	0/4697
2	D	0.55	0/3429	0.73	2/4643 (0.0%)
3	E	0.51	0/1136	0.65	0/1510
All	All	0.52	0/14960	0.69	2/20254 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	247	GLN	C-N-CA	8.51	142.98	121.70
2	D	249	ASN	N-CA-C	-5.67	95.68	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3380	0	3290	21	0
1	C	3393	0	3304	36	0
2	B	3393	0	3270	46	0
2	D	3356	0	3234	32	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	1125	0	1134	12	0
4	A	10	0	0	0	0
4	B	15	0	0	0	0
4	C	10	0	0	0	0
4	D	15	0	0	0	0
4	E	5	0	0	0	0
5	A	32	0	12	0	0
5	C	32	0	12	1	0
6	A	1	0	0	0	0
6	C	1	0	0	0	0
7	B	28	0	12	0	0
7	D	28	0	12	0	0
8	B	27	20	0	2	0
8	D	27	20	0	3	0
9	A	51	0	0	2	0
9	B	43	0	0	0	0
9	C	57	0	0	0	0
9	D	64	0	0	1	0
9	E	15	0	0	0	0
All	All	15108	40	14280	136	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 136 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:244:PHE:HB2	1:C:356:ASN:HD21	1.03	1.10
2:B:282:GLN:HB2	2:B:283:TYR:HA	1.48	0.94
1:C:244:PHE:HB2	1:C:356:ASN:ND2	1.87	0.88
1:A:181:VAL:HG13	8:B:505:FWH:C22	2.14	0.77
2:B:98:GLY:H	1:C:2:ARG:NH1	1.85	0.74

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	427/433 (99%)	412 (96%)	10 (2%)	5 (1%)	13	23
1	C	429/433 (99%)	402 (94%)	21 (5%)	6 (1%)	11	19
2	B	429/431 (100%)	410 (96%)	16 (4%)	3 (1%)	22	39
2	D	423/431 (98%)	411 (97%)	10 (2%)	2 (0%)	29	47
3	E	132/142 (93%)	125 (95%)	4 (3%)	3 (2%)	6	10
All	All	1840/1870 (98%)	1760 (96%)	61 (3%)	19 (1%)	15	27

5 of 19 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	247	GLN
2	B	250	ALA
2	B	286	LEU
1	C	281	ALA
1	C	349	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	365/367 (100%)	356 (98%)	9 (2%)	47	67
1	C	367/367 (100%)	352 (96%)	15 (4%)	30	50
2	B	374/374 (100%)	359 (96%)	15 (4%)	31	51
2	D	371/374 (99%)	357 (96%)	14 (4%)	33	53

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	E	120/125 (96%)	113 (94%)	7 (6%)	20	35
All	All	1597/1607 (99%)	1537 (96%)	60 (4%)	33	53

5 of 60 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	178	SER
3	E	25	LYS
1	C	338	LYS
3	E	15	THR
3	E	125	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	193	GLN
2	D	247	GLN
3	E	71	HIS
2	D	294	GLN
1	C	249	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 2 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	SO4	C	504	-	4,4,4	0.19	0	6,6,6	0.13	0
5	GTP	A	503	6	26,34,34	0.75	1 (3%)	32,54,54	0.98	2 (6%)
4	SO4	B	502	-	4,4,4	0.21	0	6,6,6	0.17	0
4	SO4	C	501	-	4,4,4	0.22	0	6,6,6	0.09	0
8	FWH	B	505	-	28,31,31	0.81	1 (3%)	41,46,46	1.09	4 (9%)
7	GDP	B	504	-	24,30,30	0.85	2 (8%)	30,47,47	0.94	2 (6%)
4	SO4	D	502	-	4,4,4	0.12	0	6,6,6	0.28	0
5	GTP	C	502	6	26,34,34	0.82	2 (7%)	32,54,54	0.80	0
4	SO4	B	501	-	4,4,4	0.08	0	6,6,6	0.22	0
4	SO4	B	503	-	4,4,4	0.17	0	6,6,6	0.31	0
4	SO4	A	501	-	4,4,4	0.19	0	6,6,6	0.25	0
4	SO4	E	201	-	4,4,4	0.13	0	6,6,6	0.15	0
7	GDP	D	504	-	24,30,30	0.87	1 (4%)	30,47,47	0.78	1 (3%)
4	SO4	D	503	-	4,4,4	0.18	0	6,6,6	0.56	0
8	FWH	D	505	-	28,31,31	0.78	0	41,46,46	1.04	4 (9%)
4	SO4	A	502	-	4,4,4	0.13	0	6,6,6	0.15	0
4	SO4	D	501	-	4,4,4	0.29	0	6,6,6	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GTP	A	503	6	-	5/18/38/38	0/3/3/3
8	FWH	B	505	-	-	1/8/8/8	0/5/5/5
7	GDP	B	504	-	-	5/12/32/32	0/3/3/3
5	GTP	C	502	6	-	8/18/38/38	0/3/3/3
7	GDP	D	504	-	-	5/12/32/32	0/3/3/3
8	FWH	D	505	-	-	1/8/8/8	0/5/5/5

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	B	505	FWH	C25-N2	-2.52	1.45	1.48
7	B	504	GDP	C8-N7	-2.16	1.31	1.35
7	D	504	GDP	C8-N7	-2.08	1.31	1.35
5	A	503	GTP	C8-N7	-2.06	1.31	1.35
5	C	502	GTP	C8-N7	-2.06	1.31	1.35

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	B	505	FWH	C9-C16-C11	-4.00	113.34	118.32
8	D	505	FWH	C9-C16-C11	-3.45	114.02	118.32
8	B	505	FWH	C9-C16-C24	3.35	125.58	120.22
8	D	505	FWH	C9-C16-C24	3.05	125.10	120.22
8	D	505	FWH	C22-C18-N2	-2.73	128.99	132.29

There are no chirality outliers.

5 of 25 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	503	GTP	C5'-O5'-PA-O1A
5	A	503	GTP	C5'-O5'-PA-O2A
5	C	502	GTP	PB-O3B-PG-O3G
5	C	502	GTP	C5'-O5'-PA-O1A
5	C	502	GTP	C5'-O5'-PA-O2A


There are no ring outliers.


3 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	B	505	FWH	2	0
5	C	502	GTP	1	0
8	D	505	FWH	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	C	1
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	C	38:SER	C	46:ASP	N	10.49
1	A	38:SER	C	46:ASP	N	10.40

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	431/433 (99%)	0.46	37 (8%) 10 13	79, 107, 150, 186	0
1	C	433/433 (100%)	0.35	29 (6%) 17 21	71, 104, 147, 182	0
2	B	431/431 (100%)	0.61	53 (12%) 4 4	75, 102, 159, 191	0
2	D	427/431 (99%)	0.23	10 (2%) 60 69	67, 93, 141, 171	0
3	E	136/142 (95%)	1.25	33 (24%) 0 0	97, 132, 190, 200	0
All	All	1858/1870 (99%)	0.48	162 (8%) 10 12	67, 105, 155, 200	0

The worst 5 of 162 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	E	4	ALA	14.5
1	A	282	TYR	7.9
3	E	33	PRO	7.7
1	A	130	THR	7.2
1	A	283	HIS	7.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	SO4	C	501	5/5	0.68	0.32	213,213,214,215	0
4	SO4	D	501	5/5	0.68	0.24	201,201,202,203	0
4	SO4	A	502	5/5	0.78	0.42	215,216,216,217	0
4	SO4	B	503	5/5	0.79	0.40	177,178,178,180	0
4	SO4	E	201	5/5	0.79	0.33	175,177,177,181	0
4	SO4	D	502	5/5	0.87	0.20	170,172,173,174	0
4	SO4	C	504	5/5	0.88	0.13	184,188,191,192	0
4	SO4	B	502	5/5	0.88	0.22	181,182,183,188	0
4	SO4	A	501	5/5	0.89	0.24	179,179,180,182	0
4	SO4	D	503	5/5	0.91	0.19	146,152,154,156	0
4	SO4	B	501	5/5	0.91	0.11	143,149,150,155	0
6	MG	A	504	1/1	0.93	0.13	86,86,86,86	0
8	FWH	D	505	27/27	0.93	0.20	19,51,63,69	0
8	FWH	B	505	27/27	0.96	0.20	71,80,86,88	0
7	GDP	B	504	28/28	0.97	0.21	85,94,101,104	0
5	GTP	C	502	32/32	0.97	0.20	85,92,98,100	0
5	GTP	A	503	32/32	0.97	0.20	89,96,107,111	0
6	MG	C	503	1/1	0.98	0.10	99,99,99,99	0
7	GDP	D	504	28/28	0.98	0.15	71,82,92,95	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around FWH D 505:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around FWH B 505:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







**Electron density around GDP B 504:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around GTP C 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.