wwPDB X-ray Structure Validation Summary Report (i)

Aug 8, 2020 – 08:46 PM BST

PDB ID	:	6I96
Title	:	Structure of the ferrioxamine B transporter FoxA from Pseudomonas aeruginosa in complex with ferrioxamine B
Authors	:	Josts, I.; Tidow, H.
Deposited on	:	2018-11-22
Resolution	:	1.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	677	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	0UE	A	901	X	-	-	-

2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 5945 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ferric hydroxamate uptake.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	677	5325	3345	910	1059	11	0	0	0

- Molecule 2 is Ferrioxamine B (three-letter code: 0UE) (formula: $C_{25}H_{45}FeN_6O_8$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	40	25	1	6	8	0	0

- Molecule 3 is nonyl beta-D-glucopyranoside (three-letter code: BNG) (formula: $C_{15}H_{30}O_6$).

BNG



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	12	10	2	0	0
			Total	C	O		
3	A	1	21	15	6	0	0
			Total	C	O		
3	A	1	21	15	6	0	0
			Total	C	O		
3	A	1	21	15	6	0	0
			Total	C	O		
3	A	1	21	15	6	0	0
			Total	C	O		
3	A	1	21	15	6	0	0
			Total	C	O		
3	A	1	21	15	6	0	0
			Total	C	O		
3	A	1	21	15	6	0	0
			Total	C	O		
3	A	1	10	9	1	0	0
			Total	C	O		
3	A	1	21	15	6	0	0
			Total	C	O		
3	A	1	21	15	6	0	0
			Total	C	O		
3	A	1	21	15	6	0	0
			Total	C	O		
3	A	1	21	15	6	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O			
4	A	1	6	3	3		0	0
			Total	C	O			
4	A	1	6	3	3		0	0
			Total	C	O			
4	A	1	6	3	3		0	0
			Total	C	O			
4	A	1	6	3	3		0	0
			Total	C	O			
4	A	1	6	3	3		0	0
			Total	C	O			
4	A	1	6	3	3		0	0
			Total	C	O			
4	A	1	6	3	3		0	0
			Total	C	O			
4	A	1	6	3	3		0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0

- Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	2	Total Na 2 2	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	257	Total O 257 257	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ferric hydroxamate uptake

Chain A:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α, β, γ	94.89Å 94.89Å 177.61Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	82.18 – 1.85 82.18 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.2 (82.18-1.85) 99.2 (82.18-1.85)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.87 (at 1.84Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.181 , 0.219 0.191 , 0.230	Depositor DCC
R_{free} test set	3949 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å^2)	29.6	Xtriage
Anisotropy	0.665	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.36 , 60.7	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.026 for -h,-k,l	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	5945	wwPDB-VP
Average B, all atoms (Å^2)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.39% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, 0UE, BNG, SO4, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	1.36	21/5451 (0.4%)	1.34	57/7397 (0.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	421	ASP	CB-CG	-8.78	1.33	1.51
1	A	671	ARG	CZ-NH1	7.88	1.43	1.33
1	A	333	THR	N-CA	7.78	1.61	1.46
1	A	524	SER	CB-OG	-7.68	1.32	1.42
1	A	145	PHE	CE1-CZ	7.18	1.50	1.37

The worst 5 of 57 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	421	ASP	CB-CG-OD1	-27.14	93.87	118.30
1	A	421	ASP	CB-CG-OD2	19.82	136.13	118.30
1	A	410	ARG	NE-CZ-NH1	13.91	127.25	120.30
1	A	671	ARG	NE-CZ-NH2	-12.74	113.93	120.30
1	A	810	ARG	NE-CZ-NH2	-12.48	114.06	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	144	VAL	Peptide
1	A	145	PHE	Mainchain
1	A	637	GLY	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5325	0	5038	53	0
2	A	40	0	41	10	0
3	A	253	0	368	7	0
4	A	48	0	64	1	0
5	A	20	0	0	0	0
6	A	2	0	0	0	0
7	A	257	0	0	6	0
All	All	5945	0	5511	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 62 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:901:0UE:C31	2:A:901:0UE:N31	1.70	1.51
2:A:901:0UE:N21	2:A:901:0UE:C21	1.81	1.44
2:A:901:0UE:C11	2:A:901:0UE:N11	1.85	1.39
1:A:757:ARG:HH22	3:A:913:BNG:H2	1.39	0.86
2:A:901:0UE:C19	2:A:901:0UE:C21	2.52	0.85

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	675/677 (100%)	652 (97%)	22 (3%)	1 (0%)	51 36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	145	PHE

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	566/566 (100%)	534 (94%)	32 (6%)	20 6

5 of 32 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	419	ARG
1	A	514	PHE
1	A	768	ILE
1	A	499	THR
1	A	577	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	387	ASN
1	A	682	GLN
1	A	390	HIS
1	A	385	HIS
1	A	653	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 28 ligands modelled in this entry, 2 are monoatomic - leaving 26 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	GOL	A	919	-	5,5,5	0.27	0	5,5,5	0.55	0
3	BNG	A	908	-	21,21,21	0.62	0	26,26,26	0.99	2 (7%)
3	BNG	A	902	-	11,11,21	0.57	0	9,10,26	0.21	0
4	GOL	A	921	-	5,5,5	0.85	0	5,5,5	1.62	2 (40%)
3	BNG	A	904	-	21,21,21	0.67	0	26,26,26	1.11	1 (3%)
4	GOL	A	922	-	5,5,5	0.73	0	5,5,5	0.85	0
5	SO4	A	924	-	4,4,4	1.31	1 (25%)	6,6,6	0.68	0
4	GOL	A	920	-	5,5,5	1.05	0	5,5,5	0.88	0
4	GOL	A	918	-	5,5,5	0.36	0	5,5,5	0.74	0
4	GOL	A	916	-	5,5,5	0.64	0	5,5,5	1.51	0
3	BNG	A	911	-	21,21,21	1.12	1 (4%)	26,26,26	1.44	3 (11%)
3	BNG	A	906	-	21,21,21	0.76	1 (4%)	26,26,26	1.51	4 (15%)
2	0UE	A	901	-	44,44,44	10.11	26 (59%)	44,66,66	8.69	27 (61%)
3	BNG	A	910	-	9,9,21	0.28	0	8,8,26	0.73	0
3	BNG	A	909	-	21,21,21	0.86	1 (4%)	26,26,26	1.30	3 (11%)
3	BNG	A	913	-	21,21,21	1.45	3 (14%)	26,26,26	3.24	7 (26%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	SO4	A	926	-	4,4,4	0.43	0	6,6,6	0.38	0
3	BNG	A	912	-	21,21,21	0.86	1 (4%)	26,26,26	1.36	4 (15%)
5	SO4	A	925	-	4,4,4	0.59	0	6,6,6	0.34	0
3	BNG	A	905	-	21,21,21	0.98	1 (4%)	26,26,26	1.81	6 (23%)
4	GOL	A	917	-	5,5,5	0.24	0	5,5,5	0.65	0
5	SO4	A	923	-	4,4,4	0.59	0	6,6,6	0.48	0
3	BNG	A	914	-	21,21,21	0.54	0	26,26,26	1.20	3 (11%)
3	BNG	A	907	-	21,21,21	0.71	1 (4%)	26,26,26	1.48	4 (15%)
4	GOL	A	915	-	5,5,5	0.39	0	5,5,5	0.81	0
3	BNG	A	903	-	21,21,21	0.98	1 (4%)	26,26,26	1.62	3 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	919	-	-	0/4/4/4	-
3	BNG	A	908	-	-	9/12/32/32	0/1/1/1
3	BNG	A	902	-	-	3/8/9/32	-
4	GOL	A	921	-	-	0/4/4/4	-
3	BNG	A	904	-	-	8/12/32/32	0/1/1/1
4	GOL	A	922	-	-	0/4/4/4	-
4	GOL	A	920	-	-	4/4/4/4	-
4	GOL	A	918	-	-	2/4/4/4	-
4	GOL	A	916	-	-	3/4/4/4	-
3	BNG	A	911	-	-	2/12/32/32	0/1/1/1
3	BNG	A	906	-	-	11/12/32/32	0/1/1/1
2	0UE	A	901	-	1/1/9/11	12/34/88/88	0/3/5/5
3	BNG	A	910	-	-	3/7/7/32	-
3	BNG	A	909	-	-	6/12/32/32	0/1/1/1
3	BNG	A	913	-	-	4/12/32/32	0/1/1/1
3	BNG	A	912	-	-	9/12/32/32	0/1/1/1
3	BNG	A	905	-	-	7/12/32/32	0/1/1/1
4	GOL	A	917	-	-	2/4/4/4	-
3	BNG	A	914	-	-	5/12/32/32	0/1/1/1
3	BNG	A	907	-	-	3/12/32/32	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	915	-	-	0/4/4/4	-
3	BNG	A	903	-	-	5/12/32/32	0/1/1/1

The worst 5 of 37 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	901	0UE	C11-N11	34.07	1.85	1.31
2	A	901	0UE	C21-N21	31.59	1.81	1.31
2	A	901	0UE	C39-N11	-25.61	1.02	1.47
2	A	901	0UE	C19-N21	-23.93	1.05	1.47
2	A	901	0UE	O22-C21	14.53	1.52	1.28

The worst 5 of 69 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	901	0UE	O21-N21-C21	-37.33	92.03	116.42
2	A	901	0UE	O11-N11-C11	-21.34	102.48	116.42
2	A	901	0UE	O22-C21-N21	-19.23	103.71	118.38
2	A	901	0UE	O22-C21-C22	16.72	137.74	120.13
2	A	901	0UE	O12-C11-N11	-13.95	107.73	118.38

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	901	0UE	N31

5 of 98 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	904	BNG	O5-C1-O1-C1'
4	A	920	GOL	O1-C1-C2-C3
4	A	920	GOL	C1-C2-C3-O3
4	A	920	GOL	O2-C2-C3-O3
4	A	918	GOL	C1-C2-C3-O3

There are no ring outliers.

7 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	904	BNG	1	0
4	A	920	GOL	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	911	BNG	1	0
3	A	906	BNG	2	0
2	A	901	0UE	10	0
3	A	913	BNG	2	0
3	A	905	BNG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand BNG A 908



Ligand BNG A 902



Ligand BNG A 904



Ligand BNG A 911



Ligand BNG A 906



Ligand 0UE A 901



Ligand BNG A 910



Ligand BNG A 909



Ligand BNG A 913



Ligand BNG A 912



Ligand BNG A 905



Ligand BNG A 914



Ligand BNG A 907





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	677/677 (100%)	-0.17	6 (0%) 84 84	29, 45, 78, 112	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	614	ALA	4.3
1	A	663	PHE	3.5
1	A	684	SER	3.5
1	A	144	VAL	2.3
1	A	617	THR	2.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	BNG	A	912	21/21	0.42	0.26	67,107,133,135	0
3	BNG	A	905	21/21	0.51	0.29	61,102,119,131	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	BNG	A	909	21/21	0.55	0.26	64,89,115,118	0
3	BNG	A	902	12/21	0.56	0.33	70,81,105,106	0
3	BNG	A	911	21/21	0.57	0.24	55,77,96,103	0
3	BNG	A	904	21/21	0.61	0.31	53,76,98,104	0
3	BNG	A	908	21/21	0.64	0.23	61,87,98,102	0
4	GOL	A	917	6/6	0.66	0.30	77,87,98,102	0
4	GOL	A	920	6/6	0.67	0.14	57,61,66,70	0
3	BNG	A	903	21/21	0.67	0.20	64,80,100,111	0
3	BNG	A	906	21/21	0.68	0.28	61,78,121,127	0
4	GOL	A	922	6/6	0.69	0.20	72,84,86,89	0
3	BNG	A	913	21/21	0.73	0.21	51,77,88,99	0
3	BNG	A	907	21/21	0.76	0.23	60,109,127,129	0
3	BNG	A	910	10/21	0.77	0.15	72,78,90,93	0
5	SO4	A	925	5/5	0.79	0.37	104,105,118,118	0
3	BNG	A	914	21/21	0.81	0.23	65,83,99,102	0
5	SO4	A	923	5/5	0.82	0.22	77,95,110,113	0
4	GOL	A	916	6/6	0.83	0.23	54,64,73,99	0
6	NA	A	928	1/1	0.84	0.21	76,76,76,76	0
5	SO4	A	926	5/5	0.87	0.24	99,113,117,120	0
4	GOL	A	921	6/6	0.90	0.17	47,56,66,68	0
4	GOL	A	919	6/6	0.93	0.10	45,79,89,92	0
4	GOL	A	918	6/6	0.94	0.09	43,49,51,54	0
2	0UE	A	901	40/40	0.95	0.13	37,47,66,68	0
4	GOL	A	915	6/6	0.97	0.09	31,34,36,38	0
6	NA	A	927	1/1	0.98	0.12	42,42,42,42	0
5	SO4	A	924	5/5	0.99	0.07	42,46,56,59	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around BNG A 912:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around BNG A 905:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around BNG A 909:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around BNG A 902:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around BNG A 911:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around BNG A 904:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around BNG A 908:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around BNG A 903:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around BNG A 906:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around BNG A 913:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around BNG A 907:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around BNG A 910:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around BNG A 914:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around 0UE A 901:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative)
and green (positive)







6.5 Other polymers (i)

There are no such residues in this entry.