wwPDB X-ray Structure Validation Summary Report (i)

Aug 7, 2020 – 07:19 AM BST

PDB ID	:	2JG1
Title	:	STRUCTURE OF Staphylococcus aureus D-TAGATOSE-6-PHOSPHATE KINASE with cofactor and substrate
Authors	:	Miallau, L.; Hunter, W.N.; McSweeney, S.M.; Leonard, G.A.
Deposited on	:	2007-02-07
Resolution	:	2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	330	3% 83% 12% • •
1	B	330	6% 85% 11% •
1	C	330	4% 84% 11% • 5%
1	D	330	2% 85% 9% • 5%

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 10894 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TAGATOSE-6-PHOSPHATE KINASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	318	2498	1583	424	482	4	5	0	13	0
1	B	318	2475	1567	418	481	4	5	0	7	0
1	C	315	2451	1552	416	476	3	4	0	7	0
1	D	313	2450	1553	411	478	3	5	0	12	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	124	MSE	LEU	engineered mutation	UNP P0A0B9
A	125	MSE	LEU	engineered mutation	UNP P0A0B9
B	124	MSE	LEU	engineered mutation	UNP P0A0B9
B	125	MSE	LEU	engineered mutation	UNP P0A0B9
C	124	MSE	LEU	engineered mutation	UNP P0A0B9
C	125	MSE	LEU	engineered mutation	UNP P0A0B9
D	124	MSE	LEU	engineered mutation	UNP P0A0B9
D	125	MSE	LEU	engineered mutation	UNP P0A0B9

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
2	B	1	1	1	0	0
2	A	1	1	1	0	0
2	D	1	1	1	0	0
2	C	2	2	2	0	0

- Molecule 3 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: $C_{10}H_{17}N_6O_{12}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	31	10	6	12	3	0	0
			Total	C	N	O	P		
3	B	1	31	10	6	12	3	0	0
			Total	C	N	O	P		
3	C	1	31	10	6	12	3	0	0
			Total	C	N	O	P		
3	D	1	31	10	6	12	3	0	0

- Molecule 4 is 6-O-phosphono-beta-D-tagatofuranose (three-letter code: TA6) (formula: $C_6H_{13}O_9P$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	P		
4	C	1	16	6	9	1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	221	221	221	0	0
			Total	O		
5	B	214	214	214	0	0
			Total	O		
5	C	201	201	201	0	0
			Total	O		
5	D	239	239	239	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: TAGATOSE-6-PHOSPHATE KINASE

Chain A:



- Molecule 1: TAGATOSE-6-PHOSPHATE KINASE

Chain B:



- Molecule 1: TAGATOSE-6-PHOSPHATE KINASE

Chain C:



- Molecule 1: TAGATOSE-6-PHOSPHATE KINASE

Chain D:




P196
E202
W218
S222
K231
I244
L247
T256
I260
T261
A277
N278
M282
Q286
L295
V310

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	95.20Å 97.09Å 154.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.85 – 2.00 24.85 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (24.85-2.00) 99.8 (24.85-2.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.55 (at 1.99Å)	Xtriage
Refinement program	REFMAC 5.3.0011	Depositor
R, R_{free}	0.190 , 0.234 0.189 , 0.230	Depositor DCC
R_{free} test set	4857 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å^2)	28.7	Xtriage
Anisotropy	0.010	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 50.7	EDS
L-test for twinning[2]	$< \|L\| > = 0.47$, $< L^2 > = 0.30$	Xtriage
Estimated twinning fraction	0.029 for k,h,-l	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	10894	wwPDB-VP
Average B, all atoms (Å^2)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.56% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: TA6, ANP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.66	2/2585 (0.1%)	0.56	2/3495 (0.1%)
1	B	0.44	0/2538	0.56	0/3437
1	C	0.45	0/2517	0.61	0/3409
1	D	0.52	2/2540 (0.1%)	0.55	0/3440
All	All	0.53	4/10180 (0.0%)	0.57	2/13781 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	77[A]	LYS	CD-CE	17.30	1.94	1.51
1	A	77[B]	LYS	CD-CE	17.30	1.94	1.51
1	D	202[A]	GLU	CB-CG	8.83	1.69	1.52
1	D	202[B]	GLU	CB-CG	8.83	1.69	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	77[A]	LYS	CD-CE-NZ	-5.16	99.84	111.70
1	A	77[B]	LYS	CD-CE-NZ	-5.16	99.84	111.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2498	0	2519	33	0
1	B	2475	0	2487	17	0
1	C	2451	0	2450	22	0
1	D	2450	0	2453	21	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	2	0	0	0	0
2	D	1	0	0	0	0
3	A	31	0	13	1	0
3	B	31	0	13	2	0
3	C	31	0	13	2	0
3	D	31	0	13	0	0
4	C	16	0	11	0	0
5	A	221	0	0	1	0
5	B	214	0	0	2	0
5	C	201	0	0	2	0
5	D	239	0	0	3	0
All	All	10894	0	9972	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 95 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:77[A]:LYS:CE	1:A:77[A]:LYS:CD	1.94	1.45
1:C:211:PRO:O	1:C:214:GLU:HG3	1.88	0.73
1:D:150[B]:ILE:HD12	1:D:160:VAL:HG11	1.72	0.72
1:C:70:LYS:O	1:C:73[B]:HIS:HD2	1.73	0.71
1:D:142:ASN:HD22	1:D:144:ASP:H	1.41	0.68

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	329/330 (100%)	326 (99%)	2 (1%)	1 (0%)	41	37
1	B	323/330 (98%)	317 (98%)	5 (2%)	1 (0%)	41	37
1	C	320/330 (97%)	311 (97%)	9 (3%)	0	100	100
1	D	323/330 (98%)	318 (98%)	4 (1%)	1 (0%)	41	37
All	All	1295/1320 (98%)	1272 (98%)	20 (2%)	3 (0%)	47	44

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	233	ASN
1	D	-1	SER
1	B	248	ASN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	281/275 (102%)	275 (98%)	6 (2%)	53	57
1	B	276/275 (100%)	270 (98%)	6 (2%)	52	55
1	C	272/275 (99%)	258 (95%)	14 (5%)	24	19
1	D	276/275 (100%)	272 (99%)	4 (1%)	67	72
All	All	1105/1100 (100%)	1075 (97%)	30 (3%)	44	46

5 of 30 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	54	LEU
1	C	119	LYS
1	D	175	ASN
1	C	108	GLU
1	C	137	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	148	GLN
1	C	191	GLN
1	D	142	ASN
1	C	111	ASN
1	D	155	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 10 ligands modelled in this entry, 5 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	ANP	D	1312	2	29,33,33	1.88	8 (27%)	31,52,52	1.85	5 (16%)
3	ANP	A	1312	2	29,33,33	1.95	9 (31%)	31,52,52	1.69	5 (16%)
3	ANP	C	1313	2	29,33,33	1.91	6 (20%)	31,52,52	1.69	5 (16%)
3	ANP	B	1312	2	29,33,33	1.94	9 (31%)	31,52,52	1.50	4 (12%)
4	TA6	C	1312	-	15,16,16	1.28	1 (6%)	17,25,25	0.96	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ANP	D	1312	2	-	2/14/38/38	0/3/3/3
3	ANP	A	1312	2	-	1/14/38/38	0/3/3/3
3	ANP	C	1313	2	-	2/14/38/38	0/3/3/3
3	ANP	B	1312	2	-	1/14/38/38	0/3/3/3
4	TA6	C	1312	-	-	1/9/28/28	0/1/1/1

The worst 5 of 33 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1313	ANP	PG-N3B	5.14	1.76	1.63
3	B	1312	ANP	PG-N3B	4.94	1.76	1.63
3	B	1312	ANP	PB-N3B	4.74	1.75	1.63
3	A	1312	ANP	PB-N3B	4.74	1.75	1.63
3	D	1312	ANP	PG-N3B	4.71	1.75	1.63

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	D	1312	ANP	O1G-PG-N3B	-6.99	101.48	111.77
3	C	1313	ANP	O1G-PG-N3B	-5.94	103.03	111.77
3	A	1312	ANP	O1G-PG-N3B	-5.09	104.28	111.77
3	B	1312	ANP	O1G-PG-N3B	-4.74	104.80	111.77
3	D	1312	ANP	N3-C2-N1	-3.83	122.70	128.68

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	1313	ANP	PB-N3B-PG-O1G
3	C	1313	ANP	PG-N3B-PB-O1B
3	B	1312	ANP	PG-N3B-PB-O1B
3	D	1312	ANP	PG-N3B-PB-O1B
3	A	1312	ANP	PG-N3B-PB-O1B

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1312	ANP	1	0
3	C	1313	ANP	2	0
3	B	1312	ANP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand ANP D 1312



Ligand ANP A 1312



Ligand ANP C 1313



Bond lengths



Bond angles



Torsions



Rings

Ligand ANP B 1312



Ligand TA6 C 1312



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	314/330 (95%)	0.14	10 (3%)	47	46	14, 27, 51, 66	10 (3%)
1	B	314/330 (95%)	0.31	21 (6%)	17	17	16, 28, 55, 70	7 (2%)
1	C	311/330 (94%)	0.19	12 (3%)	39	38	15, 30, 49, 63	6 (1%)
1	D	309/330 (93%)	0.01	7 (2%)	60	59	16, 26, 41, 47	16 (5%)
All	All	1248/1320 (94%)	0.16	50 (4%)	38	37	14, 28, 50, 70	39 (3%)

The worst 5 of 50 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	-7	GLY	6.1
1	B	246	VAL	5.9
1	B	247	LEU	5.0
1	B	194	ASN	4.5
1	B	206	GLN	4.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	MG	B	1311	1/1	0.92	0.07	34,34,34,34	0
3	ANP	B	1312	31/31	0.92	0.14	34,41,47,48	0
3	ANP	C	1313	31/31	0.95	0.12	27,28,30,32	4
3	ANP	A	1312	31/31	0.96	0.11	28,33,38,38	0
2	MG	D	1311	1/1	0.97	0.04	26,26,26,26	0
2	MG	A	1311	1/1	0.97	0.04	27,27,27,27	0
4	TA6	C	1312	16/16	0.97	0.08	24,26,28,29	0
3	ANP	D	1312	31/31	0.97	0.08	19,22,24,26	0
2	MG	C	1310	1/1	0.97	0.07	37,37,37,37	0
2	MG	C	1311	1/1	0.98	0.05	42,42,42,42	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ANP B 1312:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ANP C 1313:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around ANP A 1312:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around TA6 C 1312:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around ANP D 1312:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.