wwPDB NMR Structure Validation Summary Report (i)

Dec 13, 2023 – 08:04 PM EST

PDB ID	:	2JMF
Title	:	Solution structure of the Su(dx) WW4- Notch PY peptide complex
Authors	:	Avis, J.M.; Blankley, R.T.; Jennings, M.D.; Golovanov, A.P.
Deposited on	:	2006-11-05

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	53	49% · 28% 19%
2	B	21	38% 33% 29%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 16 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:523-A:550, B:2325-B:2332 (36)	0.19	16

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 6 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 5, 6, 12, 15, 16
2	8, 10, 11, 14, 18, 20
Single-model clusters	4; 7; 9; 13; 17; 19

3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 893 atoms, of which 434 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called E3 ubiquitin-protein ligase suppressor of deltex.

Mol	Chain	Residues	Atoms					Trace
			Total	C	H	N	O	
1	A	43	670	219	325	63	63	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	505	GLY	-	expression tag	UNP Q9Y0H4
A	506	PRO	-	expression tag	UNP Q9Y0H4
A	507	LEU	-	expression tag	UNP Q9Y0H4
A	508	GLY	-	expression tag	UNP Q9Y0H4
A	509	SER	-	expression tag	UNP Q9Y0H4
A	510	PRO	-	expression tag	UNP Q9Y0H4
A	511	GLU	-	expression tag	UNP Q9Y0H4
A	512	PHE	-	expression tag	UNP Q9Y0H4
A	513	HIS	-	expression tag	UNP Q9Y0H4
A	514	MET	-	expression tag	UNP Q9Y0H4

- Molecule 2 is a protein called Neurogenic locus Notch protein.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
2	B	15	223	70	109	19	24	1	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	2313	GLY	-	expression tag	UNP P07207
B	2314	PRO	-	expression tag	UNP P07207
B	2315	LEU	-	expression tag	UNP P07207
B	2316	GLY	-	expression tag	UNP P07207
B	2317	SER	-	expression tag	UNP P07207

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: E3 ubiquitin-protein ligase suppressor of deltex

Chain A:



• Molecule 2: Neurogenic locus Notch protein

Chain B:



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 16. Colouring as in section 4.1 above.

• Molecule 1: E3 ubiquitin-protein ligase suppressor of deltex

Chain A:



• Molecule 2: Neurogenic locus Notch protein

Chain B:



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing*.

Of the 20 calculated structures, 20 were deposited, based on the following criterion: *all calculated structures submitted*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CARA	structure solution	
CYANA	structure solution	2.1
CYANA	refinement	2.1
NMRView	structure solution	
Amber	refinement	

No chemical shift data was provided.

6 Model quality (i)

6.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.86±0.01	0±0/248 (0.0± 0.0%)	1.10±0.03	1±1/338 (0.3± 0.2%)
2	B	0.79±0.01	0±0/66 (0.0± 0.0%)	0.87±0.02	0±0/91 (0.0± 0.0%)
All	All	0.84	0/6280 (0.0%)	1.06	23/8580 (0.3%)

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})	Models	
								Worst	Total
1	A	530	ARG	NE-CZ-NH1	7.32	123.96	120.30	7	3
1	A	545	ARG	NE-CZ-NH1	7.17	123.89	120.30	4	16
1	A	546	ARG	NE-CZ-NH1	6.93	123.77	120.30	17	2
1	A	537	ARG	NE-CZ-NH1	5.46	123.03	120.30	14	2

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	239	221	222	0±0
All	All	6040	5500	5520	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	28/53 (53%)	27±1 (98±3%)	1±1 (2±3%)	0±0 (0±0%)	100 100
2	B	8/21 (38%)	6±1 (71±8%)	2±1 (26±8%)	0±0 (3±5%)	7 39
All	All	720/1480 (49%)	660 (92%)	55 (8%)	5 (1%)	26 73

All 2 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	2332	ILE	4
2	B	2328	TYR	1

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	24/44 (55%)	22±1 (93±4%)	2±1 (7±4%)	20 68
2	B	8/17 (47%)	8±0 (99±3%)	0±0 (1±3%)	86 97
All	All	640/1220 (52%)	607 (95%)	33 (5%)	27 76

5 of 7 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	523	LEU	20
1	A	530	ARG	4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	529	ILE	3
1	A	537	ARG	2
1	A	542	HIS	2

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

No chemical shift data were provided