



wwPDB X-ray Structure Validation Summary Report ⓘ

May 22, 2020 – 07:20 pm BST

PDB ID : 5K3J
Title : Crystals structure of Acyl-CoA oxidase-2 in Caenorhabditis elegans bound with FAD, ascaroside-CoA, and ATP
Authors : Zhang, X.; Li, K.; Jones, R.A.; Bruner, S.D.; Butcher, R.A.
Deposited on : 2016-05-19
Resolution : 2.68 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3863 (2.70-2.66)
Clashscore	141614	4210 (2.70-2.66)
Ramachandran outliers	138981	4141 (2.70-2.66)
Sidechain outliers	138945	4141 (2.70-2.66)
RSRZ outliers	127900	3780 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	674	<div> <div>%</div> <div> <div></div> <div>76%</div> <div>19%</div> <div>• •</div> </div> </div>
1	B	674	<div> <div>3%</div> <div> <div></div> <div>77%</div> <div>19%</div> <div>• •</div> </div> </div>

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 10979 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acyl-coenzyme A oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	662	Total	C	N	O	S	0	0	0
			5275	3344	930	974	27			
1	B	659	Total	C	N	O	S	0	0	0
			5243	3323	924	969	27			

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	432	ALA	GLU	conflict	UNP O62137
A	662	ALA	-	expression tag	UNP O62137
A	663	ALA	-	expression tag	UNP O62137
A	664	ALA	-	expression tag	UNP O62137
A	665	HIS	-	expression tag	UNP O62137
A	666	HIS	-	expression tag	UNP O62137
A	667	HIS	-	expression tag	UNP O62137
A	668	HIS	-	expression tag	UNP O62137
A	669	HIS	-	expression tag	UNP O62137
A	670	HIS	-	expression tag	UNP O62137
A	671	HIS	-	expression tag	UNP O62137
A	672	HIS	-	expression tag	UNP O62137
A	673	HIS	-	expression tag	UNP O62137
A	674	HIS	-	expression tag	UNP O62137
B	432	ALA	GLU	conflict	UNP O62137
B	662	ALA	-	expression tag	UNP O62137
B	663	ALA	-	expression tag	UNP O62137
B	664	ALA	-	expression tag	UNP O62137
B	665	HIS	-	expression tag	UNP O62137
B	666	HIS	-	expression tag	UNP O62137
B	667	HIS	-	expression tag	UNP O62137
B	668	HIS	-	expression tag	UNP O62137
B	669	HIS	-	expression tag	UNP O62137
B	670	HIS	-	expression tag	UNP O62137
B	671	HIS	-	expression tag	UNP O62137

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	672	HIS	-	expression tag	UNP O62137
B	673	HIS	-	expression tag	UNP O62137
B	674	HIS	-	expression tag	UNP O62137

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 4 is {S}-[2-[3-[(2 {R})-4-[[[(2 {R}),3 {S},4 {R}),5 {R})-5-(6-aminopurin-9-yl)-4-oxidany-3-phosphonooxy-oxolan-2-yl]methoxy-oxidanyl-phosphoryl]oxy-oxidanyl-phosphoryl]oxy-3,3-dimethyl-2-oxidanyl-butanoyl]amino]propanoylamino]ethyl] 5-[(2 {R}),3 {R}),5 {R}),6 {S})-6-methyl-3,5-bis(oxidanyl)oxan-2-yl]oxypentanethioate (three-letter code: 6QA) (formula: C₃₂H₅₄N₇O₂₁P₃S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	S	0	0
			64	32	7	21	3	1		
4	B	1	Total	C	N	O	P	S	0	0
			64	32	7	21	3	1		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	2	Total	Mg	0	0
			2	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	64	Total	O	0	0
			64	64		
6	B	99	Total	O	0	0
			99	99		

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Chain A:

Chain B:

3% 77% 19%

• •

Y593	L594	E595	K596	H625	L639	L645	V648	E649	K650	Y651	L652	K653	P654	H665	HIS	HIS	HIS	HIS	HIS	HIS	HIS	HIS	HIS	HIS	HIS
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	76.04Å 85.57Å 106.90Å 90.00° 91.43° 90.00°	Depositor
Resolution (Å)	39.72 – 2.68 39.72 – 2.68	Depositor EDS
% Data completeness (in resolution range)	99.8 (39.72-2.68) 99.8 (39.72-2.68)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	0.20	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.54 (at 2.69Å)	Xtriage
Refinement program	PHENIX 1.8.4_1496	Depositor
R, R_{free}	0.213 , 0.245 0.227 , 0.263	Depositor DCC
R_{free} test set	1938 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	34.0	Xtriage
Anisotropy	1.297	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 16.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.090 for h,-k,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	10979	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, 6QA, FAD, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/5379	0.47	0/7264
1	B	0.25	0/5347	0.46	0/7221
All	All	0.25	0/10726	0.47	0/14485

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5275	0	5299	83	0
1	B	5243	0	5249	98	0
2	A	53	0	30	3	0
2	B	53	0	30	5	0
3	A	31	0	12	1	0
3	B	31	0	12	3	0
4	A	64	0	0	0	0
4	B	64	0	0	1	0
5	B	2	0	0	0	0
6	A	64	0	0	3	0
6	B	99	0	0	6	0
All	All	10979	0	10632	171	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 171 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:140:ARG:CG	1:B:140:ARG:HH11	1.65	1.06
1:B:289:MET:HB3	1:B:293:ARG:NH2	1.74	1.03
1:B:140:ARG:NH1	1:B:140:ARG:HG2	1.51	1.00
1:A:136:GLU:O	1:A:140:ARG:HG3	1.75	0.86
1:B:470:ASP:O	1:B:471:LYS:HB3	1.77	0.84

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	660/674 (98%)	610 (92%)	34 (5%)	16 (2%)	6	13
1	B	655/674 (97%)	615 (94%)	26 (4%)	14 (2%)	7	16
All	All	1315/1348 (98%)	1225 (93%)	60 (5%)	30 (2%)	6	14

5 of 30 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	187	TRP
1	A	266	ARG
1	A	283	LYS
1	A	580	PHE
1	B	187	TRP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	562/574 (98%)	526 (94%)	36 (6%)	17	36
1	B	557/574 (97%)	522 (94%)	35 (6%)	18	37
All	All	1119/1148 (98%)	1048 (94%)	71 (6%)	18	37

5 of 71 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	580	PHE
1	B	175	VAL
1	B	558	LEU
1	A	588	ASP
1	B	19	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	410	HIS
1	A	435	ASN
1	B	267	HIS
1	B	435	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	B	702	-	51,58,58	4.32	18 (35%)	60,89,89	2.27	11 (18%)
3	ATP	A	702	5	26,33,33	0.92	1 (3%)	31,52,52	1.57	5 (16%)
4	6QA	A	703	-	58,67,67	3.69	18 (31%)	72,98,98	1.59	15 (20%)
3	ATP	B	703	5	26,33,33	0.94	1 (3%)	31,52,52	1.44	4 (12%)
4	6QA	B	705	-	58,67,67	3.71	18 (31%)	72,98,98	1.59	13 (18%)
2	FAD	A	701	-	51,58,58	4.36	18 (35%)	60,89,89	2.27	12 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	B	702	-	-	3/30/50/50	0/6/6/6
3	ATP	A	702	5	-	4/18/38/38	0/3/3/3
4	6QA	A	703	-	-	24/55/91/91	0/4/4/4
3	ATP	B	703	5	-	5/18/38/38	0/3/3/3
4	6QA	B	705	-	-	24/55/91/91	0/4/4/4
2	FAD	A	701	-	-	6/30/50/50	0/6/6/6

The worst 5 of 74 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	702	FAD	C2B-C1B	-16.49	1.28	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	701	FAD	C2B-C1B	-16.48	1.28	1.53
4	B	705	6QA	O34-C35	15.80	1.63	1.41
4	A	703	6QA	O34-C35	15.77	1.63	1.41
4	B	705	6QA	C46-C35	-15.58	1.30	1.53

The worst 5 of 60 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	701	FAD	C5A-C6A-N6A	9.59	134.92	120.35
2	B	702	FAD	C5A-C6A-N6A	9.53	134.84	120.35
2	A	701	FAD	N6A-C6A-N1A	-6.46	105.17	118.57
2	B	702	FAD	N6A-C6A-N1A	-6.45	105.18	118.57
2	B	702	FAD	C4-N3-C2	5.76	120.00	115.14

There are no chirality outliers.

5 of 66 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	702	FAD	C5'-O5'-P-O2P
4	A	703	6QA	C32-O31-P29-O30
4	A	703	6QA	C32-O31-P29-O54
4	A	703	6QA	O59-C03-O04-C05
4	A	703	6QA	C06-C05-O04-C03


There are no ring outliers.

5 monomers are involved in 13 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	702	FAD	5	0
3	A	702	ATP	1	0
3	B	703	ATP	3	0
4	B	705	6QA	1	0
2	A	701	FAD	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the


average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand ATP A 702




Ligand 6QA A 703




Ligand ATP B 703



Ligand 6QA B 705





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	662/674 (98%)	-0.18	9 (1%) 75 76	10, 45, 91, 148	0
1	B	659/674 (97%)	-0.14	18 (2%) 54 54	11, 45, 103, 167	0
All	All	1321/1348 (97%)	-0.16	27 (2%) 65 65	10, 45, 99, 167	0

The worst 5 of 27 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	454	LYS	4.6
1	B	460	LYS	3.9
1	B	496	TRP	3.8
1	B	507	GLU	3.8
1	A	455	ALA	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	6QA	B	705	64/64	0.74	0.35	39,81,96,112	0
5	MG	B	701	1/1	0.88	0.14	44,44,44,44	0
4	6QA	A	703	64/64	0.88	0.23	39,81,96,112	0
3	ATP	B	703	31/31	0.94	0.14	41,52,71,79	0
5	MG	B	704	1/1	0.95	0.12	41,41,41,41	0
3	ATP	A	702	31/31	0.95	0.12	40,44,59,62	0
2	FAD	B	702	53/53	0.95	0.14	25,34,51,54	0
2	FAD	A	701	53/53	0.98	0.13	21,26,46,49	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around 6QA A 703:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ATP B 703:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ATP A 702:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD B 702:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD A 701:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.