wwPDB X-ray Structure Validation Summary Report (i)

Feb 19, 2024 – 12:01 PM EST

PDB ID	:	4KHZ
Title	:	Crystal structure of the maltose-binding protein/maltose transporter complex in an pre-translocation conformation bound to maltoheptaose
Authors	:	Oldham, M.L.; Chen, S.; Chen, J.
Deposited on	:	2013-05-01
Resolution	:	2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	380	2%, 91%, 6% • •
2	F	514	6%, 89%, 5% 6%
3	G	296	4%, 83%, 11% • 5%
4	A	381	4%, 88%, 8% • •
4	B	381	7%, 90%, 7% •

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	C	4	50% 50%
6	D	3	33% 33% 33%

2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 14715 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltose-binding periplasmic protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	E	370	2877	1853	469	549	6	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	371	ALA	-	expression tag	UNP P0AEX9
E	372	SER	-	expression tag	UNP P0AEX9
E	373	ALA	-	expression tag	UNP P0AEX9
E	374	SER	-	expression tag	UNP P0AEX9
E	375	HIS	-	expression tag	UNP P0AEX9
E	376	HIS	-	expression tag	UNP P0AEX9
E	377	HIS	-	expression tag	UNP P0AEX9
E	378	HIS	-	expression tag	UNP P0AEX9
E	379	HIS	-	expression tag	UNP P0AEX9
E	380	HIS	-	expression tag	UNP P0AEX9

- Molecule 2 is a protein called Maltose transport system permease protein MalF.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	F	483	3750	2464	597	672	17	0	0	0

- Molecule 3 is a protein called Binding-protein-dependent transport systems inner membrane component.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	G	282	2182	1461	348	364	9	0	0	0

- Molecule 4 is a protein called Binding-protein-dependent transport systems inner membrane component.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	A	370	2879	1820	515	531	13	0	1	0
4	B	368	2855	1806	511	525	13	0	0	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	372	ALA	-	expression tag	UNP C9QV42
A	373	SER	-	expression tag	UNP C9QV42
A	374	ALA	-	expression tag	UNP C9QV42
A	375	SER	-	expression tag	UNP C9QV42
A	376	HIS	-	expression tag	UNP C9QV42
A	377	HIS	-	expression tag	UNP C9QV42
A	378	HIS	-	expression tag	UNP C9QV42
A	379	HIS	-	expression tag	UNP C9QV42
A	380	HIS	-	expression tag	UNP C9QV42
A	381	HIS	-	expression tag	UNP C9QV42
B	372	ALA	-	expression tag	UNP C9QV42
B	373	SER	-	expression tag	UNP C9QV42
B	374	ALA	-	expression tag	UNP C9QV42
B	375	SER	-	expression tag	UNP C9QV42
B	376	HIS	-	expression tag	UNP C9QV42
B	377	HIS	-	expression tag	UNP C9QV42
B	378	HIS	-	expression tag	UNP C9QV42
B	379	HIS	-	expression tag	UNP C9QV42
B	380	HIS	-	expression tag	UNP C9QV42
B	381	HIS	-	expression tag	UNP C9QV42

- Molecule 5 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
5	C	4	45	24	21	0	0	0

- Molecule 6 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
6	D	3	23	12	11	0	0	1

- Molecule 7 is (1R)-2-{[{[(2S)-2,3-DIHYDROXYPROPYL]OXY}(HYDROXY)PHOSPHORYL]OXY}-1-[(PALMITOYLOXY)METHYL]ETHYL (11E)-OCTADEC-11-ENOATE (three-letter code: PGV) (formula: $C_{40}H_{77}O_{10}P$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	P		
7	F	1	51	40	10	1	0	0

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
8	E	21	21	21	0	0
			Total	O		
8	F	11	11	11	0	0
			Total	O		
8	G	5	5	5	0	0
			Total	O		
8	A	9	9	9	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
8	B	7	7	7	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Maltose-binding periplasmic protein

Chain E:





• Molecule 2: Maltose transport system permease protein MalF

Chain F:



• Molecule 3: Binding-protein-dependent transport systems inner membrane component

Chain G:



• Molecule 4: Binding-protein-dependent transport systems inner membrane component

Chain A:



• Molecule 4: Binding-protein-dependent transport systems inner membrane component

Chain B:





• Molecule 5: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain C:

• Molecule 6: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α, β, γ	76.25Å 92.22Å 117.81Å 90.73° 101.68° 103.60°	Depositor
Resolution (Å)	19.81 – 2.90 19.80 – 2.89	Depositor EDS
% Data completeness (in resolution range)	71.7 (19.81-2.90) 71.5 (19.80-2.89)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.18 (at 2.88Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.243 , 0.283 0.241 , 0.279	Depositor DCC
R_{free} test set	2438 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å^2)	76.1	Xtriage
Anisotropy	0.006	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.28 , 34.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.51$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.91	EDS
Total number of atoms	14715	wwPDB-VP
Average B, all atoms (Å^2)	98.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.02% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, PGV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	E	0.26	0/2946	0.42	0/3998
2	F	0.27	0/3841	0.42	0/5228
3	G	0.27	0/2242	0.44	0/3065
4	A	0.25	0/2929	0.45	0/3972
4	B	0.25	0/2904	0.44	0/3936
All	All	0.26	0/14862	0.43	0/20199

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	2877	0	2859	10	0
2	F	3750	0	3783	14	0
3	G	2182	0	2271	14	0
4	A	2879	0	2940	20	0
4	B	2855	0	2919	14	0
5	C	45	0	39	0	0
6	D	23	0	19	1	0
7	F	51	0	76	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	A	9	0	0	0	0
8	B	7	0	0	0	0
8	E	21	0	0	0	0
8	F	11	0	0	0	0
8	G	5	0	0	0	0
All	All	14715	0	14906	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 66 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:136:ASP:HA	1:E:146:ALA:HB2	1.81	0.61
4:A:247:THR:HG21	4:A:265:GLN:HE21	1.68	0.59
1:E:43:LEU:HD11	1:E:60:ILE:CD1	2.34	0.58
4:A:243:PRO:HD3	4:A:323:ILE:HD12	1.85	0.57
3:G:210:LEU:HD21	4:A:88:PRO:HD2	1.85	0.56

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	368/380 (97%)	353 (96%)	12 (3%)	3 (1%)	19	51
2	F	479/514 (93%)	447 (93%)	31 (6%)	1 (0%)	47	78
3	G	280/296 (95%)	260 (93%)	17 (6%)	3 (1%)	14	42
4	A	369/381 (97%)	348 (94%)	19 (5%)	2 (0%)	29	61
4	B	364/381 (96%)	345 (95%)	18 (5%)	1 (0%)	41	71
All	All	1860/1952 (95%)	1753 (94%)	97 (5%)	10 (0%)	29	61

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	173	ASN
3	G	137	LEU
3	G	7	LYS
4	A	21	ASP
4	A	322	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	297/305 (97%)	294 (99%)	3 (1%)	76	92
2	F	394/424 (93%)	389 (99%)	5 (1%)	69	90
3	G	228/237 (96%)	217 (95%)	11 (5%)	25	58
4	A	315/323 (98%)	306 (97%)	9 (3%)	42	76
4	B	312/323 (97%)	305 (98%)	7 (2%)	52	81
All	All	1546/1612 (96%)	1511 (98%)	35 (2%)	50	80

5 of 35 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	B	31	PHE
4	B	35	VAL
4	B	181	LYS
3	G	54	ILE
3	G	21	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
3	G	250	GLN
4	A	116	GLN
4	B	305	GLN
4	A	265	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	A	305	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

Of 7 monosaccharides modelled in this entry, 6 were used for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	GLC	C	1	5	12,12,12	0.52	0	17,17,17	0.50	0
5	GLC	C	2	5	11,11,12	0.65	0	15,15,17	0.88	0
5	GLC	C	3	5	11,11,12	0.72	0	15,15,17	0.90	1 (6%)
5	GLC	C	4	5	11,11,12	0.52	0	15,15,17	1.24	1 (6%)
6	GLC	D	2	6	11,11,12	0.54	0	15,15,17	1.20	1 (6%)
6	GLC	D	3	6	11,11,12	0.62	0	15,15,17	0.98	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GLC	C	1	5	-	0/2/22/22	0/1/1/1
5	GLC	C	2	5	-	0/2/19/22	0/1/1/1
5	GLC	C	3	5	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GLC	C	4	5	-	2/2/19/22	0/1/1/1
6	GLC	D	2	6	-	2/2/19/22	0/1/1/1
6	GLC	D	3	6	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	C	4	GLC	C1-O5-C5	3.92	117.50	112.19
6	D	2	GLC	C1-O5-C5	3.38	116.77	112.19
5	C	3	GLC	O5-C1-C2	-2.46	106.97	110.77
6	D	3	GLC	C1-O5-C5	2.23	115.22	112.19

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	4	GLC	O5-C5-C6-O6
6	D	2	GLC	O5-C5-C6-O6
5	C	4	GLC	C4-C5-C6-O6
6	D	2	GLC	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	D	3	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

Oligosaccharide Chain C



Oligosaccharide Chain D



5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
7	PGV	F	604	-	50,50,50	1.08	3 (6%)	53,56,56	0.97	2 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	PGV	F	604	-	-	23/55/55/55	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	F	604	PGV	O01-C1	4.24	1.46	1.34
7	F	604	PGV	O03-C19	4.22	1.45	1.33
7	F	604	PGV	C12-C11	3.69	1.53	1.31

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
7	F	604	PGV	O01-C1-C2	3.97	120.06	111.50
7	F	604	PGV	O03-C19-C20	2.63	120.17	111.91

There are no chirality outliers.

5 of 23 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	F	604	PGV	C03-O11-P-O13
7	F	604	PGV	C04-O12-P-O13
7	F	604	PGV	C2-C1-O01-C02

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
7	F	604	PGV	O02-C1-O01-C02
7	F	604	PGV	C20-C19-O03-C01

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand PGV F 604



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	E	370/380 (97%)	-0.16	8 (2%)	62	59	57, 82, 117, 133	0
2	F	483/514 (93%)	0.22	29 (6%)	21	18	63, 101, 154, 189	0
3	G	282/296 (95%)	-0.03	11 (3%)	39	35	58, 86, 145, 170	0
4	A	370/381 (97%)	-0.01	14 (3%)	40	36	69, 91, 123, 176	0
4	B	368/381 (96%)	0.24	27 (7%)	15	11	66, 100, 170, 218	0
All	All	1873/1952 (95%)	0.07	89 (4%)	30	27	57, 94, 150, 218	0

The worst 5 of 89 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	B	280	ASN	6.2
4	A	15	GLU	5.9
4	B	340	GLY	5.9
2	F	40	TYR	5.7
4	B	105	ALA	5.4

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	GLC	C	4	11/12	0.93	0.27	78,81,84,84	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
6	GLC	D	2	11/12	0.95	0.10	62,62,64,64	0
5	GLC	C	3	11/12	0.96	0.11	73,74,77,77	0
5	GLC	C	2	11/12	0.97	0.12	67,70,72,72	0
6	GLC	D	3	11/12	0.97	0.18	61,62,63,63	0
5	GLC	C	1	12/12	0.98	0.13	59,62,63,65	0
6	GLC	D	1	1/12	0.98	0.08	66,66,66,66	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain D:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
7	PGV	F	604	51/51	0.79	0.29	100,108,121,122	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around PGV F 604:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.