



# wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 29, 2024 – 03:04 pm BST

PDB ID : 5LAY  
Title : Discovery of New Natural-product-inspired Spiro-oxindole Compounds as Orally Active Inhibitors of the MDM2-p53 Interaction: HDM2 (MDM2) IN COMPLEX WITH COMPOUND 6g  
Authors : Kessler, D.; Gollner, A.  
Deposited on : 2016-06-15  
Resolution : 2.71 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36.2
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3359 (2.74-2.70)
Clashscore	141614	3686 (2.74-2.70)
Ramachandran outliers	138981	3622 (2.74-2.70)
Sidechain outliers	138945	3623 (2.74-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	95	94% 5% .
1	B	95	95% . .
1	C	95	94% 5% .
1	D	95	86% 13% .
1	E	95	93% 6% .
1	F	95	79% 6% 15%

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 4950 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E3 ubiquitin-protein ligase Mdm2.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	95	Total	C	N	O	S	11	1	0
			801	518	137	142	4			
1	B	94	Total	C	N	O	S	12	3	0
			796	517	132	143	4			
1	C	94	Total	C	N	O	S	31	0	0
			773	503	128	138	4			
1	D	94	Total	C	N	O	S	11	0	0
			771	502	127	138	4			
1	E	94	Total	C	N	O	S	46	0	0
			774	503	128	139	4			
1	F	81	Total	C	N	O	S	108	0	0
			677	441	111	121	4			

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0
2	E	1	Total C O 6 3 3	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0
3	E	1	Total O S 5 4 1	0	0

- Molecule 4 is (3 {S},3' {S},4' {S},5' {S})-4'-azanyl-6-chloranyl-3'-(3-chloranyl-2-fluoranyl-phenyl)-1'-[(3-ethoxyphenyl)methyl]-5'-methyl-spiro[1 {H}-indole-3,2'-pyrrolidine]-2-one (three-letter code: 6SS) (formula: C<sub>27</sub>H<sub>26</sub>Cl<sub>2</sub>FN<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total 35	C 27	Cl 2	F 1	N 3	O 2	0	0
4	B	1	Total 35	C 27	Cl 2	F 1	N 3	O 2	0	0
4	C	1	Total 35	C 27	Cl 2	F 1	N 3	O 2	0	0
4	D	1	Total 35	C 27	Cl 2	F 1	N 3	O 2	0	0
4	E	1	Total 35	C 27	Cl 2	F 1	N 3	O 2	0	0
4	F	1	Total 35	C 27	Cl 2	F 1	N 3	O 2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	26	Total	O	0	0
			26	26		
5	B	23	Total	O	0	0
			23	23		
5	C	14	Total	O	0	0
			14	14		
5	D	26	Total	O	0	0
			26	26		
5	E	6	Total	O	0	0
			6	6		
5	F	3	Total	O	0	0
			3	3		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: E3 ubiquitin-protein ligase Mdm2

Chain A:  94% 5%



- Molecule 1: E3 ubiquitin-protein ligase Mdm2

Chain B:  95%




- Molecule 1: E3 ubiquitin-protein ligase Mdm2

Chain C:  94% 5%




- Molecule 1: E3 ubiquitin-protein ligase Mdm2

Chain D:  86% 13%




- Molecule 1: E3 ubiquitin-protein ligase Mdm2

Chain E:  93% 6%



- Molecule 1: E3 ubiquitin-protein ligase Mdm2

Chain F:  79% 6% 15%

SER	GLN	ILE	PRO	ALA	SER	E23	Q24	K39	A43	Y48	Y56	S92	VAL	LYS	E95	Y104	R105	ASN	LEU	VAL	VAL	VAL	ASN

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.45Å 71.45Å 520.98Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	87.45 – 2.71 44.86 – 2.71	Depositor EDS
% Data completeness (in resolution range)	99.0 (87.45-2.71) 99.0 (44.86-2.71)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.16	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.94 (at 2.73Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.251 , 0.292 0.319 , 0.339	Depositor DCC
$R_{free}$ test set	806 reflections (3.55%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.4	Xtriage
Anisotropy	0.435	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 32.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.79	EDS
Total number of atoms	4950	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	8.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.45% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, 6SS, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.42	0/816	0.54	0/1101
1	B	0.42	0/811	0.58	0/1095
1	C	0.48	0/788	0.56	0/1063
1	D	0.41	0/786	0.56	0/1060
1	E	0.43	0/789	0.54	0/1063
1	F	0.65	2/690 (0.3%)	0.80	3/926 (0.3%)
All	All	0.47	2/4680 (0.0%)	0.60	3/6308 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	104	TYR	CG-CD1	8.86	1.50	1.39
1	F	104	TYR	CG-CD2	-7.22	1.29	1.39

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	104	TYR	CB-CG-CD1	-11.13	114.32	121.00
1	F	104	TYR	CB-CG-CD2	11.09	127.65	121.00
1	F	56	TYR	CB-CG-CD1	5.86	124.52	121.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	801	0	836	5	0
1	B	796	0	826	4	0
1	C	773	0	805	5	0
1	D	771	0	804	8	0
1	E	774	0	805	3	0
1	F	677	0	696	2	0
2	A	18	0	24	0	0
2	C	6	0	8	1	0
2	E	6	0	8	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
3	D	5	0	0	0	0
3	E	5	0	0	0	0
4	A	35	0	0	0	0
4	B	35	0	0	0	0
4	C	35	0	0	0	0
4	D	35	0	0	0	0
4	E	35	0	0	0	0
4	F	35	0	0	0	0
5	A	26	0	0	0	0
5	B	23	0	0	0	0
5	C	14	0	0	0	0
5	D	26	0	0	0	0
5	E	6	0	0	0	0
5	F	3	0	0	0	0
All	All	4950	0	4812	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 23 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:18:GLN:N	2:C:201:GOL:HO2	1.95	0.64
1:C:37:LEU:HD13	1:C:85:LEU:HD21	1.88	0.56
1:D:41:VAL:HG11	1:D:60:TYR:HA	1.89	0.54
1:F:39:LYS:HA	1:F:43:ALA:HB3	1.90	0.53
1:D:38:LEU:HD22	1:D:56:TYR:HB2	1.92	0.52

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	95/95 (100%)	92 (97%)	3 (3%)	0	100	100
1	B	95/95 (100%)	93 (98%)	2 (2%)	0	100	100
1	C	92/95 (97%)	89 (97%)	3 (3%)	0	100	100
1	D	92/95 (97%)	90 (98%)	2 (2%)	0	100	100
1	E	92/95 (97%)	88 (96%)	4 (4%)	0	100	100
1	F	77/95 (81%)	72 (94%)	5 (6%)	0	100	100
All	All	543/570 (95%)	524 (96%)	19 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	91/89 (102%)	90 (99%)	1 (1%)	73	89
1	B	91/89 (102%)	91 (100%)	0	100	100
1	C	88/89 (99%)	88 (100%)	0	100	100
1	D	88/89 (99%)	88 (100%)	0	100	100
1	E	88/89 (99%)	88 (100%)	0	100	100
1	F	76/89 (85%)	75 (99%)	1 (1%)	69	86
All	All	522/534 (98%)	520 (100%)	2 (0%)	91	96

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	78	SER
1	F	24	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	18	GLN
1	F	79	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	D	201	-	4,4,4	0.34	0	6,6,6	0.09	0
2	GOL	E	201	-	5,5,5	0.31	0	5,5,5	0.35	0
2	GOL	A	203	-	5,5,5	0.34	0	5,5,5	0.23	0
2	GOL	A	202	-	5,5,5	0.28	0	5,5,5	0.18	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	6SS	C	202	-	39,39,39	1.07	2 (5%)	48,59,59	1.66	11 (22%)
4	6SS	D	202	-	39,39,39	1.17	2 (5%)	48,59,59	1.75	9 (18%)
3	SO4	A	204	-	4,4,4	0.33	0	6,6,6	0.07	0
4	6SS	A	205	-	39,39,39	1.24	2 (5%)	48,59,59	1.88	9 (18%)
4	6SS	F	201	-	39,39,39	1.17	3 (7%)	48,59,59	1.60	7 (14%)
2	GOL	C	201	-	5,5,5	0.30	0	5,5,5	0.29	0
2	GOL	A	201	-	5,5,5	0.43	0	5,5,5	0.43	0
3	SO4	B	201	-	4,4,4	0.34	0	6,6,6	0.08	0
4	6SS	B	202	-	39,39,39	1.21	2 (5%)	48,59,59	1.95	11 (22%)
3	SO4	E	202	-	4,4,4	0.36	0	6,6,6	0.18	0
4	6SS	E	203	-	39,39,39	1.09	2 (5%)	48,59,59	1.69	9 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	E	201	-	-	0/4/4/4	-
2	GOL	A	203	-	-	2/4/4/4	-
2	GOL	A	202	-	-	2/4/4/4	-
4	6SS	C	202	-	-	2/11/51/51	0/5/5/5
4	6SS	D	202	-	-	2/11/51/51	0/5/5/5
4	6SS	A	205	-	-	2/11/51/51	0/5/5/5
4	6SS	F	201	-	-	2/11/51/51	0/5/5/5
2	GOL	C	201	-	-	2/4/4/4	-
2	GOL	A	201	-	-	2/4/4/4	-
4	6SS	E	203	-	-	3/11/51/51	0/5/5/5
4	6SS	B	202	-	-	2/11/51/51	0/5/5/5

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	205	6SS	C26-C35	-5.87	1.51	1.55
4	B	202	6SS	C26-C35	-5.31	1.51	1.55
4	E	203	6SS	C26-C35	-5.14	1.51	1.55
4	D	202	6SS	C26-C35	-5.01	1.51	1.55
4	C	202	6SS	C26-C35	-4.54	1.52	1.55

The worst 5 of 56 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	205	6SS	C27-C26-C35	7.71	104.92	101.59
4	B	202	6SS	C27-C26-C35	7.40	104.79	101.59
4	E	203	6SS	C27-C26-C35	6.87	104.56	101.59
4	D	202	6SS	C27-C26-C35	6.56	104.43	101.59
4	C	202	6SS	C27-C26-C35	5.56	103.99	101.59

There are no chirality outliers.

5 of 21 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	201	GOL	O1-C1-C2-C3
2	A	202	GOL	O1-C1-C2-C3
2	C	201	GOL	C1-C2-C3-O3
4	E	203	6SS	C25-C21-O22-C23
2	C	201	GOL	O2-C2-C3-O3


There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	201	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


## Ligand 6SS C 202




## Ligand 6SS D 202




## Ligand 6SS A 205




## Ligand 6SS F 201



## Ligand 6SS B 202



## Ligand 6SS E 203



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.




### 6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around 6SS A 205:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around 6SS B 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around 6SS C 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around 6SS D 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around 6SS E 203:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around 6SS F 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.