



Full wwPDB X-ray Structure Validation Report i

May 26, 2020 – 01:27 am BST

PDB ID : 6LI2
Title : Crystal structure of GPR52 ligand free form with rubredoxin fusion
Authors : Luo, Z.P.; Lin, X.; Xu, F.; Han, G.W.
Deposited on : 2019-12-10
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	372	 4% (Red) 80% (Green) 12% (Yellow) • 7% (Grey)

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 2956 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chimera of G-protein coupled receptor 52 and Rubredoxin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	347	Total	C 2761	N 1825	O 425	S 483	28	0	0

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	16	GLY	-	expression tag	UNP Q9Y2T5
A	130	TRP	ALA	conflict	UNP Q9Y2T5
A	1001	MET	-	linker	UNP Q9Y2T5
A	263	TYR	-	linker	UNP P00268
A	264	LEU	-	linker	UNP P00268
A	278	GLN	TRP	engineered mutation	UNP Q9Y2T5
A	314	PRO	CYS	engineered mutation	UNP Q9Y2T5
A	318	ALA	SER	engineered mutation	UNP Q9Y2T5
A	321	ASP	ASN	engineered mutation	UNP Q9Y2T5
A	323	THR	VAL	engineered mutation	UNP Q9Y2T5
A	601	GLU	-	expression tag	UNP Q9Y2T5
A	602	PHE	-	expression tag	UNP Q9Y2T5
A	603	LEU	-	expression tag	UNP Q9Y2T5
A	604	GLU	-	expression tag	UNP Q9Y2T5
A	605	VAL	-	expression tag	UNP Q9Y2T5
A	606	LEU	-	expression tag	UNP Q9Y2T5
A	607	PHE	-	expression tag	UNP Q9Y2T5
A	608	GLN	-	expression tag	UNP Q9Y2T5
A	609	GLY	-	expression tag	UNP Q9Y2T5
A	610	PRO	-	expression tag	UNP Q9Y2T5
A	611	HIS	-	expression tag	UNP Q9Y2T5
A	612	HIS	-	expression tag	UNP Q9Y2T5
A	613	HIS	-	expression tag	UNP Q9Y2T5
A	614	HIS	-	expression tag	UNP Q9Y2T5
A	615	HIS	-	expression tag	UNP Q9Y2T5
A	616	HIS	-	expression tag	UNP Q9Y2T5
A	617	HIS	-	expression tag	UNP Q9Y2T5

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	618	HIS	-	expression tag	UNP Q9Y2T5
A	619	HIS	-	expression tag	UNP Q9Y2T5
A	620	HIS	-	expression tag	UNP Q9Y2T5

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0

- Molecule 3 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLC) (formula: C₂₁H₄₀O₄).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 14 10 4	0	0
3	A	1	Total C O 15 11 4	0	0
3	A	1	Total C O 15 11 4	0	0
3	A	1	Total C O 21 17 4	0	0
3	A	1	Total C O 12 8 4	0	0
3	A	1	Total C O 15 11 4	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 12 8 4	0	0
3	A	1	Total C O 15 11 4	0	0
3	A	1	Total C O 13 9 4	0	0
3	A	1	Total C O 12 8 4	0	0
3	A	1	Total C O 25 21 4	0	0

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 7 4 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	18	Total O 18 18	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Chimera of G-protein coupled receptor 52 and Rubredoxin



4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	77.22Å 113.23Å 138.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.25 – 2.80 29.23 – 2.80	Depositor EDS
% Data completeness (in resolution range)	81.1 (29.25-2.80) 81.2 (29.23-2.80)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	3.50 (at 2.80Å)	Xtriage
Refinement program	REFMAC 5.8.0253	Depositor
R , R_{free}	0.241 , 0.263 0.245 , 0.263	Depositor DCC
R_{free} test set	612 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	49.8	Xtriage
Anisotropy	0.148	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 50.2	EDS
L-test for twinning ²	$< L > = 0.45$, $< L^2 > = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	2956	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OLC, ZN, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.69	0/2845	0.77	0/3887

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbit. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2761	0	2691	36	0
2	A	1	0	0	0	0
3	A	169	0	216	6	0
4	A	7	0	10	0	0
5	A	18	0	0	0	0
All	All	2956	0	2917	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:182:LYS:HE2	1:A:199:THR:HG23	1.69	0.73
1:A:149:TYR:O	1:A:153:VAL:HG22	1.93	0.68
1:A:98:THR:O	1:A:102:LEU:HG	1.95	0.67
1:A:208:VAL:HG21	1:A:286:LEU:HD21	1.81	0.61
1:A:53:LEU:HB3	1:A:311:PHE:CD1	2.38	0.59
1:A:47:ILE:HD12	1:A:101:LEU:HD11	1.86	0.56
1:A:59:LEU:HD23	1:A:62:ILE:HD11	1.88	0.56
1:A:182:LYS:HB3	1:A:199:THR:HG22	1.88	0.55
1:A:1044:VAL:HB	1:A:1048:GLN:HG3	1.89	0.54
1:A:1011:TYR:HB2	1:A:1041:LEU:HD12	1.92	0.51
1:A:168:TYR:O	1:A:172:ILE:HG12	2.11	0.51
1:A:339:CYS:O	1:A:603:LEU:HB2	2.11	0.51
1:A:79:PHE:CZ	1:A:134:CYS:HB3	2.46	0.51
1:A:47:ILE:HG22	1:A:98:THR:CG2	2.41	0.50
1:A:174:LEU:N	1:A:175:PRO:CD	2.75	0.50
1:A:33:HIS:HB3	3:A:1211:OLC:H22	1.92	0.50
1:A:53:LEU:HB3	1:A:311:PHE:CE1	2.46	0.50
1:A:266:VAL:O	1:A:270:ILE:HG13	2.12	0.49
1:A:103:HIS:NE2	3:A:1204:OLC:H24A	2.29	0.48
1:A:324:PHE:CE1	3:A:1206:OLC:H2	2.50	0.47
1:A:64:VAL:HG13	3:A:1206:OLC:H24A	1.98	0.46
1:A:272:SER:HB2	3:A:1202:OLC:H21	1.98	0.45
1:A:315:VAL:O	1:A:319:LEU:HB2	2.16	0.45
1:A:47:ILE:HG22	1:A:98:THR:HG22	1.98	0.45
1:A:1008:VAL:HG21	1:A:1044:VAL:HG11	1.99	0.44
1:A:31:PHE:CD1	1:A:31:PHE:N	2.85	0.44
1:A:292:VAL:HG12	1:A:292:VAL:O	2.18	0.44
1:A:75:THR:OG1	1:A:138:ASP:OD2	2.36	0.44
1:A:76:THR:HG23	1:A:135:ILE:HG12	2.00	0.42
1:A:47:ILE:HD12	1:A:101:LEU:CD1	2.49	0.42
1:A:185:TYR:OH	1:A:288:GLU:HB3	2.20	0.42
1:A:182:LYS:HB3	1:A:199:THR:CG2	2.50	0.41
1:A:72:HIS:HB3	1:A:73:HIS:H	1.74	0.41
1:A:603:LEU:HA	1:A:603:LEU:HD12	1.86	0.41
1:A:38:ASP:OD2	3:A:1211:OLC:O25	2.25	0.41
1:A:1042:CYS:SG	1:A:1044:VAL:HG22	2.62	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	343/372 (92%)	335 (98%)	8 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	309/338 (91%)	307 (99%)	2 (1%)	86 96

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	PHE
1	A	72	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 13 ligands modelled in this entry, 1 is monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	OLC	A	1212	-	24,24,24	0.90	1 (4%)	25,25,25	0.96	2 (8%)
3	OLC	A	1209	-	14,14,24	1.23	1 (7%)	15,15,25	1.19	2 (13%)
3	OLC	A	1204	-	14,14,24	1.20	1 (7%)	15,15,25	1.12	1 (6%)
3	OLC	A	1207	-	14,14,24	1.07	1 (7%)	15,15,25	1.07	2 (13%)
3	OLC	A	1203	-	14,14,24	1.34	1 (7%)	15,15,25	0.92	1 (6%)
3	OLC	A	1206	-	11,11,24	1.28	1 (9%)	12,12,25	1.02	1 (8%)
3	OLC	A	1210	-	12,12,24	1.19	1 (8%)	13,13,25	1.14	2 (15%)
3	OLC	A	1208	-	11,11,24	1.38	1 (9%)	12,12,25	1.24	2 (16%)
3	OLC	A	1202	-	13,13,24	1.21	1 (7%)	14,14,25	1.44	2 (14%)
4	PEG	A	1213	-	6,6,6	0.17	0	5,5,5	0.21	0
3	OLC	A	1205	-	20,20,24	1.10	1 (5%)	21,21,25	0.96	1 (4%)
3	OLC	A	1211	-	11,11,24	1.48	1 (9%)	12,12,25	1.07	1 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OLC	A	1212	-	-	9/24/24/24	-
3	OLC	A	1209	-	-	4/14/14/24	-
3	OLC	A	1204	-	-	4/14/14/24	-
3	OLC	A	1207	-	-	6/14/14/24	-
3	OLC	A	1203	-	-	7/14/14/24	-
3	OLC	A	1206	-	-	4/11/11/24	-
3	OLC	A	1210	-	-	3/12/12/24	-
3	OLC	A	1208	-	-	3/11/11/24	-
3	OLC	A	1202	-	-	3/13/13/24	-
4	PEG	A	1213	-	-	1/4/4/4	-
3	OLC	A	1205	-	-	10/20/20/24	-
3	OLC	A	1211	-	-	6/11/11/24	-

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1211	OLC	O20-C1	4.73	1.47	1.33
3	A	1205	OLC	O20-C1	4.66	1.47	1.33
3	A	1203	OLC	O20-C1	4.61	1.46	1.33
3	A	1208	OLC	O20-C1	4.41	1.46	1.33
3	A	1209	OLC	O20-C1	4.41	1.46	1.33
3	A	1204	OLC	O20-C1	4.28	1.45	1.33
3	A	1202	OLC	O20-C1	4.20	1.45	1.33
3	A	1212	OLC	O20-C1	4.17	1.45	1.33
3	A	1210	OLC	O20-C1	3.84	1.44	1.33
3	A	1206	OLC	O20-C1	3.83	1.44	1.33
3	A	1207	OLC	O20-C1	3.74	1.44	1.33

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1202	OLC	O20-C1-C2	3.72	123.58	111.91
3	A	1204	OLC	O20-C1-C2	3.22	122.02	111.91
3	A	1205	OLC	O20-C1-C2	3.11	121.65	111.91
3	A	1209	OLC	O20-C1-C2	3.10	121.64	111.91
3	A	1208	OLC	O20-C1-C2	3.01	121.35	111.91
3	A	1207	OLC	O20-C1-C2	2.80	120.68	111.91
3	A	1202	OLC	O20-C1-O19	-2.78	116.57	123.59
3	A	1210	OLC	O20-C1-C2	2.71	120.43	111.91
3	A	1212	OLC	O20-C1-C2	2.71	120.40	111.91

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	A	1211	OLC	O20-C1-C2	2.56	119.95	111.91
3	A	1206	OLC	O20-C1-C2	2.44	119.57	111.91
3	A	1203	OLC	O20-C1-O19	-2.39	117.57	123.59
3	A	1207	OLC	O20-C1-O19	-2.33	117.72	123.59
3	A	1210	OLC	O20-C1-O19	-2.31	117.76	123.59
3	A	1208	OLC	O20-C1-O19	-2.28	117.83	123.59
3	A	1209	OLC	O20-C1-O19	-2.13	118.23	123.59
3	A	1212	OLC	O20-C1-O19	-2.00	118.54	123.59

There are no chirality outliers.

All (60) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1209	OLC	O20-C21-C22-O23
3	A	1207	OLC	C21-C22-C24-O25
3	A	1205	OLC	O20-C21-C22-C24
3	A	1211	OLC	O20-C21-C22-C24
3	A	1211	OLC	O20-C21-C22-O23
3	A	1209	OLC	C2-C1-O20-C21
3	A	1211	OLC	C2-C1-O20-C21
3	A	1211	OLC	O19-C1-O20-C21
3	A	1209	OLC	O19-C1-O20-C21
3	A	1212	OLC	O20-C21-C22-O23
3	A	1205	OLC	O20-C21-C22-O23
3	A	1203	OLC	C2-C1-O20-C21
3	A	1203	OLC	O19-C1-O20-C21
3	A	1212	OLC	C2-C1-O20-C21
3	A	1212	OLC	O20-C21-C22-C24
3	A	1210	OLC	C2-C3-C4-C5
3	A	1202	OLC	C1-C2-C3-C4
3	A	1205	OLC	C1-C2-C3-C4
3	A	1212	OLC	C1-C2-C3-C4
3	A	1207	OLC	C1-C2-C3-C4
3	A	1203	OLC	C1-C2-C3-C4
3	A	1211	OLC	C1-C2-C3-C4
3	A	1212	OLC	O19-C1-O20-C21
3	A	1210	OLC	C2-C1-O20-C21
3	A	1205	OLC	C2-C1-O20-C21
3	A	1203	OLC	C3-C4-C5-C6
3	A	1205	OLC	C2-C3-C4-C5
3	A	1206	OLC	C1-C2-C3-C4
3	A	1207	OLC	O23-C22-C24-O25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	1210	OLC	O19-C1-O20-C21
3	A	1205	OLC	O19-C1-O20-C21
3	A	1209	OLC	O20-C21-C22-C24
3	A	1212	OLC	C15-C16-C17-C18
3	A	1204	OLC	C4-C5-C6-C7
3	A	1204	OLC	C2-C3-C4-C5
3	A	1208	OLC	C2-C1-O20-C21
3	A	1202	OLC	C4-C5-C6-C7
3	A	1212	OLC	C5-C6-C7-C8
3	A	1208	OLC	O20-C21-C22-C24
3	A	1208	OLC	O19-C1-O20-C21
3	A	1206	OLC	C2-C3-C4-C5
3	A	1205	OLC	C10-C11-C12-C13
3	A	1211	OLC	C2-C3-C4-C5
3	A	1212	OLC	C10-C11-C12-C13
4	A	1213	PEG	O2-C3-C4-O4
3	A	1206	OLC	O20-C21-C22-C24
3	A	1207	OLC	C2-C3-C4-C5
3	A	1207	OLC	C4-C5-C6-C7
3	A	1205	OLC	C5-C6-C7-C8
3	A	1207	OLC	O20-C21-C22-C24
3	A	1203	OLC	C5-C6-C7-C8
3	A	1202	OLC	C2-C3-C4-C5
3	A	1203	OLC	O23-C22-C24-O25
3	A	1206	OLC	O20-C21-C22-O23
3	A	1212	OLC	C7-C8-C9-C10
3	A	1205	OLC	C9-C10-C11-C12
3	A	1203	OLC	C21-C22-C24-O25
3	A	1204	OLC	O20-C1-C2-C3
3	A	1204	OLC	C5-C6-C7-C8
3	A	1205	OLC	O20-C1-C2-C3


There are no ring outliers.


4 monomers are involved in 6 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1204	OLC	1	0
3	A	1206	OLC	2	0
3	A	1202	OLC	1	0
3	A	1211	OLC	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,


bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	347/372 (93%)	0.02	15 (4%) 35 25	43, 62, 106, 134	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	107	GLY	5.1
1	A	109	HIS	3.4
1	A	337	THR	3.4
1	A	604	GLU	3.4
1	A	151	GLN	2.9
1	A	339	CYS	2.9
1	A	340	MET	2.9
1	A	234	THR	2.8
1	A	104	TYR	2.5
1	A	155	PRO	2.3
1	A	603	LEU	2.3
1	A	156	CYS	2.1
1	A	233	HIS	2.1
1	A	108	VAL	2.0
1	A	601	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates i


There are no carbohydrates in this entry.


6.4 Ligands [\(i\)](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	OLC	A	1204	15/25	0.75	0.31	61,100,111,125	0
3	OLC	A	1208	12/25	0.79	0.40	69,118,124,124	0
3	OLC	A	1207	15/25	0.81	0.26	65,83,99,103	0
3	OLC	A	1203	15/25	0.82	0.23	58,69,101,101	0
3	OLC	A	1210	13/25	0.85	0.45	66,89,111,132	0
3	OLC	A	1202	14/25	0.85	0.30	58,81,105,106	0
3	OLC	A	1205	21/25	0.86	0.19	52,75,88,98	0
3	OLC	A	1211	12/25	0.86	0.24	60,76,109,115	0
4	PEG	A	1213	7/7	0.88	0.15	36,53,60,71	0
3	OLC	A	1209	15/25	0.89	0.17	69,84,106,112	0
3	OLC	A	1212	25/25	0.89	0.36	53,68,82,92	0
3	OLC	A	1206	12/25	0.93	0.18	53,74,89,94	0
2	ZN	A	1201	1/1	0.99	0.05	56,56,56,56	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.














6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.