



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 2, 2023 – 12:57 AM EDT

PDB ID : 6MJI
Title : Crystal structure of the mCD1d/xxs (JJ304) /iNKTCR ternary complex
Authors : Zajonc, D.M.; Bitra, A.; Janssens, J.
Deposited on : 2018-09-20
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : **FAILED**
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : **FAILED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 7014 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called T cell receptor alpha variable 11,T cell receptor alpha joining 18,Human nkt tcr alpha chain, CHIMERIC PROTEIN.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| | | | Total | C | N | O | S | | | |
| 1 | C | 203 | 1570 | 972 | 270 | 320 | 8 | 0 | 2 | 0 |

There are 2 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|-----------------------|----------------|
| C | 0 | MET | - | initiating methionine | UNP A0A0B4J1J9 |
| C | 113 | ILE | - | linker | UNP A0A0B4J1J9 |

- Molecule 2 is a protein called Beta-chain,Tcell receptor chain,T cell receptor beta constant 2.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|---------|-------|
| | | | Total | C | N | O | S | | | |
| 2 | D | 239 | 1884 | 1182 | 336 | 360 | 6 | 0 | 1 | 0 |

There are 7 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|---------|------------|
| D | 95 | ASP | - | linker | UNP A2NTY6 |
| D | 96 | GLU | - | linker | UNP A2NTY6 |
| D | 97 | GLY | - | linker | UNP A2NTY6 |
| D | 98 | TYR | - | linker | UNP A2NTY6 |
| D | 130 | ALA | ALA | linker | UNP A0N8J3 |
| D | 168 | CYS | SER | variant | UNP A0A5B9 |
| D | 186 | SER | CYS | variant | UNP A0A5B9 |

- Molecule 3 is a protein called Antigen-presenting glycoprotein CD1d1.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|---------|-------|
| | | | Total | C | N | O | S | | | |
| 3 | A | 270 | 2167 | 1384 | 373 | 397 | 13 | 0 | 0 | 0 |

There are 6 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|----------------|----------------|
| A | 280 | HIS | - | expression tag | UNP A0A0R4J090 |
| A | 281 | HIS | - | expression tag | UNP A0A0R4J090 |
| A | 282 | HIS | - | expression tag | UNP A0A0R4J090 |
| A | 283 | HIS | - | expression tag | UNP A0A0R4J090 |
| A | 284 | HIS | - | expression tag | UNP A0A0R4J090 |
| A | 285 | HIS | - | expression tag | UNP A0A0R4J090 |

- Molecule 4 is a protein called Beta-2-microglobulin.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|---------|-------|
| | | | Total | C | N | O | S | | | |
| 4 | B | 97 | 793 | 506 | 134 | 147 | 6 | 0 | 0 | 0 |

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|----|---|----|---------|---------|-------|
| | | | Total | C | N | O | | | |
| 5 | E | 2 | 28 | 16 | 2 | 10 | 0 | 0 | 0 |


- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.

| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|----|---|----|---------|---------|-------|
| | | | Total | C | N | O | | | |
| 6 | F | 5 | 60 | 34 | 2 | 24 | 0 | 0 | 0 |

- Molecule 7 is SODIUM ION (three-letter code: NA) (formula: Na).

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---------|---------|
| | | | Total | Na | | |
| 7 | C | 1 | 1 | 1 | 0 | 0 |
| 7 | D | 1 | 1 | 1 | 0 | 0 |

- Molecule 8 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




| Mol | Chain | Residues | Atoms | | | ZeroOcc | AltConf |
|-----|-------|----------|-------|---|---|---------|---------|
| 8 | D | 1 | Total | C | O | 0 | 0 |
| | | | 6 | 3 | 3 | | |
| 8 | D | 1 | Total | C | O | 0 | 0 |
| | | | 6 | 3 | 3 | | |
| 8 | D | 1 | Total | C | O | 0 | 0 |
| | | | 6 | 3 | 3 | | |
| 8 | D | 1 | Total | C | O | 0 | 0 |
| | | | 6 | 3 | 3 | | |
| 8 | A | 1 | Total | C | O | 0 | 0 |
| | | | 6 | 3 | 3 | | |
| 8 | B | 1 | Total | C | O | 0 | 0 |
| | | | 6 | 3 | 3 | | |

- Molecule 9 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf |
|-----|-------|----------|-------|---|---|---|---------|---------|
| | | | Total | C | N | O | | |
| 9 | A | 1 | 14 | 8 | 1 | 5 | 0 | 0 |

- Molecule 10 is N-[(2S,3S,4R)-1-{[4-O-(cyclopropylmethyl)-alpha-D-galactopyranosyl]oxy}-3,4-dihydroxyoctadecan-2-yl]hexacosanamide (three-letter code: JTD) (formula: $C_{54}H_{105}NO_9$).



| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---|---|---------|---------|
| | | | Total | C | N | O | | |
| 10 | A | 1 | 64 | 54 | 1 | 9 | 0 | 0 |

- Molecule 11 is water.

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|--------------|----------|---------|---------|
| 11 | C | 86 | Total 86 | O 86 | 0 | 0 |
| 11 | D | 151 | Total 151 | O 151 | 0 | 0 |
| 11 | A | 115 | Total 115 | O 115 | 0 | 0 |
| 11 | B | 44 | Total 44 | O 44 | 0 | 0 |

MolProbity and EDS failed to run properly - this section is therefore empty.

3 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

| Property | Value | Source |
|--|---|-----------|
| Space group | C 2 2 21 | Depositor |
| Cell constants a, b, c, α , β , γ | 79.22Å 191.16Å 151.09Å 90.00° 90.00° 90.00° | Depositor |
| Resolution (Å) | 45.61 – 2.30 | Depositor |
| % Data completeness (in resolution range) | 99.5 (45.61-2.30) | Depositor |
| R_{merge} | 0.11 | Depositor |
| R_{sym} | (Not available) | Depositor |
| $\langle I/\sigma(I) \rangle$ ¹ | 1.96 (at 2.29Å) | Xtrriage |
| Refinement program | REFMAC 5.8.0222 | Depositor |
| R, R_{free} | 0.189 , 0.227 | Depositor |
| Wilson B-factor (Å ²) | 41.7 | Xtrriage |
| Anisotropy | 0.052 | Xtrriage |
| L-test for twinning ² | $\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$ | Xtrriage |
| Estimated twinning fraction | No twinning to report. | Xtrriage |
| Total number of atoms | 7014 | wwPDB-VP |
| Average B, all atoms (Å ²) | 46.0 | wwPDB-VP |

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

7 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 5 | NAG | E | 1 | 3,5 | 14,14,15 | 0.39 | 0 | 17,19,21 | 0.77 | 0 |
| 5 | NAG | E | 2 | 5 | 14,14,15 | 0.37 | 0 | 17,19,21 | 0.93 | 1 (5%) |
| 6 | NAG | F | 1 | 3,6 | 14,14,15 | 0.33 | 0 | 17,19,21 | 0.96 | 1 (5%) |
| 6 | NAG | F | 2 | 6 | 14,14,15 | 0.31 | 0 | 17,19,21 | 0.75 | 0 |
| 6 | BMA | F | 3 | 6 | 11,11,12 | 0.42 | 0 | 15,15,17 | 0.75 | 0 |
| 6 | MAN | F | 4 | 6 | 11,11,12 | 0.32 | 0 | 15,15,17 | 1.09 | 1 (6%) |
| 6 | FUC | F | 5 | 6 | 10,10,11 | 0.51 | 0 | 14,14,16 | 0.69 | 0 |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|-----------|---------|
| 5 | NAG | E | 1 | 3,5 | - | 0/6/23/26 | 0/1/1/1 |
| 5 | NAG | E | 2 | 5 | - | 2/6/23/26 | 0/1/1/1 |
| 6 | NAG | F | 1 | 3,6 | - | 0/6/23/26 | 0/1/1/1 |
| 6 | NAG | F | 2 | 6 | - | 2/6/23/26 | 0/1/1/1 |
| 6 | BMA | F | 3 | 6 | - | 2/2/19/22 | 0/1/1/1 |
| 6 | MAN | F | 4 | 6 | - | 2/2/19/22 | 0/1/1/1 |
| 6 | FUC | F | 5 | 6 | - | - | 0/1/1/1 |

There are no bond length outliers.

All (3) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|----------|------|-------------|----------|
| 5 | E | 2 | NAG | O5-C5-C6 | 2.98 | 111.87 | 107.20 |
| 6 | F | 4 | MAN | C3-C4-C5 | 2.68 | 115.01 | 110.24 |
| 6 | F | 1 | NAG | O5-C5-C6 | 2.62 | 111.31 | 107.20 |

There are no chirality outliers.

All (8) torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|-------------|
| 5 | E | 2 | NAG | O5-C5-C6-O6 |
| 6 | F | 3 | BMA | O5-C5-C6-O6 |
| 6 | F | 2 | NAG | O5-C5-C6-O6 |
| 6 | F | 2 | NAG | C4-C5-C6-O6 |
| 6 | F | 3 | BMA | C4-C5-C6-O6 |

Continued on next page...


Continued from previous page...


| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|-------------|
| 5 | E | 2 | NAG | C4-C5-C6-O6 |
| 6 | F | 4 | MAN | C4-C5-C6-O6 |
| 6 | F | 4 | MAN | O5-C5-C6-O6 |

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





4.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 8 | GOL | D | 602 | - | 5,5,5 | 0.19 | 0 | 5,5,5 | 0.36 | 0 |
| 8 | GOL | A | 301 | - | 5,5,5 | 0.30 | 0 | 5,5,5 | 0.38 | 0 |
| 10 | JTD | A | 310 | - | 65,65,65 | 0.50 | 0 | 71,76,76 | 0.81 | 1 (1%) |
| 8 | GOL | D | 603 | - | 5,5,5 | 0.27 | 0 | 5,5,5 | 0.40 | 0 |
| 8 | GOL | D | 604 | - | 5,5,5 | 0.35 | 0 | 5,5,5 | 0.21 | 0 |
| 8 | GOL | D | 601 | - | 5,5,5 | 0.36 | 0 | 5,5,5 | 0.18 | 0 |
| 9 | NAG | A | 302 | 3 | 14,14,15 | 0.44 | 0 | 17,19,21 | 0.90 | 1 (5%) |

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 8 | GOL | B | 101 | - | 5,5,5 | 0.43 | 0 | 5,5,5 | 0.41 | 0 |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|-------------|---------|
| 8 | GOL | D | 602 | - | - | 1/4/4/4 | - |
| 8 | GOL | A | 301 | - | - | 2/4/4/4 | - |
| 10 | JTD | A | 310 | - | - | 12/63/85/85 | 0/2/2/2 |
| 8 | GOL | D | 603 | - | - | 4/4/4/4 | - |
| 8 | GOL | D | 604 | - | - | 0/4/4/4 | - |
| 8 | GOL | D | 601 | - | - | 4/4/4/4 | - |
| 9 | NAG | A | 302 | 3 | - | 0/6/23/26 | 0/1/1/1 |
| 8 | GOL | B | 101 | - | - | 4/4/4/4 | - |

There are no bond length outliers.

All (2) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-------------|-------|-------------|----------|
| 9 | A | 302 | NAG | C1-O5-C5 | 2.49 | 115.56 | 112.19 |
| 10 | A | 310 | JTD | CAN-NAO-CBE | -2.19 | 119.79 | 123.48 |

There are no chirality outliers.

All (27) torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|-----------------|
| 8 | D | 601 | GOL | O1-C1-C2-O2 |
| 8 | D | 601 | GOL | O1-C1-C2-C3 |
| 8 | D | 603 | GOL | C1-C2-C3-O3 |
| 8 | A | 301 | GOL | C1-C2-C3-O3 |
| 8 | B | 101 | GOL | O1-C1-C2-C3 |
| 8 | B | 101 | GOL | C1-C2-C3-O3 |
| 10 | A | 310 | JTD | CCO-CCP-CCQ-CCR |
| 8 | D | 601 | GOL | C1-C2-C3-O3 |
| 8 | D | 603 | GOL | O1-C1-C2-C3 |
| 10 | A | 310 | JTD | CCL-CCM-CCN-CCO |
| 10 | A | 310 | JTD | CCB-CCC-CCD-CCE |
| 10 | A | 310 | JTD | CBP-CBQ-CBR-CBS |

Continued on next page...


Continued from previous page...

| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|-----------------|
| 10 | A | 310 | JTD | CCE-CCF-CCG-CCH |
| 10 | A | 310 | JTD | CCJ-CCK-CCL-CCM |
| 8 | D | 603 | GOL | O2-C2-C3-O3 |
| 8 | A | 301 | GOL | O2-C2-C3-O3 |
| 8 | B | 101 | GOL | O1-C1-C2-O2 |
| 8 | B | 101 | GOL | O2-C2-C3-O3 |
| 10 | A | 310 | JTD | CBZ-CCA-CCB-CCC |
| 10 | A | 310 | JTD | CCM-CCN-CCO-CCP |
| 10 | A | 310 | JTD | O6-C2-C3-O2 |
| 10 | A | 310 | JTD | CCI-CCJ-CCK-CCL |
| 10 | A | 310 | JTD | CBK-CBL-CBM-CBN |
| 8 | D | 601 | GOL | O2-C2-C3-O3 |
| 8 | D | 602 | GOL | O1-C1-C2-C3 |
| 8 | D | 603 | GOL | O1-C1-C2-O2 |
| 10 | A | 310 | JTD | CCK-CCL-CCM-CCN |

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data [i](#)

5.1 Protein, DNA and RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates [i](#)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands [i](#)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers [i](#)

EDS failed to run properly - this section is therefore empty.