



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 10, 2020 – 10:17 AM BST

PDB ID : 5MUJ
Title : BT0996 RGII Chain B Complex
Authors : Cartmell, A.; Basle, A.; Ndeh, D.; Luis, A.S.; Venditto, I.; Labourel, A.; Rogowski, A.; Gilbert, H.J.
Deposited on : 2017-01-13
Resolution : 1.37 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2907 (1.40-1.36)
Clashscore	141614	3037 (1.40-1.36)
Ramachandran outliers	138981	2970 (1.40-1.36)
Sidechain outliers	138945	2969 (1.40-1.36)
RSRZ outliers	127900	2846 (1.40-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	381	 9% 85% 7% 8%
2	B	8	 38% 50% 13%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3281 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-galactosidase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	352	2834	1819	474	529	3	9	0	2	0


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	-	initiating methionine	UNP Q8A921

- Molecule 2 is an oligosaccharide called alpha-L-rhamnopyranose-(1-2)-[alpha-L-rhamnopyranose-(1-3)]alpha-L-arabinopyranose-(1-4)-[4-O-[(1R)-1-hydroxyethyl]-2-O-methyl-alpha-L-fucopyranose-(1-2)]beta-D-galactopyranose-(1-2)-alpha-D-aceric acid-(1-4)-alpha-L-rhamnopyranose-(1-3)-3-C-(hydroxymethyl)-alpha-D-erythrofuranoose.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	B	8	85	49	36	0	0	0

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	358	Total O 358 358	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-galactosidase



- Molecule 2: alpha-L-rhamnopyranose-(1-2)-[alpha-L-rhamnopyranose-(1-3)]alpha-L-arabinopyranose-(1-4)-[4-O-[(1R)-1-hydroxyethyl]-2-O-methyl-alpha-L-fucopyranose-(1-2)]beta-D-galactopyranose-(1-2)-alpha-D-aceric acid-(1-4)-alpha-L-rhamnopyranose-(1-3)-3-C-(hydroxymethyl)-alpha-D-erythrofuranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	49.46 Å 80.00 Å 50.81 Å 90.00° 114.94° 90.00°	Depositor
Resolution (Å)	44.85 – 1.37 44.85 – 1.37	Depositor EDS
% Data completeness (in resolution range)	99.4 (44.85-1.37) 99.4 (44.85-1.37)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.81 (at 1.37 Å)	Xtrriage
Refinement program	REFMAC 5.8.0155	Depositor
R, R_{free}	0.198 , 0.252 0.206 , 0.259	Depositor DCC
R_{free} test set	3603 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	15.2	Xtrriage
Anisotropy	0.735	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.035 for l,-k,h	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3281	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 8PK, 8OQ, RAM, ARA, GAL, ACT, XXM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.69	0/2918	0.86	3/3954 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	145	ARG	NE-CZ-NH2	6.13	123.37	120.30
1	A	61	ASP	CB-CG-OD1	5.44	123.20	118.30
1	A	145	ARG	NE-CZ-NH1	-5.16	117.72	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2834	0	2657	18	1
2	B	85	0	43	0	1
3	A	4	0	3	1	0
4	A	358	0	0	7	2
All	All	3281	0	2703	19	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:56:ASN:HD21	1:A:356:GLN:HE21	1.37	0.71
1:A:195:LYS:CB	4:A:795:HOH:O	2.45	0.65
1:A:190:LEU:HD23	4:A:508:HOH:O	1.98	0.62
1:A:104:ARG:HD2	4:A:774:HOH:O	2.07	0.54
1:A:56:ASN:ND2	1:A:356:GLN:HE21	2.04	0.54
1:A:189:ILE:C	1:A:190:LEU:HD12	2.28	0.53
1:A:232:HIS:CD2	1:A:234:GLU:H	2.29	0.51
1:A:146:HIS:HD2	4:A:535:HOH:O	1.96	0.49
3:A:409:ACT:H1	4:A:724:HOH:O	2.11	0.49
1:A:92:THR:HB	1:A:110:LEU:HB2	1.95	0.48
1:A:64:THR:HG23	1:A:336:TYR:CE2	2.49	0.47
1:A:67:ARG:NH1	1:A:69:GLY:O	2.47	0.47
1:A:66:PHE:HB2	4:A:512:HOH:O	2.15	0.46
1:A:232:HIS:HD2	1:A:234:GLU:H	1.63	0.45
1:A:191[A]:SER:HA	1:A:205:GLN:HE22	1.83	0.44
1:A:191[B]:SER:HA	1:A:205:GLN:HE22	1.84	0.43
1:A:199:TRP:CD2	1:A:233:PRO:HD3	2.54	0.42
1:A:224:MSE:O	1:A:244:ILE:HA	2.22	0.40
1:A:255:ARG:NE	4:A:508:HOH:O	2.54	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:379:THR:C	4:A:839:HOH:O[1_656]	1.90	0.30
2:B:1:XXM:O1	4:A:659:HOH:O[1_655]	1.95	0.25

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	352/381 (92%)	336 (96%)	16 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	289/309 (94%)	286 (99%)	3 (1%)	76	52

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	67	ARG
1	A	190	LEU
1	A	333	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	56	ASN
1	A	58	HIS
1	A	68	GLN
1	A	146	HIS
1	A	205	GLN
1	A	232	HIS
1	A	316	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	XXM	B	1	2	10,10,10	0.66	0	11,15,15	2.95	3 (27%)
2	RAM	B	2	2	10,10,11	0.46	0	14,14,16	2.41	4 (28%)
2	8OQ	B	3	2	7,11,12	1.02	1 (14%)	5,17,19	1.79	2 (40%)
2	GAL	B	4	2	11,11,12	0.43	0	15,15,17	0.96	0
2	ARA	B	5	2	9,9,10	0.73	0	10,12,14	1.81	2 (20%)
2	RAM	B	6	2	10,10,11	0.59	0	14,14,16	2.38	4 (28%)
2	RAM	B	7	2	10,10,11	0.28	0	14,14,16	0.83	0
2	8PK	B	8	2	14,14,15	0.64	0	18,19,21	0.93	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XXM	B	1	2	-	3/3/19/19	0/1/1/1
2	RAM	B	2	2	-	-	0/1/1/1
2	8OQ	B	3	2	-	0/0/23/26	0/1/1/1
2	GAL	B	4	2	-	0/2/19/22	0/1/1/1
2	ARA	B	5	2	-	-	0/1/1/1
2	RAM	B	6	2	-	-	0/1/1/1
2	RAM	B	7	2	-	-	0/1/1/1
2	8PK	B	8	2	-	1/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	3	8OQ	C3-C2	-2.30	1.51	1.54

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	XXM	O3-C3-C2	7.66	133.79	109.52
2	B	2	RAM	O4-C4-C3	-7.54	92.93	110.35
2	B	6	RAM	O5-C1-C2	6.11	120.20	110.77
2	B	1	XXM	O3-C3-C4	-4.64	99.65	109.05
2	B	6	RAM	C3-C4-C5	3.83	115.73	109.77
2	B	5	ARA	C5-O5-C1	3.47	116.85	111.52
2	B	5	ARA	C1-C2-C3	3.37	113.81	109.67
2	B	6	RAM	O5-C5-C4	3.31	115.46	109.52
2	B	3	8OQ	O2-C2-C1	3.03	118.34	110.26
2	B	2	RAM	C1-O5-C5	2.51	118.46	112.78
2	B	1	XXM	O4-C4-C3	-2.39	102.33	105.29
2	B	2	RAM	C3-C4-C5	2.34	113.42	109.77
2	B	2	RAM	O2-C2-C1	-2.23	104.58	109.15
2	B	3	8OQ	O2-C2-C3	2.19	117.45	112.57
2	B	6	RAM	C1-O5-C5	2.12	117.57	112.78

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	1	XXM	O3-C3-C5-O5
2	B	1	XXM	C4-C3-C5-O5
2	B	1	XXM	C2-C3-C5-O5
2	B	8	8PK	C1-C2-O2-C7

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1	XXM	0	1

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ACT	A	409	-	1,3,3	0.30	0	0,3,3	0.00	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	409	ACT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	343/381 (90%)	1.16	36 (10%) 6 6	10, 15, 25, 33	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	212	TRP	3.5
1	A	137	TYR	3.4
1	A	49[A]	LEU	3.3
1	A	176	ILE	3.0
1	A	281	ASP	2.9
1	A	307	TYR	2.9
1	A	70	ASP	2.8
1	A	199	TRP	2.8
1	A	94	ILE	2.7
1	A	101	LEU	2.7
1	A	103	TRP	2.6
1	A	125	PHE	2.6
1	A	326	ASN	2.5
1	A	237	LEU	2.5
1	A	349	CYS	2.4
1	A	75	THR	2.4
1	A	158	TYR	2.4
1	A	361	ILE	2.4
1	A	235	THR	2.4
1	A	32	GLU	2.4
1	A	47	TYR	2.4
1	A	312	TRP	2.3
1	A	253	TRP	2.3
1	A	214	LYS	2.2
1	A	293	PRO	2.2
1	A	256	TYR	2.2
1	A	93	TRP	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	203	LEU	2.1
1	A	65	VAL	2.1
1	A	138	GLU	2.1
1	A	30	THR	2.1
1	A	62	CYS	2.1
1	A	351	VAL	2.1
1	A	338	ILE	2.0
1	A	234	GLU	2.0
1	A	200	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	RAM	B	2	10/11	0.70	0.27	30,40,48,49	0
2	XXM	B	1	10/10	0.73	0.31	22,23,32,36	0
2	8PK	B	8	14/15	0.78	0.22	24,27,30,31	0
2	RAM	B	7	10/11	0.80	0.19	21,24,26,29	0
2	GAL	B	4	11/12	0.87	0.19	19,21,24,24	0
2	8OQ	B	3	11/12	0.90	0.16	24,29,36,39	0
2	ARA	B	5	9/10	0.90	0.11	15,16,20,22	0
2	RAM	B	6	10/11	0.91	0.17	14,15,16,16	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ACT	A	409	4/4	0.73	0.16	18,20,21,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.