



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 19, 2023 – 06:43 AM EDT

PDB ID : 2NS8  
Title : How an in vitro selected peptide mimics the antibiotic tetracycline to induce TET repressor  
Authors : Luckner, S.R.; Klotzsche, M.; Berens, C.; Hillen, W.; Muller, Y.A.  
Deposited on : 2006-11-03  
Resolution : 2.55 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1284 (2.56-2.52)
Clashscore	141614	1332 (2.56-2.52)
Ramachandran outliers	138981	1315 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)
RSRZ outliers	127900	1272 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	208	
1	B	208	
1	C	208	
1	D	208	
2	E	17	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	17	
2	G	17	
2	H	17	
2	Z	17	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6884 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tetracycline repressor protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	200	1599	1009	285	303	2	0	0	0
1	B	204	1626	1029	287	308	2	0	0	0
1	C	206	1643	1038	292	311	2	0	0	0
1	D	198	1577	996	280	299	2	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	68	SER	CYS	engineered mutation	UNP P04483
A	88	ASN	CYS	engineered mutation	UNP P04483
A	121	THR	CYS	engineered mutation	UNP P04483
A	144	SER	CYS	engineered mutation	UNP P04483
B	68	SER	CYS	engineered mutation	UNP P04483
B	88	ASN	CYS	engineered mutation	UNP P04483
B	121	THR	CYS	engineered mutation	UNP P04483
B	144	SER	CYS	engineered mutation	UNP P04483
C	68	SER	CYS	engineered mutation	UNP P04483
C	88	ASN	CYS	engineered mutation	UNP P04483
C	121	THR	CYS	engineered mutation	UNP P04483
C	144	SER	CYS	engineered mutation	UNP P04483
D	68	SER	CYS	engineered mutation	UNP P04483
D	88	ASN	CYS	engineered mutation	UNP P04483
D	121	THR	CYS	engineered mutation	UNP P04483
D	144	SER	CYS	engineered mutation	UNP P04483


- Molecule 2 is a protein called 16 residue peptide Tip (Transcription inducing peptide).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	H	13	Total	C	N	O	0	0	0
			102	70	15	17			
2	E	12	Total	C	N	O	0	0	0
			96	67	14	15			
2	F	13	Total	C	N	O	0	0	0
			102	70	15	17			
2	G	12	Total	C	N	O	0	0	0
			96	67	14	15			
2	Z	4	Total	C	N	O	0	0	0
			43	30	7	6			


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tetracycline repressor protein



- Molecule 1: Tetracycline repressor protein




- Molecule 1: Tetracycline repressor protein



- Molecule 1: Tetracycline repressor protein





- Molecule 2: 16 residue peptide Tip (Transcription inducing peptide)



- Molecule 2: 16 residue peptide Tip (Transcription inducing peptide)



- Molecule 2: 16 residue peptide Tip (Transcription inducing peptide)



- Molecule 2: 16 residue peptide Tip (Transcription inducing peptide)



- Molecule 2: 16 residue peptide Tip (Transcription inducing peptide)



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	77.89Å 200.59Å 65.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.55 38.23 – 2.55	Depositor EDS
% Data completeness (in resolution range)	100.0 (30.00-2.55) 99.5 (38.23-2.55)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.57 (at 2.54Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.206 , 0.270 0.214 , 0.274	Depositor DCC
$R_{free}$ test set	2715 reflections (8.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	39.1	Xtrriage
Anisotropy	0.439	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 46.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	6884	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.38% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.66	0/1631	0.70	0/2206
1	B	0.70	0/1658	0.79	2/2244 (0.1%)
1	C	0.61	0/1675	0.71	1/2266 (0.0%)
1	D	0.60	0/1607	0.69	0/2171
2	E	0.60	0/100	0.66	0/140
2	F	0.58	0/106	0.53	0/148
2	G	0.53	0/100	0.63	0/140
2	H	0.70	0/106	0.64	0/148
2	Z	0.89	0/46	0.59	0/64
All	All	0.64	0/7029	0.72	3/9527 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	171	ARG	NE-CZ-NH2	-6.69	116.95	120.30
1	C	195	ARG	NE-CZ-NH1	5.82	123.21	120.30
1	B	171	ARG	NE-CZ-NH1	5.46	123.03	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	164	ASP	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1599	0	1583	19	0
1	B	1626	0	1616	30	0
1	C	1643	0	1634	26	0
1	D	1577	0	1555	26	0
2	E	96	0	81	0	0
2	F	102	0	86	4	0
2	G	96	0	81	6	0
2	H	102	0	86	2	0
2	Z	43	0	35	0	0
All	All	6884	0	6757	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 94 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:0:ACE:H1	2:F:1:TRP:HB2	1.53	0.91
1:C:10:ILE:HG21	1:C:58:GLU:HG3	1.61	0.83
1:C:206:GLN:O	1:C:207:ILE:HG23	1.88	0.74
1:A:26:THR:OG1	1:A:29:LYS:HD2	1.90	0.71
1:C:200:GLN:HE21	1:D:188:HIS:HE1	1.39	0.70

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	198/208 (95%)	183 (92%)	13 (7%)	2 (1%)	15	22
1	B	202/208 (97%)	197 (98%)	3 (2%)	2 (1%)	15	22
1	C	204/208 (98%)	197 (97%)	5 (2%)	2 (1%)	15	22
1	D	194/208 (93%)	181 (93%)	10 (5%)	3 (2%)	10	14
2	E	10/17 (59%)	9 (90%)	1 (10%)	0	100	100
2	F	11/17 (65%)	9 (82%)	2 (18%)	0	100	100
2	G	10/17 (59%)	9 (90%)	1 (10%)	0	100	100
2	H	11/17 (65%)	11 (100%)	0	0	100	100
2	Z	2/17 (12%)	2 (100%)	0	0	100	100
All	All	842/917 (92%)	798 (95%)	35 (4%)	9 (1%)	14	19

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	160	THR
1	B	165	SER
1	A	162	THR
1	C	64	HIS
1	C	161	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	172/179 (96%)	155 (90%)	17 (10%)	8	9
1	B	175/179 (98%)	156 (89%)	19 (11%)	6	6
1	C	177/179 (99%)	157 (89%)	20 (11%)	6	5
1	D	168/179 (94%)	149 (89%)	19 (11%)	6	5

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	E	7/9 (78%)	7 (100%)	0	100	100
2	F	8/9 (89%)	8 (100%)	0	100	100
2	G	7/9 (78%)	7 (100%)	0	100	100
2	H	8/9 (89%)	8 (100%)	0	100	100
2	Z	4/9 (44%)	3 (75%)	1 (25%)	0	0
All	All	726/761 (95%)	650 (90%)	76 (10%)	7	7

5 of 76 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	14	LEU
1	D	156	GLU
1	D	30	LEU
1	D	65	THR
2	Z	1	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	100	HIS
1	C	206	GLN
1	D	188	HIS
1	C	180	GLN
1	D	11	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	200/208 (96%)	0.57	16 (8%) 12 15	38, 48, 63, 73	0
1	B	204/208 (98%)	0.58	13 (6%) 19 22	42, 47, 56, 60	0
1	C	206/208 (99%)	0.74	22 (10%) 6 7	36, 46, 58, 70	0
1	D	198/208 (95%)	0.80	20 (10%) 7 8	38, 47, 59, 71	0
2	E	11/17 (64%)	1.27	2 (18%) 1 1	48, 50, 57, 59	0
2	F	12/17 (70%)	1.49	4 (33%) 0 0	49, 53, 60, 61	0
2	G	11/17 (64%)	0.95	1 (9%) 9 11	48, 50, 54, 58	0
2	H	12/17 (70%)	1.63	4 (33%) 0 0	45, 47, 54, 55	0
2	Z	4/17 (23%)	2.93	3 (75%) 0 0	53, 54, 55, 56	0
All	All	858/917 (93%)	0.72	85 (9%) 7 9	36, 47, 59, 73	0

The worst 5 of 85 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	163	THR	6.3
1	A	161	PRO	6.1
2	E	10	ALA	6.0
1	C	2	SER	5.6
2	Z	2	THR	5.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.