Full wwPDB X-ray Structure Validation Report (i)

Jan 7, 2024 – 06:26 pm GMT

PDB ID	:	5O22
Title	:	E. coli FolD in complex with carolacton
Authors	:	Koehnke, J.; Sikandar, A.
Deposited on	:	2017-05-19
Resolution	:	2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	289	2%, 90%, 7%
2	B	289	3%, 93%
3	C	289	%, 94%
4	D	289	6%, 87%, 8%, 6%

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 18203 atoms, of which 8881 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bifunctional protein FolD.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	280	4338	1353	2199	379	397	10	0	3	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP P24186
A	1	GLY	-	expression tag	UNP P24186

- Molecule 2 is a protein called Bifunctional protein FolD.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	B	280	4342	1356	2203	376	396	11	8	3	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P24186
B	1	GLY	-	expression tag	UNP P24186

- Molecule 3 is a protein called Bifunctional protein FolD.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
3	C	284	4353	1357	2206	380	401	9	0	1	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	MET	-	initiating methionine	UNP P24186
C	1	GLY	-	expression tag	UNP P24186

- Molecule 4 is a protein called Bifunctional protein FolD.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
4	D	273	4180	1305	2117	366	384	8	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	0	MET	-	initiating methionine	UNP P24186
D	1	GLY	-	expression tag	UNP P24186

- Molecule 5 is Carolacton (three-letter code: C3R) (formula: $C_{25}H_{40}O_8$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
5	A	1	72	25	39	8	0	0
5	B	1	72	25	39	8	0	0
5	C	1	72	25	39	8	0	0
5	D	1	72	25	39	8	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	216	216	216	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	B	173	173	173	0	0
			Total	O		
6	C	216	216	216	0	0
			Total	O		
6	D	97	97	97	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bifunctional protein FolD

Chain A:





- Molecule 2: Bifunctional protein FolD

Chain B:





- Molecule 3: Bifunctional protein FolD

Chain C:





- Molecule 4: Bifunctional protein FolD

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	99.64Å 81.02Å 100.94Å 90.00° 112.95° 90.00°	Depositor
Resolution (Å)	45.88 – 2.10 45.88 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.8 (45.88-2.10) 98.8 (45.88-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.97 (at 2.10Å)	Xtriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.175 , 0.206 0.175 , 0.205	Depositor DCC
R_{free} test set	4258 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å^2)	30.8	Xtriage
Anisotropy	0.214	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.38 , 46.9	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.023 for l,-k,h	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	18203	wwPDB-VP
Average B, all atoms (Å^2)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.11% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: C3R, MLY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.32	0/2161	0.50	0/2943
2	B	0.34	0/2152	0.53	1/2930 (0.0%)
3	C	0.32	0/2175	0.48	0/2964
4	D	0.29	0/2096	0.48	0/2852
All	All	0.32	0/8584	0.50	1/11689 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1
3	C	0	1
All	All	0	2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	194	MLY	O-C-N	-6.58	112.18	122.70

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	194	MLY	Mainchain
3	C	22	MLY	Mainchain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2139	2199	2199	14	0
2	B	2139	2203	2205	7	0
3	C	2147	2206	2203	9	0
4	D	2063	2117	2119	17	0
5	A	33	39	0	0	0
5	B	33	39	0	0	0
5	C	33	39	0	0	0
5	D	33	39	0	0	0
6	A	216	0	0	5	0
6	B	173	0	0	2	0
6	C	216	0	0	7	0
6	D	97	0	0	6	0
All	All	9322	8881	8726	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:60:GLU:OE2	6:C:401:HOH:O	1.92	0.86
2:B:194:MLY:O	6:B:401:HOH:O	1.94	0.84
4:D:159:GLY:O	6:D:401:HOH:O	1.96	0.83
4:D:139:ARG:NH1	6:D:405:HOH:O	2.13	0.82
4:D:76:GLU:OE2	6:D:402:HOH:O	2.03	0.76
3:C:236:GLU:OE1	6:C:403:HOH:O	2.04	0.75
3:C:60:GLU:OE2	6:C:402:HOH:O	2.03	0.75
2:B:49:ILE:N	6:B:402:HOH:O	2.21	0.73
1:A:14:GLN:OE1	6:A:401:HOH:O	2.07	0.72
3:C:2:ALA:O	6:C:404:HOH:O	2.06	0.72
1:A:50:TYR:N	6:A:405:HOH:O	2.23	0.71
4:D:273:GLU:OE1	6:D:403:HOH:O	2.08	0.71
1:A:76:GLU:OE1	6:A:402:HOH:O	2.09	0.70
4:D:42:GLY:O	6:D:404:HOH:O	2.13	0.67
1:A:151:GLU:OE2	6:A:403:HOH:O	2.11	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:139:ARG:NH2	1:A:178:GLU:OE1	2.34	0.60
3:C:55:ARG:NH1	3:C:67:SER:OG	2.36	0.59
1:A:204:ASP:OD1	2:B:194:MLY:CH1	2.51	0.58
4:D:100:PRO:HA	4:D:101:LEU:HB2	1.84	0.58
3:C:53:SER:HA	3:C:56:LYS:HD3	1.87	0.56
1:A:132[B]:CYS:SG	1:A:177[B]:MET:SD	3.06	0.54
4:D:14:GLN:O	4:D:18:GLU:HG3	2.08	0.53
3:C:107:ASN:ND2	6:C:405:HOH:O	2.10	0.52
3:C:161:ASN:HB3	6:C:567:HOH:O	2.09	0.51
4:D:53:SER:HA	6:D:429:HOH:O	2.10	0.51
4:D:149:LEU:HG	4:D:266:MET:HG3	1.94	0.49
4:D:18:GLU:OE2	4:D:152:ARG:NH2	2.44	0.47
1:A:174:PRO:HA	1:A:177[B]:MET:HE3	1.96	0.47
2:B:66:ARG:CZ	2:B:92:ILE:HD11	2.45	0.47
4:D:195:ASN:OD1	4:D:195:ASN:N	2.48	0.47
3:C:255:TYR:OH	6:C:406:HOH:O	2.21	0.46
4:D:100:PRO:HA	4:D:101:LEU:CB	2.46	0.45
1:A:73:THR:OG1	6:A:404:HOH:O	2.20	0.45
4:D:245:VAL:HG23	4:D:245:VAL:O	2.17	0.45
4:D:142:THR:HB	4:D:143:PRO:HD3	1.98	0.45
4:D:176:SER:HB2	4:D:186:THR:HG21	1.99	0.44
2:B:37:ALA:HB2	2:B:92:ILE:HD12	2.01	0.43
1:A:204:ASP:OD1	2:B:194:MLY:HH12	2.17	0.43
1:A:142:THR:HB	1:A:143:PRO:HD3	2.00	0.43
4:D:144:ARG:HD3	4:D:270:THR:HG23	2.02	0.42
2:B:142:THR:HB	2:B:143:PRO:HD3	2.01	0.41
4:D:133:GLN:O	4:D:134:ARG:HG2	2.19	0.41
1:A:230:VAL:HG12	1:A:230:VAL:O	2.20	0.41
1:A:217:PRO:HG2	1:A:220:TRP:CE2	2.56	0.41
1:A:179:LEU:HD11	1:A:207:ILE:HD12	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	277/289 (96%)	268 (97%)	9 (3%)	0	100	100
2	B	277/289 (96%)	269 (97%)	8 (3%)	0	100	100
3	C	282/289 (98%)	275 (98%)	7 (2%)	0	100	100
4	D	267/289 (92%)	258 (97%)	8 (3%)	1 (0%)	34	32
All	All	1103/1156 (95%)	1070 (97%)	32 (3%)	1 (0%)	51	54

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	D	101	LEU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/235 (98%)	229 (99%)	2 (1%)	78	84
2	B	230/234 (98%)	225 (98%)	5 (2%)	52	57
3	C	232/236 (98%)	230 (99%)	2 (1%)	78	84
4	D	223/237 (94%)	222 (100%)	1 (0%)	91	94
All	All	916/942 (97%)	906 (99%)	10 (1%)	76	79

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	65	SER
1	A	107	ASN
2	B	49	ILE
2	B	123[A]	PHE
2	B	123[B]	PHE
2	B	219	ASP
2	B	284	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	C	55	ARG
3	C	219	ASP
4	D	101	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
1	MLY	A	194	1	9,10,11	0.58	0	6,11,13	0.55	0
2	MLY	B	212	2	9,10,11	0.70	0	6,11,13	0.53	0
2	MLY	B	194	2	9,10,11	0.60	0	6,11,13	1.32	1 (16%)
3	MLY	C	22	3	9,10,11	0.59	0	6,11,13	0.75	0
1	MLY	A	222	1	9,10,11	0.61	0	6,11,13	0.75	0
2	MLY	B	222	2	9,10,11	0.59	0	6,11,13	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MLY	A	194	1	-	1/8/9/11	-
2	MLY	B	212	2	-	0/8/9/11	-
2	MLY	B	194	2	-	3/8/9/11	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MLY	C	22	3	-	1/8/9/11	-
1	MLY	A	222	1	-	0/8/9/11	-
2	MLY	B	222	2	-	0/8/9/11	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	194	MLY	CH2-NZ-CH1	-2.01	104.53	109.73

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	194	MLY	CD-CE-NZ-CH1
2	B	194	MLY	CD-CE-NZ-CH2
2	B	194	MLY	CG-CD-CE-NZ
1	A	194	MLY	CG-CD-CE-NZ
3	C	22	MLY	CG-CD-CE-NZ

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	194	MLY	3	0

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	C3R	B	301	-	33,33,33	0.88	2 (6%)	35,45,45	1.24	4 (11%)
5	C3R	A	301	-	33,33,33	0.87	0	35,45,45	1.18	4 (11%)
5	C3R	D	301	-	33,33,33	1.01	2 (6%)	35,45,45	1.22	4 (11%)
5	C3R	C	301	-	33,33,33	1.22	4 (12%)	35,45,45	1.54	5 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	C3R	B	301	-	-	8/53/53/53	0/0/1/1
5	C3R	A	301	-	-	3/53/53/53	0/0/1/1
5	C3R	D	301	-	-	6/53/53/53	0/0/1/1
5	C3R	C	301	-	-	12/53/53/53	0/0/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	301	C3R	C19-C18	-2.78	1.50	1.54
5	D	301	C3R	C21-C23	2.77	1.58	1.53
5	B	301	C3R	O06-C05	-2.51	1.38	1.43
5	C	301	C3R	C21-C23	2.33	1.57	1.53
5	B	301	C3R	C07-C05	2.30	1.61	1.53
5	C	301	C3R	C18-C20	2.16	1.55	1.53
5	D	301	C3R	C26-C27	2.14	1.56	1.51
5	C	301	C3R	O06-C05	-2.09	1.39	1.43

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	C	301	C3R	C19-C18-C17	3.94	114.44	110.75
5	C	301	C3R	C14-C15-C17	3.42	131.37	120.93
5	B	301	C3R	O08-C07-C32	-3.25	101.20	110.17
5	C	301	C3R	C16-C15-C17	-3.07	115.38	123.45
5	D	301	C3R	O08-C07-C32	-2.91	102.13	110.17
5	A	301	C3R	O08-C07-C32	-2.84	102.32	110.17

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	B	301	C3R	C19-C18-C20	2.61	113.30	109.66
5	A	301	C3R	C19-C18-C20	2.59	113.28	109.66
5	C	301	C3R	O30-C20-C18	-2.58	116.28	120.76
5	D	301	C3R	O30-C20-C18	-2.43	116.55	120.76
5	C	301	C3R	O08-C07-C32	-2.40	103.54	110.17
5	A	301	C3R	C05-C04-C03	2.32	128.58	124.93
5	D	301	C3R	C19-C18-C20	2.17	112.69	109.66
5	D	301	C3R	O31-C32-C07	2.16	114.69	111.15
5	A	301	C3R	O31-C32-C07	2.16	114.69	111.15
5	B	301	C3R	O30-C20-C18	-2.15	117.02	120.76
5	B	301	C3R	C05-C04-C03	2.07	128.18	124.93

There are no chirality outliers.

All (29) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	301	C3R	C15-C17-C18-C19
5	C	301	C3R	C17-C18-C20-O30
5	C	301	C3R	C09-C02-C03-C04
5	C	301	C3R	C03-C04-C05-O06
5	C	301	C3R	C03-C04-C05-C07
5	C	301	C3R	C11-C12-C14-O31
5	D	301	C3R	C03-C04-C05-C07
5	D	301	C3R	C11-C12-C14-O31
5	D	301	C3R	C02-C09-C10-C11
5	C	301	C3R	C02-C09-C10-C11
5	C	301	C3R	C14-C15-C17-C18
5	A	301	C3R	C10-C11-C12-C14
5	C	301	C3R	C01-C02-C03-C04
5	D	301	C3R	C03-C04-C05-O06
5	C	301	C3R	C16-C15-C17-C18
5	C	301	C3R	O24-C23-C26-C27
5	D	301	C3R	C13-C12-C14-O31
5	C	301	C3R	C17-C18-C20-C21
5	B	301	C3R	C02-C09-C10-C11
5	B	301	C3R	C04-C05-C07-O08
5	B	301	C3R	C03-C04-C05-O06
5	B	301	C3R	C03-C04-C05-C07
5	B	301	C3R	C09-C02-C03-C04
5	A	301	C3R	C05-C07-C32-O31
5	D	301	C3R	O08-C07-C32-O31
5	B	301	C3R	O06-C05-C07-O08

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	B	301	C3R	O08-C07-C32-O31
5	A	301	C3R	C05-C07-C32-O33
5	B	301	C3R	C03-C02-C09-C10

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand C3R B 301



Ligand C3R A 301



Ligand C3R D 301



Ligand C3R C 301



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	278/289 (96%)	-0.22	7 (2%)	57	62	20, 30, 64, 97	0
2	B	277/289 (95%)	-0.14	8 (2%)	51	57	21, 35, 66, 91	0
3	C	283/289 (97%)	-0.18	4 (1%)	75	78	23, 35, 68, 93	0
4	D	273/289 (94%)	0.20	17 (6%)	20	25	27, 51, 79, 100	0
All	All	1111/1156 (96%)	-0.09	36 (3%)	47	54	20, 37, 73, 100	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	D	240	VAL	5.8
4	D	241	VAL	4.3
4	D	28	ALA	4.1
1	A	236	GLU	4.0
1	A	237	ASN	4.0
4	D	51	VAL	3.8
1	A	239	LYS	3.7
2	B	236	GLU	3.6
4	D	31	LEU	3.4
4	D	285	PRO	3.2
2	B	55	ARG	3.1
2	B	49	ILE	3.1
1	A	103	ALA	3.0
4	D	50	TYR	3.0
4	D	234	ARG	2.9
4	D	55	ARG	2.9
2	B	43	SER	2.7
3	C	50	TYR	2.7
1	A	235	LEU	2.7
4	D	259	VAL	2.7
4	D	232	ILE	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	239	LYS	2.6
4	D	246	PHE	2.6
4	D	27	ILE	2.6
4	D	1	GLY	2.5
3	C	55	ARG	2.5
4	D	53	SER	2.4
3	C	28	ALA	2.4
4	D	284	ASP	2.4
2	B	51	VAL	2.3
1	A	286	GLN	2.2
2	B	237	ASN	2.1
4	D	30	GLY	2.1
1	A	285	PRO	2.0
2	B	123[A]	PHE	2.0
3	C	30	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	MLY	B	194	11/12	0.89	0.16	32,66,85,85	0
3	MLY	C	22	11/12	0.91	0.20	36,69,83,85	0
2	MLY	B	212	11/12	0.94	0.10	31,48,59,59	0
1	MLY	A	194	11/12	0.95	0.13	22,36,60,60	0
2	MLY	B	222	11/12	0.98	0.13	30,44,59,59	0
1	MLY	A	222	11/12	0.98	0.11	25,37,50,50	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	C3R	C	301	33/33	0.81	0.26	41,113,143,152	0
5	C3R	D	301	33/33	0.83	0.27	62,106,130,131	0
5	C3R	A	301	33/33	0.93	0.13	30,51,68,72	0
5	C3R	B	301	33/33	0.93	0.13	32,48,62,64	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around C3R C 301:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)







Electron density around C3R D 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around C3R A 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around C3R B 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.