



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 12, 2023 – 11:02 PM EDT

PDB ID : 4OFW
Title : Crystal Structure of Arabidopsis thaliana DJ-1d
Authors : Choi, D.; Kim, J.; Ryu, K.-S.; Park, C.
Deposited on : 2014-01-15
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	387	
1	B	387	
1	C	387	
1	D	387	
1	E	387	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	387	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into two segments: a green segment on the left labeled '63%' and a yellow segment on the right labeled '34%'. A small red square is at the beginning of the bar, and a small black dot is at the end. A '%' symbol is positioned above the start of the bar.</p>

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 18086 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein DJ-1 homolog D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	387	2924	1864	487	554	19	0	0	0
1	B	387	2924	1864	487	554	19	0	0	0
1	C	387	2924	1864	487	554	19	0	0	0
1	D	387	2924	1864	487	554	19	0	0	0
1	E	387	2924	1864	487	554	19	0	0	0
1	F	387	2924	1864	487	554	19	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	238	GLU	ASP	conflict	UNP Q9M8R4
A	389	HIS	-	expression tag	UNP Q9M8R4
B	238	GLU	ASP	conflict	UNP Q9M8R4
B	389	HIS	-	expression tag	UNP Q9M8R4
C	238	GLU	ASP	conflict	UNP Q9M8R4
C	389	HIS	-	expression tag	UNP Q9M8R4
D	238	GLU	ASP	conflict	UNP Q9M8R4
D	389	HIS	-	expression tag	UNP Q9M8R4
E	238	GLU	ASP	conflict	UNP Q9M8R4
E	389	HIS	-	expression tag	UNP Q9M8R4
F	238	GLU	ASP	conflict	UNP Q9M8R4
F	389	HIS	-	expression tag	UNP Q9M8R4


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	97	Total O 97 97	0	0
2	B	81	Total O 81 81	0	0
2	C	96	Total O 96 96	0	0
2	D	100	Total O 100 100	0	0
2	E	82	Total O 82 82	0	0
2	F	86	Total O 86 86	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Protein DJ-1 homolog D




- Molecule 1: Protein DJ-1 homolog D



- Molecule 1: Protein DJ-1 homolog D





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	105.56Å 115.05Å 115.53Å 90.00° 91.51° 90.00°	Depositor
Resolution (Å)	50.00 – 2.30 47.96 – 2.30	Depositor EDS
% Data completeness (in resolution range)	95.9 (50.00-2.30) 95.9 (47.96-2.30)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.62 (at 2.29Å)	Xtrriage
Refinement program	CNS, REFMAC	Depositor
R, R_{free}	0.243 , 0.284 0.243 , 0.284	Depositor DCC
R_{free} test set	11742 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	24.0	Xtrriage
Anisotropy	0.678	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 43.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.000 for -h,l,k 0.000 for -h,-l,-k 0.029 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	18086	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 46.23 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1795e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/2994	0.62	0/4075
1	B	0.37	0/2994	0.62	0/4075
1	C	0.39	0/2994	0.64	0/4075
1	D	0.39	0/2994	0.62	0/4075
1	E	0.38	0/2994	0.63	0/4075
1	F	0.39	0/2994	0.64	0/4075
All	All	0.38	0/17964	0.63	0/24450

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2924	0	2872	131	0
1	B	2924	0	2872	119	0
1	C	2924	0	2872	110	0
1	D	2924	0	2872	122	0
1	E	2924	0	2872	116	0
1	F	2924	0	2872	134	0
2	A	97	0	0	22	0
2	B	81	0	0	11	0
2	C	96	0	0	14	0
2	D	100	0	0	20	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	82	0	0	13	0
2	F	86	0	0	15	0
All	All	18086	0	17232	701	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 701 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:197:ASN:HA	2:B:468:HOH:O	1.59	1.00
1:F:133:ASN:HA	2:F:460:HOH:O	1.66	0.95
1:B:20:VAL:HB	2:B:455:HOH:O	1.66	0.94
1:F:353:ILE:HD12	1:F:353:ILE:H	1.28	0.94
1:D:7:VAL:HG23	2:D:486:HOH:O	1.70	0.91

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	385/387 (100%)	353 (92%)	27 (7%)	5 (1%)	12	12
1	B	385/387 (100%)	356 (92%)	27 (7%)	2 (0%)	29	35
1	C	385/387 (100%)	356 (92%)	28 (7%)	1 (0%)	41	50
1	D	385/387 (100%)	358 (93%)	24 (6%)	3 (1%)	19	23
1	E	385/387 (100%)	352 (91%)	31 (8%)	2 (0%)	29	35
1	F	385/387 (100%)	356 (92%)	24 (6%)	5 (1%)	12	12
All	All	2310/2322 (100%)	2131 (92%)	161 (7%)	18 (1%)	19	23

5 of 18 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	75	ASP
1	E	238	GLU
1	A	74	PHE
1	A	236	ALA
1	D	275	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	313/313 (100%)	298 (95%)	15 (5%)	25	36
1	B	313/313 (100%)	300 (96%)	13 (4%)	30	42
1	C	313/313 (100%)	297 (95%)	16 (5%)	24	33
1	D	313/313 (100%)	304 (97%)	9 (3%)	42	58
1	E	313/313 (100%)	304 (97%)	9 (3%)	42	58
1	F	313/313 (100%)	300 (96%)	13 (4%)	30	42
All	All	1878/1878 (100%)	1803 (96%)	75 (4%)	31	44

5 of 75 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	147	LEU
1	F	223	LEU
1	E	197	ASN
1	F	64	ARG
1	B	251	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 49 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	259	HIS
1	E	52	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	266	ASN
1	D	377	GLN
1	E	183	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	387/387 (100%)	-0.05	3 (0%) 86 89	16, 30, 48, 69	0
1	B	387/387 (100%)	-0.09	2 (0%) 91 94	14, 33, 49, 68	0
1	C	387/387 (100%)	-0.20	2 (0%) 91 94	15, 28, 46, 68	0
1	D	387/387 (100%)	-0.08	4 (1%) 82 86	18, 31, 49, 69	0
1	E	387/387 (100%)	-0.21	1 (0%) 94 96	15, 31, 47, 68	0
1	F	387/387 (100%)	-0.10	5 (1%) 77 81	14, 31, 48, 67	0
All	All	2322/2322 (100%)	-0.12	17 (0%) 87 91	14, 31, 48, 69	0

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	389	HIS	4.1
1	B	389	HIS	4.1
1	A	3	ASN	3.4
1	D	389	HIS	3.0
1	C	389	HIS	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.