wwPDB X-ray Structure Validation Summary Report (i)

Dec 16, 2023 – 08:36 AM EST

PDB ID	:	2OPZ
Title	:	AVPF bound to BIR3-XIAP
Authors	:	Wist, A.D.
Deposited on	:	2007-01-30
Resolution	:	3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	109	59% 33% 6% •
1	B	109	55% 40% • •
1	C	109	42% 52% • •
1	D	109	54% 40% 5% •
2	E	4	50% 50%
2	F	4	50% 50%
2	G	4	50% 50%
2	H	4	75% 25%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3680 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Baculoviral IAP repeat-containing protein 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	109	888	566	151	166	5	0	0	0
1	B	109	888	566	151	166	5	0	0	0
1	C	109	888	566	151	166	5	0	0	0
1	D	109	888	566	151	166	5	0	0	0

- Molecule 2 is a protein called AVPF (Smac homologue, N-terminal tetrapeptide).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	E	4	31	22	4	5	0	0	0
2	F	4	31	22	4	5	0	0	0
2	G	4	31	22	4	5	0	0	0
2	H	4	31	22	4	5	0	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
3	A	1	1	1	0	0
3	B	1	1	1	0	0
3	C	1	1	1	0	0
3	D	1	1	1	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: Baculoviral IAP repeat-containing protein 4

Chain A:



• Molecule 1: Baculoviral IAP repeat-containing protein 4

Chain B:



• Molecule 1: Baculoviral IAP repeat-containing protein 4

Chain C:



• Molecule 1: Baculoviral IAP repeat-containing protein 4

Chain D:





• Molecule 2: AVPF (Smac homologue, N-terminal tetrapeptide)

Chain E:

• Molecule 2: AVPF (Smac homologue, N-terminal tetrapeptide)

Chain F:

• Molecule 2: AVPF (Smac homologue, N-terminal tetrapeptide)

Chain G:

• Molecule 2: AVPF (Smac homologue, N-terminal tetrapeptide)

Chain H:

4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	I 21 3	Depositor
Cell constants a, b, c, α, β, γ	170.37Å 170.37Å 170.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.00	Depositor
% Data completeness (in resolution range)	95.9 (50.00-3.00)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.231 , 0.277	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3680	wwPDB-VP
Average B, all atoms (Å^2)	72.0	wwPDB-VP

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	0.55	0/916	0.80	1/1243 (0.1%)
1	B	0.55	0/916	0.83	1/1243 (0.1%)
1	C	0.60	0/916	0.85	2/1243 (0.2%)
1	D	0.58	0/916	0.84	1/1243 (0.1%)
2	E	0.90	0/32	0.93	0/42
2	F	0.75	0/32	0.49	0/42
2	G	0.67	0/32	0.61	0/42
2	H	0.61	0/32	0.62	0/42
All	All	0.58	0/3792	0.82	5/5140 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	C	250	PHE	N-CA-C	-10.02	83.94	111.00
1	B	251	PRO	N-CA-C	9.25	136.14	112.10
1	A	252	ASN	N-CA-C	-8.52	88.01	111.00
1	C	251	PRO	N-CA-C	7.94	132.75	112.10
1	D	251	PRO	N-CA-C	7.67	132.04	112.10

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	277	TYR	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	888	0	827	43	0
1	B	888	0	827	48	0
1	C	888	0	827	62	0
1	D	888	0	827	48	0
2	E	31	0	32	2	0
2	F	31	0	32	2	0
2	G	31	0	32	2	0
2	H	31	0	32	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
All	All	3680	0	3436	200	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

The worst 5 of 200 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:265:TYR:CE2	1:C:269:ILE:HG12	2.01	0.95
1:B:254:THR:HG22	1:B:328:LYS:HD2	1.57	0.84
1:C:251:PRO:O	1:C:255:ASN:HB2	1.78	0.84
1:D:265:TYR:CE2	1:D:269:ILE:HG12	2.14	0.82
1:D:265:TYR:HE2	1:D:269:ILE:HG12	1.47	0.80

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	107/109 (98%)	93 (87%)	9 (8%)	5 (5%)	2 14
1	B	107/109 (98%)	96 (90%)	9 (8%)	2 (2%)	8 36
1	C	107/109 (98%)	99 (92%)	6 (6%)	2 (2%)	8 36
1	D	107/109 (98%)	94 (88%)	11 (10%)	2 (2%)	8 36
2	E	2/4 (50%)	2 (100%)	0	0	100 100
2	F	2/4 (50%)	2 (100%)	0	0	100 100
2	G	2/4 (50%)	2 (100%)	0	0	100 100
2	H	2/4 (50%)	2 (100%)	0	0	100 100
All	All	436/452 (96%)	390 (89%)	35 (8%)	11 (2%)	5 28

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	253	SER
1	A	254	THR
1	B	255	ASN
1	C	251	PRO
1	D	251	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	94/94 (100%)	86 (92%)	8 (8%)	10 38

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	94/94 (100%)	89 (95%)	5 (5%)	22	58
1	C	94/94 (100%)	86 (92%)	8 (8%)	10	38
1	D	94/94 (100%)	86 (92%)	8 (8%)	10	38
2	E	3/3 (100%)	3 (100%)	0	100	100
2	F	3/3 (100%)	3 (100%)	0	100	100
2	G	3/3 (100%)	3 (100%)	0	100	100
2	H	3/3 (100%)	3 (100%)	0	100	100
All	All	388/388 (100%)	359 (92%)	29 (8%)	13	43

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	251	PRO
1	D	348	LEU
1	C	279	VAL
1	D	269	ILE
1	C	269	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	319	GLN
1	C	320	HIS
1	D	320	HIS
1	C	341	ASN
1	B	283	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.