



Full wwPDB X-ray Structure Validation Report i

Aug 10, 2020 – 07:12 AM BST

PDB ID : 3OV6
Title : CD1c in complex with MPM (mannosyl-beta1-phosphomycoketide)
Authors : Scharf, L.; Li, N.S.; Hawk, A.J.; Garzon, D.; Zhang, T.; Kazen, A.R.; Shah, S.; Haddadian, E.J.; Saghatelian, A.; Faraldo-Gomez, J.D.; Meredith, S.C.; Piccirilli, J.A.; Adams, E.J.
Deposited on : 2010-09-15
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	1	-	-	-	X
3	D12	A	1104	-	-	X	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MK0	A	1105	-	-	-	X

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 3126 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-2-microglobulin, T-cell surface glycoprotein CD1c, T-cell surface glycoprotein CD1b.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	376	3014	1926	517	559	12	178	0	0

There are 26 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	998	ALA	-	expression tag	UNP P61769
A	999	ASP	-	expression tag	UNP P61769
A	1000	PRO	-	expression tag	UNP P61769
A	1100	GLY	-	linker	UNP P61769
A	1101	GLY	-	linker	UNP P61769
A	1102	GLY	-	linker	UNP P61769
A	1103	GLY	-	linker	UNP P61769
A	1104	SER	-	linker	UNP P61769
A	1105	GLY	-	linker	UNP P61769
A	1106	GLY	-	linker	UNP P61769
A	1107	SER	-	linker	UNP P61769
A	1108	GLY	-	linker	UNP P61769
A	1109	SER	-	linker	UNP P61769
A	1110	GLY	-	linker	UNP P61769
A	1111	GLY	-	linker	UNP P61769
A	1112	GLY	-	linker	UNP P61769
A	1113	SER	-	linker	UNP P61769
A	1114	SER	-	linker	UNP P61769
A	52	GLN	ASN	engineered mutation	UNP P29017
A	57	GLN	ASN	engineered mutation	UNP P29017
A	108	GLY	LYS	engineered mutation	UNP P29017
A	128	GLN	ASN	engineered mutation	UNP P29017
A	241	GLN	ASN	engineered mutation	UNP P29016
A	242	GLY	TRP	engineered mutation	UNP P29016
A	279	HIS	-	expression tag	UNP P29016
A	280	HIS	-	expression tag	UNP P29016

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C N O 14 8 1 5	0	0

- Molecule 3 is DODECANE (three-letter code: D12) (formula: C₁₂H₂₆).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C 12 12	0	0

- Molecule 4 is 1-O-[(S)-hydroxy{[(4S,8S,16S,20S)-4,8,12,16,20-pentamethylheptacosyl]oxy}p phosphoryl]-beta-D-mannopyranose (three-letter code: MK0) (formula: C₃₈H₇₇O₉P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	P		
4	A	1	48	38	9	1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O			
5	A	38	38	38		1	0

3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-2-microglobulin, T-cell surface glycoprotein CD1c, T-cell surface glycoprotein CD1b



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	53.72Å 87.09Å 88.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.54 – 2.50 43.54 – 2.50	Depositor EDS
% Data completeness (in resolution range)	(Not available) (43.54-2.50) 91.3 (43.54-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	2.47 (at 2.51Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.5_2)	Depositor
R , R_{free}	0.240 , 0.273 0.231 , 0.268	Depositor DCC
R_{free} test set	686 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	50.4	Xtriage
Anisotropy	0.292	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 49.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.017 for -h,l,k	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3126	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, D12, MK0

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.54	0/3104	0.72	0/4211

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3014	0	2858	84	0
2	A	14	0	13	0	0
3	A	12	0	26	10	0
4	A	48	0	76	10	0
5	A	38	0	0	2	0
All	All	3126	0	2973	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:148:LEU:HD21	3:A:1104:D12:H62	1.52	0.88
1:A:69:LEU:HG	4:A:1105:MK0:HACB	1.61	0.83
1:A:73:TYR:HB2	4:A:1105:MK0:HACA	1.60	0.83
1:A:28:GLY:HA3	4:A:1105:MK0:HAOA	1.67	0.75
3:A:1104:D12:H122	4:A:1105:MK0:O6	1.87	0.75
1:A:1096:ASP:H	1:A:1101:GLY:HA2	1.53	0.74
1:A:1033:SER:HB2	1:A:1054:LEU:HD21	1.71	0.72
1:A:218:TRP:HB3	1:A:265:LYS:HB2	1.72	0.72
1:A:73:TYR:CD1	4:A:1105:MK0:HBB	2.28	0.69
1:A:128:GLN:NE2	5:A:1133:HOH:O	2.22	0.68
1:A:165:SER:O	1:A:168:PRO:HD2	1.94	0.68
1:A:215:LYS:N	1:A:216:PRO:HD2	2.10	0.67
1:A:153:GLU:HA	1:A:156:THR:HG23	1.75	0.66
1:A:77:LEU:HD21	3:A:1104:D12:H91	1.78	0.66
1:A:62:GLU:O	1:A:66:LEU:HD22	1.97	0.64
1:A:220:MET:HG2	1:A:222:MET:HE2	1.80	0.63
1:A:206:VAL:HG22	1:A:249:THR:HG22	1.81	0.63
1:A:76:GLY:HA3	3:A:1104:D12:H121	1.81	0.62
1:A:1059:ASP:O	1:A:1060:TRP:HB2	1.98	0.62
1:A:175:LEU:O	1:A:179:LYS:HG2	1.98	0.62
1:A:1060:TRP:CE2	1:A:117:ALA:HB2	2.35	0.61
1:A:129:THR:HG22	1:A:129:THR:O	1.99	0.61
1:A:1060:TRP:CD2	1:A:117:ALA:HB2	2.36	0.60
1:A:209:VAL:HG21	1:A:219:VAL:HG21	1.83	0.60
1:A:188:PRO:HB3	1:A:212:PHE:HB3	1.86	0.58
1:A:204:GLN:HE21	1:A:249:THR:HB	1.69	0.57
1:A:1095:TRP:CH2	1:A:1097:ARG:HG2	2.39	0.57
1:A:220:MET:CG	1:A:222:MET:HE2	2.36	0.56
1:A:77:LEU:CD2	3:A:1104:D12:H91	2.36	0.56
1:A:182:VAL:HG22	1:A:183:HIS:HB3	1.88	0.55
1:A:92:TYR:N	1:A:93:PRO:HD3	2.21	0.55
1:A:1001:ILE:HG23	1:A:1002:GLN:N	2.23	0.53
1:A:77:LEU:O	1:A:81:ILE:HG12	2.08	0.53
1:A:161:ASN:O	1:A:165:SER:HB3	2.09	0.52
1:A:209:VAL:HG21	1:A:219:VAL:CG2	2.40	0.52
1:A:92:TYR:H	1:A:93:PRO:HD3	1.76	0.51
1:A:77:LEU:HD22	3:A:1104:D12:H123	1.92	0.51
1:A:182:VAL:HG13	1:A:183:HIS:CG	2.45	0.50
1:A:215:LYS:N	1:A:216:PRO:CD	2.75	0.50
1:A:218:TRP:HH2	1:A:222:MET:CE	2.24	0.49
1:A:76:GLY:HA3	3:A:1104:D12:C12	2.42	0.49
1:A:144:VAL:O	1:A:148:LEU:HD13	2.13	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:205:LEU:HB3	1:A:221:TRP:CZ2	2.47	0.49
1:A:8:VAL:HG21	1:A:175:LEU:HD21	1.94	0.49
1:A:218:TRP:HH2	1:A:222:MET:HE3	1.78	0.48
1:A:13:ILE:O	1:A:28:GLY:HA2	2.13	0.48
1:A:83:ASP:O	1:A:84:HIS:HB3	2.14	0.48
1:A:1027:VAL:O	1:A:1063:TYR:HA	2.14	0.48
1:A:1015:ALA:HB1	1:A:1097:ARG:HD3	1.95	0.47
1:A:130:THR:O	1:A:132:VAL:HG23	2.15	0.47
1:A:154:GLY:HA2	1:A:157:GLU:HB2	1.97	0.47
1:A:190:ALA:HA	1:A:208:HIS:O	2.14	0.47
1:A:218:TRP:CH2	1:A:222:MET:HE3	2.50	0.47
1:A:1099:MET:HA	1:A:195:GLY:HA2	1.97	0.46
1:A:73:TYR:CE1	3:A:1104:D12:H91	2.51	0.46
1:A:1055:SER:HB3	1:A:1063:TYR:CZ	2.51	0.46
1:A:160:TYR:O	1:A:164:ARG:HG2	2.16	0.45
1:A:1000:PRO:HA	1:A:1001:ILE:HA	1.76	0.45
1:A:77:LEU:CD2	3:A:1104:D12:H123	2.47	0.45
1:A:140:LEU:O	1:A:144:VAL:HG12	2.17	0.45
1:A:114:PHE:HE1	4:A:1105:MK0:HBDA	1.81	0.44
1:A:162:LEU:HG	4:A:1105:MK0:HBH	1.99	0.44
1:A:1012:ARG:HH12	1:A:241:GLN:HG2	1.82	0.44
1:A:220:MET:HG3	1:A:221:TRP:N	2.33	0.44
1:A:77:LEU:HD11	3:A:1104:D12:H32	2.00	0.44
1:A:174:LEU:HD12	1:A:174:LEU:HA	1.83	0.43
1:A:126:PHE:CZ	1:A:129:THR:HA	2.53	0.43
1:A:266:HIS:CD2	5:A:1130:HOH:O	2.72	0.43
1:A:1059:ASP:O	1:A:1060:TRP:CB	2.66	0.42
1:A:162:LEU:HD12	1:A:162:LEU:HA	1.87	0.42
1:A:197:SER:HA	1:A:198:PRO:HD3	1.88	0.42
1:A:1004:THR:CG2	1:A:1004:THR:O	2.66	0.42
1:A:145:CYS:O	1:A:149:ASN:HB2	2.19	0.42
1:A:167:CYS:HB3	1:A:168:PRO:HD3	2.01	0.42
1:A:211:GLY:HA2	1:A:243:THR:OG1	2.20	0.42
1:A:137:CYS:HA	1:A:138:GLY:HA3	1.53	0.42
1:A:40:TRP:HB2	4:A:1105:MK0:HAQA	2.01	0.42
1:A:1036:GLU:O	1:A:1082:VAL:HA	2.20	0.41
1:A:264:VAL:HB	1:A:274:ILE:HB	2.02	0.41
1:A:8:VAL:HG12	1:A:9:SER:N	2.35	0.41
1:A:83:ASP:O	1:A:84:HIS:CB	2.68	0.41
1:A:171:LEU:O	1:A:171:LEU:HD12	2.20	0.41
4:A:1105:MK0:HBE	4:A:1105:MK0:HPB	1.94	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:153:GLU:HA	1:A:156:THR:CG2	2.46	0.40
4:A:1105:MK0:HARA	4:A:1105:MK0:HACA	1.90	0.40
1:A:114:PHE:CD2	1:A:114:PHE:C	2.95	0.40
1:A:40:TRP:CZ2	1:A:45:GLY:HA2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	370/397 (93%)	349 (94%)	19 (5%)	2 (0%)	29 48

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1102	GLY
1	A	92	TYR

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	327/339 (96%)	304 (93%)	23 (7%)	15 29

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1004	THR
1	A	1019	LYS
1	A	1031	HIS
1	A	1070	PHE
1	A	1088	SER
1	A	1093	VAL
1	A	23	TRP
1	A	66	LEU
1	A	68	LEU
1	A	137	CYS
1	A	139	SER
1	A	146	HIS
1	A	156	THR
1	A	162	LEU
1	A	174	LEU
1	A	175	LEU
1	A	176	ASP
1	A	183	HIS
1	A	194	SER
1	A	205	LEU
1	A	233	LEU
1	A	251	ASP
1	A	254	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1031	HIS
1	A	105	HIS
1	A	115	GLN
1	A	161	ASN
1	A	204	GLN
1	A	232	GLN
1	A	241	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	MK0	A	1105	-	47,48,48	0.56	1 (2%)	60,61,61	1.54	11 (18%)
2	NAG	A	1	1	14,14,15	0.70	0	17,19,21	2.07	3 (17%)
3	D12	A	1104	-	11,11,11	0.30	0	10,10,10	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MK0	A	1105	-	-	19/46/66/66	0/1/1/1
2	NAG	A	1	1	-	1/6/23/26	0/1/1/1
3	D12	A	1104	-	-	2/9/9/9	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1105	MK0	PBV-O1	2.27	1.66	1.60

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	NAG	C2-N2-C7	5.97	131.40	122.90
4	A	1105	MK0	C4-C3-C2	4.42	118.54	110.82
4	A	1105	MK0	C3-C4-C5	4.09	117.54	110.24
2	A	1	NAG	O5-C1-C2	-3.91	105.11	111.29
4	A	1105	MK0	CAV-CBH-CBP	-3.39	104.96	115.92
4	A	1105	MK0	CAS-CBA-CBL	3.34	126.73	115.92
4	A	1105	MK0	PBV-OBI-CAX	-2.85	107.54	121.59
4	A	1105	MK0	C1-C2-C3	2.75	115.73	110.00
4	A	1105	MK0	OBI-CAX-CAR	2.71	118.04	108.99
4	A	1105	MK0	O5-C5-C4	2.67	114.55	109.69
4	A	1105	MK0	C1-O5-C5	2.55	118.68	113.69
4	A	1105	MK0	CAT-CBD-CBO	-2.43	108.06	115.92
2	A	1	NAG	O7-C7-N2	2.29	126.17	121.95
4	A	1105	MK0	O1-C1-C2	2.09	112.21	108.38

There are no chirality outliers.

All (22) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	1105	MK0	CAX-OBI-PBV-OAG
2	A	1	NAG	C3-C2-N2-C7
4	A	1105	MK0	CAQ-CAY-CBL-CAB
4	A	1105	MK0	CAX-OBI-PBV-O1
4	A	1105	MK0	CAX-CAR-CAZ-CBM
4	A	1105	MK0	CBD-CAT-CBB-CBM
4	A	1105	MK0	CAS-CBA-CBL-CAB
4	A	1105	MK0	CAN-CAO-CAP-CAQ
3	A	1104	D12	C3-C4-C5-C6
4	A	1105	MK0	CAS-CBA-CBL-CAY
4	A	1105	MK0	CAS-CBC-CBN-CBE
4	A	1105	MK0	CAS-CBC-CBN-CAD
4	A	1105	MK0	CBH-CAV-CBF-CBO
4	A	1105	MK0	CAZ-CAR-CAX-OBI
4	A	1105	MK0	O5-C5-C6-O6
4	A	1105	MK0	CBC-CAS-CBA-CBL
3	A	1104	D12	C9-C10-C11-C12
4	A	1105	MK0	CAV-CBH-CBP-CBG
4	A	1105	MK0	CAV-CBH-CBP-CAF
4	A	1105	MK0	CAV-CBF-CBO-CBD
4	A	1105	MK0	CAO-CAP-CAQ-CAY
4	A	1105	MK0	CAX-OBI-PBV-OAL

There are no ring outliers.

2 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1105	MK0	10	0
3	A	1104	D12	10	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	376/397 (94%)	0.37	28 (7%) 14 15	33, 53, 87, 114	46 (12%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	137	CYS	15.0
1	A	90	SER	9.4
1	A	182	VAL	7.4
1	A	136	GLY	6.6
1	A	183	HIS	6.2
1	A	146	HIS	6.0
1	A	91	LYS	5.4
1	A	152	TYR	5.0
1	A	148	LEU	4.9
1	A	151	GLN	3.5
1	A	144	VAL	3.5
1	A	180	MET	3.5
1	A	181	TYR	3.1
1	A	138	GLY	3.0
1	A	141	ALA	2.9
1	A	23	TRP	2.7
1	A	215	LYS	2.7
1	A	150	HIS	2.7
1	A	1000	PRO	2.5
1	A	154	GLY	2.4
1	A	18	PHE	2.3
1	A	85	ALA	2.3
1	A	185	GLN	2.2
1	A	86	SER	2.2
1	A	22	SER	2.2
1	A	60	ASN	2.1
1	A	139	SER	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1003	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	A	1	14/15	0.35	0.49	101,111,114,115	0
4	MK0	A	1105	48/48	0.66	0.42	42,67,114,120	0
3	D12	A	1104	12/12	0.70	0.37	66,72,80,84	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.