wwPDB X-ray Structure Validation Summary Report (i)

May 22, 2020 – 03:43 pm BST

PDB ID	:	1PDG
Title	:	CRYSTAL STRUCTURE OF HUMAN PLATELET-DERIVED GROWTH FACTOR BB
Authors	:	Oefner, C.; Darcy, A.D.; Winkler, F.K.; Eggimann, B.; Hosnag, M.
Deposited on	:	1992-07-14
Resolution	:	3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	109	14% 44% 17% 5% 20%
1	B	109	16% 29% 23% 8% 24%
1	C	109	17% 36% 19% 7% 21%

2 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 2035 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PLATELET-DERIVED GROWTH FACTOR BB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	87	692	433	129	121	9	0	0	0
			Total	C	N	O	S			
1	B	83	659	412	122	116	9	0	0	0
			Total	C	N	O	S			
1	C	86	684	430	125	120	9	0	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: PLATELET-DERIVED GROWTH FACTOR BB

Chain A:





- Molecule 1: PLATELET-DERIVED GROWTH FACTOR BB

Chain B:



- Molecule 1: PLATELET-DERIVED GROWTH FACTOR BB

Chain C:





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	147.30Å 31.80Å 90.10Å 90.00° 98.70° 90.00°	Depositor
Resolution (Å)	6.00 – 3.00	Depositor
% Data completeness (in resolution range)	(Not available) (6.00-3.00)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROLSQ	Depositor
R, R_{free}	0.209 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2035	wwPDB-VP
Average B, all atoms ($Å^2$)	25.0	wwPDB-VP

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	1.24	3/702 (0.4%)	2.07	17/947 (1.8%)
1	B	1.11	1/668 (0.1%)	1.92	20/901 (2.2%)
1	C	1.10	0/694	1.95	23/937 (2.5%)
All	All	1.15	4/2064 (0.2%)	1.98	60/2785 (2.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
All	All	0	3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	97	CYS	CB-SG	-7.88	1.68	1.82
1	B	43	CYS	CB-SG	7.01	1.94	1.82
1	A	49	CYS	CB-SG	-6.31	1.71	1.82
1	A	52	CYS	CB-SG	-5.07	1.73	1.81

The worst 5 of 60 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	C	73	ARG	CA-CB-CG	15.85	148.26	113.40
1	A	19	ARG	NE-CZ-NH1	-13.68	113.46	120.30
1	C	45	GLU	CA-CB-CG	12.25	140.34	113.40
1	A	45	GLU	CA-CB-CG	11.07	137.75	113.40
1	C	19	ARG	NE-CZ-NH2	-11.04	114.78	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	48	ARG	Sidechain
1	B	19	ARG	Sidechain
1	B	68	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	692	0	713	107	0
1	B	659	0	681	100	0
1	C	684	0	708	103	2
All	All	2035	0	2102	297	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 72.

The worst 5 of 297 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:78:VAL:HG12	1:C:81:LYS:NZ	1.52	1.21
1:C:68:ARG:NH1	1:C:91:LEU:HD12	1.62	1.15
1:A:19:ARG:HG2	1:A:19:ARG:HH11	1.09	1.13
1:C:68:ARG:HH12	1:C:91:LEU:CD1	1.66	1.07
1:A:78:VAL:HB	1:A:83:ILE:HD11	1.33	1.07

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:43:CYS:SG	1:C:52:CYS:SG[2_555]	1.72	0.48
1:C:19:ARG:NH2	1:C:45:GLU:OE1[2_555]	2.06	0.14

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	83/109 (76%)	68 (82%)	14 (17%)	1 (1%)	13 48
1	B	79/109 (72%)	61 (77%)	13 (16%)	5 (6%)	1 7
1	C	82/109 (75%)	59 (72%)	16 (20%)	7 (8%)	1 4
All	All	244/327 (75%)	188 (77%)	43 (18%)	13 (5%)	2 11

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	42	PRO
1	B	54	ASN
1	B	79	ARG
1	C	15	GLU
1	C	56	ARG

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	80/98 (82%)	54 (68%)	26 (32%)	0 1
1	B	77/98 (79%)	44 (57%)	33 (43%)	0 0
1	C	80/98 (82%)	59 (74%)	21 (26%)	0 2
All	All	237/294 (81%)	157 (66%)	80 (34%)	0 1

5 of 80 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	42	PRO
1	B	61	ARG
1	C	83	ILE
1	B	43	CYS
1	B	55	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	66	GLN
1	C	55	ASN
1	C	64	GLN
1	B	59	GLN
1	C	59	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.