



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 13, 2023 – 08:26 AM EDT

PDB ID : 4PVF
Title : Crystal structure of Homo sapiens holo serine hydroxymethyltransferase 2 (mitochondrial) (SHMT2), isoform 3, transcript variant 5, 483 aa, at 2.6 ang. resolution
Authors : Giardina, G.; Brunotti, P.; Fiascarelli, A.; Contestabile, R.; Cutruzzola, F.
Deposited on : 2014-03-17
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	483	 83% 8% • 6%
1	B	483	 82% 8% • 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	501	-	-	X	-
2	GOL	B	501	-	-	X	-
3	PEG	A	502	-	-	X	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 7229 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine hydroxymethyltransferase, mitochondrial.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	P				S
1	A	454	Total 3561	C 2244	N 637	O 663	P 1	S 16	0	0	0
1	B	447	Total 3523	C 2222	N 629	O 655	P 1	S 16	0	2	0

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 6	C 3	O 3	0	0
2	B	1	Total 6	C 3	O 3	0	0

- Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 7 4 3	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	78	Total O 78 78	0	0
4	B	48	Total O 48 48	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Serine hydroxymethyltransferase, mitochondrial

Chain A: 



- Molecule 1: Serine hydroxymethyltransferase, mitochondrial

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	160.46Å 160.46Å 211.46Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	138.96 – 2.60 49.48 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (138.96-2.60) 100.0 (49.48-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.90 (at 2.61Å)	Xtrriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.206 , 0.243 0.213 , 0.250	Depositor DCC
R_{free} test set	2531 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	30.7	Xtrriage
Anisotropy	0.331	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 14.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7229	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LLP, PEG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.74	0/3607	0.91	6/4876 (0.1%)
1	B	0.73	0/3573	0.90	9/4828 (0.2%)
All	All	0.74	0/7180	0.91	15/9704 (0.2%)

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	425	ASP	CB-CG-OD1	7.51	125.06	118.30
1	A	48	ARG	NE-CZ-NH2	-7.12	116.74	120.30
1	A	387	ASN	CB-CA-C	-7.12	96.17	110.40
1	B	387	ASN	CB-CA-C	-6.98	96.44	110.40
1	A	161	ARG	NE-CZ-NH1	6.86	123.73	120.30
1	B	265	ARG	CG-CD-NE	-6.39	98.37	111.80
1	B	425	ASP	CB-CG-OD2	-6.09	112.81	118.30
1	A	48	ARG	NE-CZ-NH1	5.96	123.28	120.30
1	A	428	ASP	CB-CG-OD1	5.81	123.53	118.30
1	B	48	ARG	NE-CZ-NH1	5.61	123.10	120.30
1	A	199	LEU	CA-CB-CG	5.34	127.58	115.30
1	B	452	LEU	CA-CB-CG	5.22	127.30	115.30
1	B	293	ARG	NE-CZ-NH2	-5.12	117.74	120.30
1	B	381	LEU	CB-CG-CD1	5.11	119.69	111.00
1	B	199	LEU	CA-CB-CG	5.05	126.93	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3561	0	3545	39	0
1	B	3523	0	3510	29	0
2	A	6	0	8	5	0
2	B	6	0	8	7	0
3	A	7	0	10	8	0
4	A	78	0	0	2	0
4	B	48	0	0	1	0
All	All	7229	0	7081	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:191:THR:HG21	4:B:605:HOH:O	1.68	0.91
1:A:85:TYR:HH	2:B:501:GOL:HO1	1.15	0.85
1:A:55:SER:OG	2:A:501:GOL:H2	1.84	0.77
1:A:127:LEU:O	1:A:131:THR:HG23	1.88	0.74
1:A:388:LYS:HD2	1:A:400:PRO:HG2	1.70	0.74
1:B:127:LEU:O	1:B:131:THR:HG23	1.88	0.74
1:B:259:LLP:H4'1	2:B:501:GOL:H2	1.72	0.71
1:A:210:LEU:HA	3:A:502:PEG:C4	2.20	0.71
1:A:259:LLP:H4'1	2:A:501:GOL:H12	1.75	0.69
1:A:234:ILE:CD1	3:A:502:PEG:H32	2.25	0.66
1:A:55:SER:OG	2:A:501:GOL:C2	2.43	0.65
1:A:55:SER:CB	2:A:501:GOL:H2	2.27	0.65
1:A:210:LEU:HA	3:A:502:PEG:H42	1.81	0.63
1:A:191:THR:HG21	4:A:654:HOH:O	1.99	0.63
1:A:210:LEU:HA	3:A:502:PEG:H41	1.80	0.62
1:B:449:SER:O	1:B:453:LYS:CG	2.50	0.59
1:A:85:TYR:OH	2:B:501:GOL:O1	2.07	0.59
1:B:55:SER:OG	2:B:501:GOL:O3	2.21	0.57
1:B:262:ARG:HH22	1:B:325:MET:HE1	1.70	0.56
1:B:422:ARG:HG3	1:B:422:ARG:HH11	1.71	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:449:SER:O	1:B:453:LYS:HG2	2.06	0.55
1:A:244:PRO:HD2	3:A:502:PEG:H12	1.89	0.53
2:B:501:GOL:O1	2:B:501:GOL:O3	2.25	0.53
1:A:55:SER:HB3	2:A:501:GOL:H2	1.92	0.52
1:B:141:MET:HE1	1:B:191:THR:HG23	1.91	0.52
1:A:234:ILE:HD12	3:A:502:PEG:H32	1.93	0.51
1:A:338:ARG:HG3	1:A:338:ARG:HH11	1.75	0.51
1:B:141:MET:HE1	1:B:191:THR:CG2	2.41	0.51
1:B:338:ARG:HH11	1:B:338:ARG:HG3	1.77	0.50
1:A:422:ARG:HH11	1:A:422:ARG:HG3	1.76	0.49
1:B:188:LEU:CD1	1:B:216:MET:CE	2.91	0.49
1:A:370:LEU:HG	1:A:371:ASP:H	1.78	0.49
1:A:188:LEU:CD1	1:A:216:MET:CE	2.91	0.49
1:A:234:ILE:HD13	3:A:502:PEG:H32	1.95	0.49
1:A:33:PRO:HD2	4:A:672:HOH:O	2.12	0.48
1:B:188:LEU:HD11	1:B:216:MET:CE	2.42	0.48
1:A:141:MET:HE1	1:A:191:THR:HG23	1.95	0.48
1:B:370:LEU:HG	1:B:371:ASP:H	1.78	0.48
1:A:445:GLN:NE2	1:A:445:GLN:HA	2.27	0.48
1:A:329:TYR:O	1:A:333:VAL:HG23	2.14	0.47
1:A:187:GLN:O	1:A:191:THR:HB	2.14	0.47
1:B:187:GLN:O	1:B:191:THR:HB	2.15	0.46
1:A:396:SER:O	1:A:397:ALA:CB	2.63	0.46
1:B:329:TYR:O	1:B:333:VAL:HG23	2.15	0.46
1:A:217:ARG:HA	1:A:217:ARG:HD3	1.74	0.46
1:B:407:ALA:N	1:B:408:PRO:HD3	2.31	0.46
1:A:188:LEU:HD11	1:A:216:MET:CE	2.46	0.45
1:A:422:ARG:HG3	1:A:422:ARG:NH1	2.32	0.45
1:B:366:ARG:HB2	1:B:367:PRO:HD3	1.98	0.44
1:B:407:ALA:N	1:B:408:PRO:CD	2.80	0.44
1:A:407:ALA:N	1:A:408:PRO:CD	2.81	0.43
1:A:407:ALA:N	1:A:408:PRO:HD3	2.33	0.43
1:A:244:PRO:CD	3:A:502:PEG:H12	2.49	0.42
1:A:141:MET:HE1	1:A:191:THR:CG2	2.48	0.42
1:B:422:ARG:NH1	1:B:422:ARG:CG	2.82	0.42
1:B:449:SER:O	1:B:453:LYS:HG3	2.19	0.42
1:B:55:SER:HG	2:B:501:GOL:HO3	1.58	0.42
1:B:118:GLN:N	1:B:119:PRO:CD	2.83	0.42
1:A:321:ALA:HA	1:A:326:PHE:CG	2.55	0.42
1:A:118:GLN:N	1:A:119:PRO:CD	2.83	0.41
1:B:259:LLP:C4'	2:B:501:GOL:H2	2.45	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:262:ARG:HH22	1:B:325:MET:CE	2.34	0.41
1:A:188:LEU:HD12	1:A:216:MET:CE	2.50	0.41
1:A:366:ARG:HB2	1:A:367:PRO:HD3	2.01	0.41
1:B:416:ARG:HG3	1:B:416:ARG:NH1	2.35	0.41
1:A:256:THR:HG21	1:A:259:LLP:HE2	2.03	0.40
1:B:407:ALA:H	1:B:408:PRO:HD3	1.86	0.40
1:B:156:MET:HB3	1:B:161:ARG:HA	2.04	0.40
1:B:347:ARG:NH2	1:B:425:ASP:HA	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	449/483 (93%)	433 (96%)	15 (3%)	1 (0%)	47 71
1	B	442/483 (92%)	424 (96%)	18 (4%)	0	100 100
All	All	891/966 (92%)	857 (96%)	33 (4%)	1 (0%)	51 75

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	397	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	373/394 (95%)	348 (93%)	25 (7%)	16	33
1	B	370/394 (94%)	342 (92%)	28 (8%)	13	26
All	All	743/788 (94%)	690 (93%)	53 (7%)	14	29

All (53) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	48	ARG
1	A	75	TYR
1	A	121	SER
1	A	127	LEU
1	A	131	THR
1	A	143	LEU
1	A	156	MET
1	A	159	VAL
1	A	176	LEU
1	A	188	LEU
1	A	191	THR
1	A	196	ARG
1	A	216	MET
1	A	217	ARG
1	A	272	ARG
1	A	331	LEU
1	A	350	SER
1	A	387	ASN
1	A	396	SER
1	A	416	ARG
1	A	422	ARG
1	A	445	GLN
1	A	449	SER
1	A	458	SER
1	A	482	GLU
1	B	37	GLU
1	B	75	TYR
1	B	121	SER
1	B	127	LEU
1	B	131	THR
1	B	143	LEU
1	B	156	MET
1	B	159	VAL
1	B	176	LEU
1	B	188	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	191	THR
1	B	196	ARG
1	B	216	MET
1	B	217	ARG
1	B	272	ARG
1	B	325	MET
1	B	331	LEU
1	B	350	SER
1	B	381	LEU
1	B	387	ASN
1	B	390	THR
1	B	422	ARG
1	B	441	THR
1	B	445	GLN
1	B	446	ASP
1	B	449	SER
1	B	452	LEU
1	B	482	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	320	GLN
1	A	445	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LLP	B	259	1	23,24,25	1.94	4 (17%)	25,32,34	2.04	6 (24%)
1	LLP	A	259	1	23,24,25	2.16	5 (21%)	25,32,34	1.91	7 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	B	259	1	-	7/16/17/19	0/1/1/1
1	LLP	A	259	1	-	7/16/17/19	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	259	LLP	C4'-NZ	5.98	1.47	1.27
1	B	259	LLP	C4'-NZ	5.57	1.45	1.27
1	B	259	LLP	C3-C2	4.91	1.45	1.40
1	A	259	LLP	C3-C2	4.71	1.45	1.40
1	A	259	LLP	C4-C5	4.40	1.47	1.42
1	A	259	LLP	C4-C3	4.40	1.47	1.40
1	B	259	LLP	C4-C3	3.68	1.46	1.40
1	B	259	LLP	C4-C5	2.55	1.45	1.42
1	A	259	LLP	C4-C4'	2.07	1.50	1.46

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	259	LLP	OP4-C5'-C5	5.59	120.01	109.35
1	B	259	LLP	OP4-C5'-C5	5.52	119.88	109.35
1	B	259	LLP	C4-C4'-NZ	-5.16	100.64	124.31
1	A	259	LLP	C4-C4'-NZ	-3.36	108.90	124.31
1	A	259	LLP	C6-N1-C2	2.81	124.38	119.17
1	B	259	LLP	C5-C4-C4'	2.77	126.12	121.56
1	A	259	LLP	C5-C4-C4'	2.45	125.58	121.56
1	B	259	LLP	CE-NZ-C4'	2.30	125.96	118.90
1	A	259	LLP	C2'-C2-N1	2.24	122.05	117.67
1	A	259	LLP	C3-C2-N1	-2.19	117.94	120.77
1	A	259	LLP	OP2-P-OP1	2.12	119.00	110.68
1	B	259	LLP	C2'-C2-N1	2.01	121.61	117.67
1	B	259	LLP	C3-C4-C4'	-2.01	116.67	120.41

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	259	LLP	C4-C4'-NZ-CE
1	A	259	LLP	C5'-OP4-P-OP2
1	A	259	LLP	O-C-CA-CB
1	B	259	LLP	C5'-OP4-P-OP2
1	B	259	LLP	O-C-CA-CB
1	B	259	LLP	CG-CD-CE-NZ
1	A	259	LLP	C5'-OP4-P-OP1
1	B	259	LLP	C5'-OP4-P-OP1
1	B	259	LLP	C3-C4-C4'-NZ
1	B	259	LLP	C5-C4-C4'-NZ
1	A	259	LLP	CD-CE-NZ-C4'
1	B	259	LLP	CD-CE-NZ-C4'
1	A	259	LLP	C3-C4-C4'-NZ
1	A	259	LLP	CG-CD-CE-NZ

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	259	LLP	2	0
1	A	259	LLP	2	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PEG	A	502	-	6,6,6	1.30	0	5,5,5	1.55	2 (40%)
2	GOL	A	501	-	5,5,5	0.50	0	5,5,5	1.32	0
2	GOL	B	501	-	5,5,5	0.55	0	5,5,5	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEG	A	502	-	-	1/4/4/4	-
2	GOL	A	501	-	-	2/4/4/4	-
2	GOL	B	501	-	-	2/4/4/4	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	PEG	O1-C1-C2	2.12	124.11	111.81
3	A	502	PEG	C3-O2-C2	2.08	122.28	113.29

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	GOL	O1-C1-C2-C3
2	B	501	GOL	O1-C1-C2-C3
2	A	501	GOL	O1-C1-C2-O2
3	A	502	PEG	O1-C1-C2-O2
2	B	501	GOL	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	502	PEG	8	0
2	A	501	GOL	5	0
2	B	501	GOL	7	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	453/483 (93%)	-0.53	1 (0%) 95 95	20, 24, 55, 89	0
1	B	446/483 (92%)	-0.17	16 (3%) 42 35	20, 35, 78, 119	0
All	All	899/966 (93%)	-0.35	17 (1%) 66 62	20, 29, 70, 119	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	450	PHE	5.4
1	B	439	SER	4.5
1	B	442	ALA	3.5
1	B	286	ILE	3.5
1	B	459	GLN	3.2
1	B	277	ALA	3.1
1	A	277	ALA	3.0
1	B	399	THR	2.7
1	B	457	THR	2.7
1	B	452	LEU	2.7
1	B	453	LYS	2.5
1	B	378	VAL	2.4
1	B	275	VAL	2.3
1	B	444	LEU	2.2
1	B	456	GLU	2.2
1	B	179	LYS	2.2
1	B	463	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
1	LLP	B	259	24/25	0.97	0.13	23,29,32,34	0
1	LLP	A	259	24/25	0.98	0.15	20,20,20,22	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	PEG	A	502	7/7	0.83	0.32	20,20,21,22	7
2	GOL	B	501	6/6	0.89	0.20	39,46,49,54	0
2	GOL	A	501	6/6	0.91	0.49	20,20,20,20	6

6.5 Other polymers [i](#)

There are no such residues in this entry.