



# wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 17, 2023 – 02:49 PM EST

PDB ID : 1QHB  
Title : VANADIUM BROMOPEROXIDASE FROM RED ALGA CORALLINA OF-  
FICINALIS  
Authors : Isupov, M.N.; Dalby, A.R.; Brindley, A.A.; Littlechild, J.A.  
Deposited on : 1999-05-11  
Resolution : 2.30 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	596	
1	B	596	
1	C	596	
1	D	596	
1	E	596	
1	F	596	

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 29624 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HALOPEROXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	595	Total 4567	C 2900	N 766	O 893	S 8	0	0	0
1	B	595	Total 4567	C 2900	N 766	O 893	S 8	0	0	0
1	C	595	Total 4567	C 2900	N 766	O 893	S 8	0	0	0
1	D	595	Total 4567	C 2900	N 766	O 893	S 8	0	0	0
1	E	595	Total 4567	C 2900	N 766	O 893	S 8	0	0	0
1	F	595	Total 4567	C 2900	N 766	O 893	S 8	0	0	0

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O P 5 4 1	0	0
2	B	1	Total O P 5 4 1	0	0
2	C	1	Total O P 5 4 1	0	0
2	D	1	Total O P 5 4 1	0	0
2	E	1	Total O P 5 4 1	0	0
2	F	1	Total O P 5 4 1	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Ca 1 1	0	0
3	B	1	Total Ca 1 1	0	0
3	C	1	Total Ca 1 1	0	0
3	D	1	Total Ca 1 1	0	0
3	E	1	Total Ca 1 1	0	0
3	F	1	Total Ca 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	335	Total O 335 335	0	0
4	B	394	Total O 394 394	0	0
4	C	372	Total O 372 372	0	0
4	D	387	Total O 387 387	0	0
4	E	352	Total O 352 352	0	0
4	F	346	Total O 346 346	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HALOPEROXIDASE



- Molecule 1: HALOPEROXIDASE




- Molecule 1: HALOPEROXIDASE





- Molecule 1: HALOPEROXIDASE







• Molecule 1: HALOPEROXIDASE



• Molecule 1: HALOPEROXIDASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 42 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	201.91Å 201.91Å 178.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.30 19.94 – 2.30	Depositor EDS
% Data completeness (in resolution range)	94.8 (20.00-2.30) 96.8 (19.94-2.30)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.42 (at 2.30Å)	Xtrriage
Refinement program	REFMAC WITH INPUT DM PHASES	Depositor
R, $R_{free}$	0.172 , 0.227 0.155 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.6	Xtrriage
Anisotropy	0.044	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 49.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.53$ , $\langle L^2 \rangle = 0.36$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	29624	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 54.01 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.8834e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.64	0/4666	1.33	27/6339 (0.4%)
1	B	0.63	0/4666	1.31	37/6339 (0.6%)
1	C	0.62	0/4666	1.29	26/6339 (0.4%)
1	D	0.64	0/4666	1.30	36/6339 (0.6%)
1	E	0.61	0/4666	1.33	39/6339 (0.6%)
1	F	0.64	0/4666	1.29	30/6339 (0.5%)
All	All	0.63	0/27996	1.31	195/38034 (0.5%)

There are no bond length outliers.

The worst 5 of 195 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	29	ARG	NE-CZ-NH1	15.46	128.03	120.30
1	A	29	ARG	NE-CZ-NH2	-15.14	112.73	120.30
1	A	321	ARG	NE-CZ-NH1	13.44	127.02	120.30
1	D	503	ASN	OD1-CG-ND2	-12.95	92.12	121.90
1	C	389	ARG	NE-CZ-NH2	-12.58	114.01	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4567	0	4453	18	0
1	B	4567	0	4453	18	0
1	C	4567	0	4453	24	0
1	D	4567	0	4453	15	0
1	E	4567	0	4453	22	0
1	F	4567	0	4453	23	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
2	E	5	0	0	0	0
2	F	5	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
4	A	335	0	0	3	0
4	B	394	0	0	3	0
4	C	372	0	0	3	0
4	D	387	0	0	2	0
4	E	352	0	0	5	0
4	F	346	0	0	1	0
All	All	29624	0	26718	112	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 112 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:203:LYS:HG2	4:E:3321:HOH:O	1.77	0.84
1:A:49:GLY:HA2	4:A:3281:HOH:O	1.83	0.78
1:F:502:ALA:HB1	1:F:531:ALA:HB2	1.72	0.72
1:E:124:GLU:O	1:E:395:ARG:NH1	2.28	0.67
1:C:369:ASP:HB3	1:F:124:GLU:HG2	1.77	0.66

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	593/596 (100%)	585 (99%)	8 (1%)	0	100	100
1	B	593/596 (100%)	583 (98%)	8 (1%)	2 (0%)	41	50
1	C	593/596 (100%)	587 (99%)	6 (1%)	0	100	100
1	D	593/596 (100%)	585 (99%)	8 (1%)	0	100	100
1	E	593/596 (100%)	586 (99%)	7 (1%)	0	100	100
1	F	593/596 (100%)	583 (98%)	9 (2%)	1 (0%)	47	58
All	All	3558/3576 (100%)	3509 (99%)	46 (1%)	3 (0%)	51	64

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	424	SER
1	B	371	PHE
1	F	371	PHE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	487/488 (100%)	477 (98%)	10 (2%)	53	70
1	B	487/488 (100%)	478 (98%)	9 (2%)	59	75
1	C	487/488 (100%)	476 (98%)	11 (2%)	50	67
1	D	487/488 (100%)	476 (98%)	11 (2%)	50	67

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	487/488 (100%)	475 (98%)	12 (2%)	47	65
1	F	487/488 (100%)	476 (98%)	11 (2%)	50	67
All	All	2922/2928 (100%)	2858 (98%)	64 (2%)	52	69

5 of 64 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	187	SER
1	F	394	VAL
1	C	423	LYS
1	C	394	VAL
1	F	440	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	368	GLN
1	E	66	ASN
1	F	341	ASN
1	D	595	GLN
1	E	341	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 6 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PO4	A	3000	-	4,4,4	1.55	1 (25%)	6,6,6	0.74	0
2	PO4	B	3002	-	4,4,4	1.60	0	6,6,6	1.57	2 (33%)
2	PO4	F	3010	-	4,4,4	1.46	1 (25%)	6,6,6	0.95	0
2	PO4	E	3008	-	4,4,4	1.39	0	6,6,6	0.26	0
2	PO4	C	3004	-	4,4,4	1.62	1 (25%)	6,6,6	1.19	0
2	PO4	D	3006	-	4,4,4	1.59	1 (25%)	6,6,6	0.57	0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	3004	PO4	P-O2	-2.43	1.47	1.54
2	A	3000	PO4	P-O3	-2.29	1.47	1.54
2	D	3006	PO4	P-O3	-2.24	1.47	1.54
2	F	3010	PO4	P-O2	-2.09	1.48	1.54

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	3002	PO4	O3-P-O2	2.51	116.03	107.97
2	B	3002	PO4	O3-P-O1	-2.28	102.54	110.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	595/596 (99%)	-0.63	10 (1%) 70 76	7, 15, 40, 74	0
1	B	595/596 (99%)	-0.72	5 (0%) 86 89	7, 15, 39, 75	0
1	C	595/596 (99%)	-0.73	4 (0%) 87 91	7, 15, 38, 66	0
1	D	595/596 (99%)	-0.72	2 (0%) 94 96	7, 15, 39, 73	0
1	E	595/596 (99%)	-0.70	6 (1%) 82 86	7, 15, 38, 66	0
1	F	595/596 (99%)	-0.70	6 (1%) 82 86	7, 15, 39, 90	0
All	All	3570/3576 (99%)	-0.70	33 (0%) 84 88	7, 15, 39, 90	0

The worst 5 of 33 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	365	LEU	4.8
1	F	102	GLU	4.5
1	B	102	GLU	3.4
1	D	102	GLU	3.3
1	C	459	VAL	3.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CA	C	3005	1/1	0.83	0.10	45,45,45,45	0
3	CA	F	3011	1/1	0.86	0.09	45,45,45,45	0
3	CA	D	3007	1/1	0.90	0.09	45,45,45,45	0
3	CA	B	3003	1/1	0.91	0.17	44,44,44,44	0
3	CA	A	3001	1/1	0.92	0.11	46,46,46,46	0
3	CA	E	3009	1/1	0.95	0.08	45,45,45,45	0
2	PO4	A	3000	5/5	0.99	0.05	11,14,15,17	0
2	PO4	B	3002	5/5	0.99	0.07	11,13,16,17	0
2	PO4	C	3004	5/5	0.99	0.06	12,14,15,17	0
2	PO4	D	3006	5/5	0.99	0.06	11,13,16,17	0
2	PO4	E	3008	5/5	0.99	0.05	10,13,14,18	0
2	PO4	F	3010	5/5	0.99	0.05	13,13,15,17	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.