wwPDB NMR Structure Validation Summary Report (i)

May 28, 2020 – 08:24 pm BST

PDB ID	:	1QJT
Title	:	SOLUTION STRUCTURE OF THE APO EH1 DOMAIN OF MOUSE EPIDERMAL GROWTH FACTOR RECEPTOR SUBSTRATE 15, EPS15
Authors	:	Whitehead, B.; Tessari, M.; Carotenuto, A.; van Bergen en Henegouwen, P.M.; Vuister, G.W.
Deposited on	:	1999-07-02

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	2.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 92%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	99	28% 56% 14% .

2 Ensemble composition and analysis (i)

This entry contains 30 models. Model 2 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:9-A:105 (97)	0.38	2

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 7 clusters and 5 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 8, 11, 25, 26, 28
2	6, 12, 15, 21, 22, 23
3	5, 13
4	7, 14
5	20, 27
6	19, 29
7	10, 17
Single-model clusters	9; 16; 18; 24; 30

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1509 atoms, of which 757 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called EPIDERMAL GROWTH FACTOR RECEPTOR SUBSTRATE SUBSTRATE 15, EPS15.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	99	1509	483	757	127	141	1	0

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: EPIDERMAL GROWTH FACTOR RECEPTOR SUBSTRATE SUBSTRATE 15, EPS15

Chain A:





4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 2. Colouring as in section 4.1 above.

• Molecule 1: EPIDERMAL GROWTH FACTOR RECEPTOR SUBSTRATE SUBSTRATE 15, EPS15

Chain A:



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *TAD*.

Of the 100 calculated structures, 30 were deposited, based on the following criterion: *LOWEST OVERALL TARGET FUNCTION*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DYANA	refinement	1.5
NMRPipe	structure solution	
XEASY	structure solution	
DYANA	structure solution	1.5

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	input_cs.cif
Number of chemical shift lists	1
Total number of shifts	1399
Number of shifts mapped to atoms	1206
Number of unparsed shifts	0
Number of shifts with mapping errors	193
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	92%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	738	741	753	70±5
All	All	22140	22230	22590	2111

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 47.

5 of 380 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:9:LEU:HD13	1:A:74:PHE:CE1	0.97	1.95	20	6
1:A:48:LEU:HD12	1:A:80:VAL:HG13	0.96	1.32	17	23
1:A:28:ALA:HB3	1:A:31:THR:HG21	0.95	1.39	29	1
1:A:9:LEU:HD13	1:A:74:PHE:CZ	0.91	2.01	30	2
1:A:59:LEU:HD12	1:A:79:LEU:HD11	0.90	1.41	12	9

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	96/99 (97%)	69±3 (72±3%)	22±3 (23±3%)	5±1 (5±1%)	4 24
All	All	2880/2970 (97%)	2076 (72%)	657 (23%)	147 (5%)	4 24

5 of 16 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	103	PHE	29
1	A	70	LYS	20
1	A	62	THR	16
1	A	88	GLU	15
1	A	13	SER	13

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	79/81 (98%)	56±3 (71±3%)	23±3 (29±3%)	1 17
All	All	2370/2430 (98%)	1679 (71%)	691 (29%)	1 17

5 of 57 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	20	GLU	30
1	A	14	SER	30
1	A	51	LEU	30
1	A	13	SER	29
1	A	38	ASP	28

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 92% for the well-defined parts and 92% for the entire structure.

7.1 Chemical shift list 1

File name: input_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1399
Number of shifts mapped to atoms	1206
Number of unparsed shifts	0
Number of shifts with mapping errors	193
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- Residue not found in structure. First 5 (of 193) occurences are reported below.

Chain	Res	Type	Atom	Shift Data		
				Value	Uncertainty	Ambiguity
A	118	ALA	HB1	1.37	-1.0	1
A	109	PRO	HD3	3.8	-1.0	2
A	119	GLU	CA	55.6	-1.0	1
A	110	LEU	HD23	0.86	-1.0	1
A	110	LEU	CA	55.2	-1.0	1

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	119	0.11 ± 0.12	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	108	0.47 ± 0.09	None needed (< 0.5 ppm)
$^{13}C'$	111	-0.26 ± 0.11	None needed (< 0.5 ppm)

Continued on next page...

Continued from previous page...

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
^{15}N	111	0.32 ± 0.28	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 92%, i.e. 1061 atoms were assigned a chemical shift out of a possible 1158. 24 out of 24 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	467/475 (98%)	188/189 (99%)	188/194 (97%)	91/92 (99%)
Sidechain	547/604 (91%)	329/352 (93%)	207/227 (91%)	11/25 (44%)
Aromatic	47/79 (59%)	25/42 (60%)	21/35 (60%)	1/2 (50%)
Overall	1061/1158 (92%)	542/583 (93%)	416/456 (91%)	103/119 (87%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, *ppm*	Expected range, *ppm*	Z-score
1	A	70	LYS	HG2	-0.45	2.67 – 0.07	-7.0

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:


RCI
0.3
0.2
0.1
0
7:LEU
10:THR
13:SER
16:ASN
19:TYR
22:TYR
25:GLN
28:ALA
31:THR
34:VAL
37:LEU
40:ALA
43:LEU
46:SER
49:PRO
52:ILE
55:LYS
58:ASP
61:ASP
64:GLY
67:VAL
70:LYS
73:PHE
76:ALA
79:LEU
82:CYS
85:ASN
88:GLU
91:LEU
94:LEU
97:ALA
100:PRO
103:PHE