



wwPDB EM Validation Summary Report ⓘ

May 25, 2024 – 09:17 am BST

PDB ID : 8QPK
EMDB ID : EMD-18555
Title : Cryo-EM Structure of Pre-B+5'ss Complex (core part)
Authors : Zhang, Z.; Kumar, V.; Dybkov, O.; Will, C.L.; Zhong, J.; Ludwig, S.; Urlaub, H.; Kastner, B.; Stark, H.; Luehrmann, R.
Deposited on : 2023-10-02
Resolution : 4.20 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis	:	0.0.1.dev92
Mogul	:	1.8.4, CSD as541be (2020)
MolProbity	:	4.02b-467
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 4.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.












Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	G	820	
2	D	142	
3	L	499	
4	A	2335	
5	R	480	
6	U	565	
7	S	800	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
8	C	972	 76%10%14%
9	5	117	 26%27%12%34%
10	4	144	 8%22%28%47%
11	X	155	 14%86%
12	z	11	 55%45%
13	6	106	 8%28%9%59%
14	N	941	 13%86%
15	B	2136	 97%
16	7	793	 95%

2 Entry composition

There are 17 unique types of molecules in this entry. The entry contains 29527 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable ATP-dependent RNA helicase DDX23.

Mol	Chain	Residues	Atoms				AltConf	Trace
1	G	102	Total	C	N	O	0	0
			658	419	128	111		

- Molecule 2 is a protein called Thioredoxin-like protein 4A.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	D	141	Total	C	N	O	S	0	0
			1008	648	184	171	5		

- Molecule 3 is a protein called U4/U6 small nuclear ribonucleoprotein Prp31.

Mol	Chain	Residues	Atoms				AltConf	Trace
3	L	50	Total	C	N	O	0	0
			251	151	50	50		

- Molecule 4 is a protein called Pre-mRNA-processing-splicing factor 8.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	A	1947	Total	C	N	O	S	0	0
			13055	8360	2412	2254	29		

- Molecule 5 is a protein called RNA-binding protein 42.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	R	106	Total	C	N	O	S	0	0
			659	412	124	121	2		

- Molecule 6 is a protein called Ubiquitin carboxyl-terminal hydrolase 39.

Mol	Chain	Residues	Atoms				AltConf	Trace
6	U	456	Total	C	N	O	0	0
			2320	1404	459	457		

- Molecule 7 is a protein called U4/U6.U5 tri-snRNP-associated protein 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	S	73	Total	C	N	O	S	0	0
			446	274	90	81	1		

- Molecule 8 is a protein called 116 kDa U5 small nuclear ribonucleoprotein component.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	C	836	Total	C	N	O	S	0	0
			5163	3262	981	913	7		

- Molecule 9 is a RNA chain called U5 snRNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	5	77	Total	C	N	O	P	0	0
			1620	726	271	546	77		

- Molecule 10 is a RNA chain called U4 snRNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	4	76	Total	C	N	O	P	0	0
			1617	723	286	532	76		

- Molecule 11 is a protein called U4/U6.U5 small nuclear ribonucleoprotein 27 kDa protein.

Mol	Chain	Residues	Atoms				AltConf	Trace
11	X	22	Total	C	N	O	0	0
			124	78	22	24		

- Molecule 12 is a RNA chain called 5'ss oligo.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	z	11	Total	C	N	O	P	0	0
			239	107	46	75	11		

- Molecule 13 is a RNA chain called U6 snRNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	6	43	Total	C	N	O	P	0	0
			927	414	175	295	43		

- Molecule 14 is a protein called Pre-mRNA-processing factor 6.

Mol	Chain	Residues	Atoms				AltConf	Trace
14	N	133	Total	C	N	O	0	0
			784	482	157	145		


- Molecule 15 is a protein called U5 small nuclear ribonucleoprotein 200 kDa helicase.

Mol	Chain	Residues	Atoms				AltConf	Trace
15	B	72	Total	C	N	O	0	0
			401	238	86	77		

- Molecule 16 is a protein called Splicing factor 3A subunit 1.

Mol	Chain	Residues	Atoms				AltConf	Trace
16	7	36	Total	C	N	O	0	0
			219	136	43	40		

- Molecule 17 is INOSITOL HEXAKISPHOSPHATE (three-letter code: IHP) (formula: $C_6H_{18}O_{24}P_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				AltConf
17	A	1	Total	C	O	P	0
			36	6	24	6	

MET	S2	L5	W12	Q13	V14	V25	R29	F30	G31	H32	D33	W34	Y46	N54	Y59	I63	N70	L75	F84	I90	M91	I92		D108	I115	I116	V119	Y120	R121	K125	G126	R127	G128	L129		R141
-----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	--	------	------	------	------	------	------	------	------	------	------	------	--	------

Chain L: 9% 90%


- Molecule 4: Pre-mRNA-processing-splicing factor 8

Chain A: 


P175	P176	D177	Y178	N181	I182	L183	L188	E199	V203	F207	P212	L213	K214	D215	S216	R217	K218	Y219	S223	F230	Y239	L241	A242	N243	Q244	L245	Y254	L260	W270	R284	D285	I286	N287	L288	Q289	D290	E291	D298	I299	N300	K301	I302	R305								
Met	Ala	Gly	Val	Phe	Pro	Tyr	Gly	Gly	Pro	Pro	Pro	Pro	Ala	Leu	Leu	Pro	Met	Ser	Glu	Glu	Lys	Leu	Gln	Lys	Ala	Ala	Arg	Lys	Lys	Trp	Gln	Gln	Leu	Gln	Lys	Lys	Arg	Tyr	Ala	Ala	Glu	Lys	Arg	Lys	Phe	Gly	Phe	Val	Asp	Asp	Met
170	D75	M76	T77	K80	D84	K85	R86	Y88	L89	G90	P96	L103	P109	L119	Y120	S127	F128	V129	N130	P133	I136	E137	P138	V139	S142	Q143	W144	G145	S146	M147	M151	R152	R153	L154	K155	R156	D157	R158	L159	F169	D170	D171	E172	E173	P174						




- Molecule 5: RNA-binding protein 42



- Molecule 6: Ubiquitin carboxyl-terminal hydrolase 39




- Molecule 7: U4/U6.U5 tri-snRNP-associated protein 1



GLN
ASP
VAL
LEU
ASN
TYR
PRO
MET


• Molecule 9: U5 snRNA

Chain 5: 26% 27% 12% 34%




• Molecule 10: U4 snRNA

Chain 4: 8% 22% 28% 47%



• Molecule 11: U4/U6.U5 small nuclear ribonucleoprotein 27 kDa protein

Chain X: 14% 86%



• Molecule 12: 5'ss oligo

Chain z: 55% 45%




• Molecule 13: U6 snRNA

Chain 6: 8% 28% 9% 59%





• Molecule 14: Pre-mRNA-processing factor 6



• Molecule 15: U5 small nuclear ribonucleoprotein 200 kDa helicase





WORLDWIDE
PDB
PROTEIN DATA BANK

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	176879	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	45	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	5000	Depositor
Magnification	Not provided	
Image detector	FEI FALCON III (4k x 4k)	Depositor
Maximum map value	0.096	Depositor
Minimum map value	-0.044	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.0165	Depositor
Map size (Å)	580.0, 580.0, 580.0	wwPDB
Map dimensions	500, 500, 500	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.16, 1.16, 1.16	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IHP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	G	0.25	0/667	0.45	0/906
2	D	0.25	0/1032	0.47	0/1409
3	L	0.23	0/253	0.35	0/352
4	A	0.25	0/13400	0.46	0/18475
5	R	0.24	0/668	0.44	0/913
6	U	0.23	0/2343	0.41	0/3285
7	S	0.24	0/450	0.46	0/610
8	C	0.25	0/5286	0.45	0/7297
9	5	0.15	0/1806	0.75	5/2807 (0.2%)
10	4	0.12	0/1805	0.66	0/2805
11	X	0.23	0/126	0.42	0/174
12	z	0.16	0/268	0.73	0/416
13	6	0.12	0/1038	0.65	0/1615
14	N	0.24	0/794	0.43	0/1086
15	B	0.23	0/401	0.49	0/544
16	7	0.23	0/220	0.47	0/300
All	All	0.23	0/30557	0.50	5/42994 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	5	68	C	N1-C2-O2	8.08	123.75	118.90
9	5	68	C	C2-N1-C1'	7.69	127.26	118.80
9	5	68	C	N3-C2-O2	-6.50	117.35	121.90
9	5	68	C	C6-N1-C1'	-5.26	114.49	120.80
9	5	68	C	C6-N1-C2	-5.22	118.21	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	658	0	505	11	0
2	D	1008	0	877	19	0
3	L	251	0	128	6	0
4	A	13055	0	10358	256	0
5	R	659	0	511	7	0
6	U	2320	0	1124	24	0
7	S	446	0	330	7	0
8	C	5163	0	3686	63	0
9	5	1620	0	822	30	0
10	4	1617	0	819	30	0
11	X	124	0	64	0	0
12	z	239	0	119	0	0
13	6	927	0	467	14	0
14	N	784	0	517	12	0
15	B	401	0	244	13	0
16	7	219	0	171	2	0
17	A	36	0	6	0	0
All	All	29527	0	20748	432	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 432 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:U:164:PHE:H	6:U:173:TYR:H	1.29	0.81
4:A:1536:LEU:HD22	4:A:1572:SER:HB3	1.64	0.77
4:A:1571:ILE:HD12	4:A:1574:ILE:HD11	1.70	0.73
4:A:136:ILE:HG13	4:A:418:THR:HG22	1.69	0.73
4:A:596:TYR:HE1	9:5:46:U:H5'	1.54	0.73

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	98/820 (12%)	96 (98%)	2 (2%)	0	100	100
2	D	139/142 (98%)	136 (98%)	3 (2%)	0	100	100
3	L	48/499 (10%)	48 (100%)	0	0	100	100
4	A	1943/2335 (83%)	1861 (96%)	81 (4%)	1 (0%)	51	85
5	R	104/480 (22%)	101 (97%)	3 (3%)	0	100	100
6	U	454/565 (80%)	441 (97%)	13 (3%)	0	100	100
7	S	69/800 (9%)	68 (99%)	1 (1%)	0	100	100
8	C	834/972 (86%)	814 (98%)	20 (2%)	0	100	100
11	X	20/155 (13%)	20 (100%)	0	0	100	100
14	N	127/941 (14%)	125 (98%)	2 (2%)	0	100	100
15	B	66/2136 (3%)	66 (100%)	0	0	100	100
16	7	34/793 (4%)	34 (100%)	0	0	100	100
All	All	3936/10638 (37%)	3810 (97%)	125 (3%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	A	80	LYS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	35/721 (5%)	35 (100%)	0	100	100
2	D	82/130 (63%)	82 (100%)	0	100	100
3	L	3/424 (1%)	3 (100%)	0	100	100
4	A	877/2108 (42%)	870 (99%)	7 (1%)	81	89
5	R	40/369 (11%)	39 (98%)	1 (2%)	47	68
6	U	26/511 (5%)	26 (100%)	0	100	100
7	S	23/681 (3%)	21 (91%)	2 (9%)	10	34
8	C	266/866 (31%)	265 (100%)	1 (0%)	91	94
11	X	3/144 (2%)	3 (100%)	0	100	100
14	N	29/792 (4%)	29 (100%)	0	100	100
15	B	12/1908 (1%)	12 (100%)	0	100	100
16	7	11/709 (2%)	11 (100%)	0	100	100
All	All	1407/9363 (15%)	1396 (99%)	11 (1%)	82	89

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	R	460	LYS
7	S	718	ARG
8	C	529	ARG
7	S	736	LYS
4	A	616	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
4	A	545	HIS
4	A	610	HIS
7	S	731	HIS
4	A	723	ASN
4	A	495	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
10	4	74/144 (51%)	16 (21%)	2 (2%)
12	z	10/11 (90%)	5 (50%)	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
13	6	41/106 (38%)	5 (12%)	0
9	5	76/117 (64%)	27 (35%)	1 (1%)
All	All	201/378 (53%)	53 (26%)	3 (1%)

5 of 53 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
9	5	8	G
9	5	9	G
9	5	10	U
9	5	12	U
9	5	14	U

All (3) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
9	5	19	A
10	4	1	A
10	4	68	A

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
17	IHP	A	2401	-	36,36,36	1.20	4 (11%)	54,60,60	1.64	6 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
17	IHP	A	2401	-	-	1/30/54/54	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
17	A	2401	IHP	P5-O15	2.60	1.64	1.59
17	A	2401	IHP	P1-O11	2.26	1.63	1.59
17	A	2401	IHP	C4-C3	2.19	1.56	1.52
17	A	2401	IHP	C5-C4	2.17	1.56	1.52

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
17	A	2401	IHP	C5-C6-C1	5.05	121.47	110.41
17	A	2401	IHP	C3-C2-C1	4.61	120.50	110.41
17	A	2401	IHP	C5-C4-C3	4.38	120.01	110.41
17	A	2401	IHP	C4-C3-C2	3.47	118.00	110.41
17	A	2401	IHP	O15-C5-C4	2.42	114.40	108.69

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
17	A	2401	IHP	C5-O15-P5-O25

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-18555. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 250




Y Index: 250




Z Index: 250


6.2.2 Raw map



X Index: 250



Y Index: 250




Z Index: 250


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 153




Y Index: 214




Z Index: 276


6.3.2 Raw map



X Index: 153



Y Index: 213




Z Index: 276


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0165. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 494 nm³; this corresponds to an approximate mass of 446 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.238 \AA^{-1}

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.238 \AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.20	-	-
Author-provided FSC curve	4.10	6.37	4.23
Unmasked-calculated*	5.46	7.69	6.18

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 5.46 differs from the reported value 4.2 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-18555 and PDB model 8QPK. Per-residue inclusion information can be found in [section 3](#) on [page 7](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.0165 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0165).



































9.4 Atom inclusion [i](#)



At the recommended contour level, 92% of all backbone atoms, 88% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.0165) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8750	 0.3650
4	 0.7590	 0.2470
5	 0.9560	 0.3240
6	 0.7370	 0.2920
7	 0.8780	 0.4000
A	 0.8620	 0.3820
B	 0.7770	 0.3570
C	 0.9500	 0.3930
D	 0.8950	 0.4320
G	 0.8130	 0.3570
L	 0.6570	 0.3490
N	 0.9380	 0.3830
R	 0.9110	 0.3380
S	 0.7060	 0.3240
U	 0.9150	 0.3240
X	 0.7540	 0.4020
z	 0.9620	 0.3970

