Full wwPDB X-ray Structure Validation Report (i)

Jan 14, 2024 – 12:10 pm GMT

PDB ID	:	6S2R
Title	:	Mycobacterial hydrolase 2
Authors	:	Cereija, T.B.; Vilela, F.; Macedo-Ribeiro, S.; Pereira, P.J.B.
Deposited on	:	2019-06-21
Resolution	:	1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1156 (1.46-1.46)
Clashscore	141614	1202 (1.46-1.46)
Ramachandran outliers	138981	1178 (1.46-1.46)
Sidechain outliers	138945	1178 (1.46-1.46)
RSRZ outliers	127900	1139 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	236	11% 89% 9%
1	B	236	8% 89% 10%

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 4230 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucosyl-3-phosphoglycerate phosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	234	1918	1194	372	347	5	0	16	0
1	B	233	1835	1138	357	336	4	0	5	0

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	224	LYS	-	expression tag	UNP P9WIC7
A	225	LEU	-	expression tag	UNP P9WIC7
A	226	ALA	-	expression tag	UNP P9WIC7
A	227	ALA	-	expression tag	UNP P9WIC7
A	228	ALA	-	expression tag	UNP P9WIC7
A	229	LEU	-	expression tag	UNP P9WIC7
A	230	GLU	-	expression tag	UNP P9WIC7
A	231	HIS	-	expression tag	UNP P9WIC7
A	232	HIS	-	expression tag	UNP P9WIC7
A	233	HIS	-	expression tag	UNP P9WIC7
A	234	HIS	-	expression tag	UNP P9WIC7
A	235	HIS	-	expression tag	UNP P9WIC7
A	236	HIS	-	expression tag	UNP P9WIC7
B	224	LYS	-	expression tag	UNP P9WIC7
B	225	LEU	-	expression tag	UNP P9WIC7
B	226	ALA	-	expression tag	UNP P9WIC7
B	227	ALA	-	expression tag	UNP P9WIC7
B	228	ALA	-	expression tag	UNP P9WIC7
B	229	LEU	-	expression tag	UNP P9WIC7
B	230	GLU	-	expression tag	UNP P9WIC7
B	231	HIS	-	expression tag	UNP P9WIC7
B	232	HIS	-	expression tag	UNP P9WIC7
B	233	HIS	-	expression tag	UNP P9WIC7
B	234	HIS	-	expression tag	UNP P9WIC7
B	235	HIS	-	expression tag	UNP P9WIC7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	236	HIS	-	expression tag	UNP P9WIC7

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	6	3	3	0	0
			Total	C	O		
2	A	1	6	3	3	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
3	B	3	3	3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	218	218	218	0	5
			Total	O		
4	B	244	244	244	0	2

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glucosyl-3-phosphoglycerate phosphatase

Chain A:





- Molecule 1: Glucosyl-3-phosphoglycerate phosphatase

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	98.95Å 48.83Å 93.48Å 90.00° 104.33° 90.00°	Depositor
Resolution (Å)	47.51 – 1.45 47.51 – 1.45	Depositor EDS
% Data completeness (in resolution range)	98.9 (47.51-1.45) 98.9 (47.51-1.45)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$< I/\sigma(I) >$ [1]	1.40 (at 1.45Å)	Xtriage
Refinement program	PHENIX (1.15.2_3472: ???)	Depositor
R, R_{free}	0.174 , 0.200 0.174 , 0.200	Depositor DCC
R_{free} test set	3677 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å^2)	19.2	Xtriage
Anisotropy	0.421	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.36 , 56.7	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	4230	wwPDB-VP
Average B, all atoms (Å^2)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.07% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.45	0/1962	0.66	0/2670
1	B	0.42	0/1874	0.65	0/2548
All	All	0.43	0/3836	0.66	0/5218

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1918	0	1890	14	0
1	B	1835	0	1814	14	0
2	A	12	0	16	0	0
3	B	3	0	0	0	0
4	A	218	0	0	2	0
4	B	244	0	0	2	0
All	All	4230	0	3720	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:200:SER:HB2	1:A:204:SER:HB2	1.78	0.64
1:B:48[B]:ARG:NH2	1:B:191:GLN:OE1	2.31	0.62
1:A:218:VAL:HG13	1:A:222:VAL:HG23	1.85	0.59
1:B:113:ALA:HB3	1:B:175:VAL:HG13	1.84	0.59
1:A:163:ILE:HG21	1:A:184[B]:MET:SD	2.44	0.57
1:B:106:ARG:O	1:B:110:ARG:HG2	2.04	0.57
1:B:10[B]:ARG:NH2	1:B:186:ASN:OD1	2.36	0.57
1:A:4:ARG:HG3	1:A:196:TRP:CZ3	2.41	0.56
1:A:123[B]:ARG:HG2	1:A:165:ALA:HB2	1.91	0.52
1:A:115[A]:TRP:CD1	1:A:117[A]:PRO:HD3	2.46	0.51
1:A:110:ARG:HD2	1:B:208[B]:ARG:NH1	2.25	0.51
1:B:131:ARG:NH1	1:B:203:GLU:OE1	2.37	0.49
1:B:10[A]:ARG:O	1:B:187:ALA:HA	2.12	0.48
1:A:219:SER:OG	1:A:222:VAL:HG22	2.15	0.47
1:B:115:TRP:CZ2	1:B:117:PRO:HB3	2.51	0.46
1:B:106:ARG:HH11	1:B:110:ARG:NH2	2.14	0.46
1:A:15:ASP:OD1	4:A:401:HOH:O	2.20	0.45
1:B:38:ALA:HB2	4:B:410:HOH:O	2.18	0.43
1:A:208:ARG:HD2	4:A:467:HOH:O	2.19	0.43
1:A:32:GLU:OE2	1:A:35:ARG:NH1	2.51	0.42
1:A:43:GLU:HG2	1:A:71[B]:ARG:CZ	2.50	0.42
1:B:12:GLY:N	4:B:410:HOH:O	2.46	0.41
1:B:20:SER:HB2	1:B:95:HIS:CD2	2.56	0.41
1:A:2:ARG:HH11	1:A:198:PRO:HG3	1.85	0.41
1:B:115:TRP:CD2	1:B:117:PRO:HD3	2.56	0.41
1:B:20:SER:HB2	1:B:95:HIS:CE1	2.56	0.40
1:A:2:ARG:HB2	1:A:196:TRP:CH2	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/236 (105%)	244 (98%)	3 (1%)	1 (0%)	34	13
1	B	236/236 (100%)	226 (96%)	8 (3%)	2 (1%)	19	4
All	All	484/472 (102%)	470 (97%)	11 (2%)	3 (1%)	25	7

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	16	TYR
1	B	158	ALA
1	A	158	ALA

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/182 (105%)	190 (100%)	1 (0%)	88	75
1	B	184/182 (101%)	180 (98%)	4 (2%)	52	18
All	All	375/364 (103%)	370 (99%)	5 (1%)	73	40

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	ARG
1	B	2	ARG
1	B	148[A]	ASP
1	B	148[B]	ASP
1	B	162	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	GOL	A	301	-	5,5,5	0.87	0	5,5,5	0.98	0
2	GOL	A	302	-	5,5,5	0.94	0	5,5,5	0.99	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	301	-	-	0/4/4/4	-
2	GOL	A	302	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	302	GOL	C1-C2-C3-O3
2	A	302	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	234/236 (99%)	0.61	26 (11%) 5 6	15, 24, 69, 126	0
1	B	233/236 (98%)	0.64	18 (7%) 13 16	15, 26, 68, 94	0
All	All	467/472 (98%)	0.62	44 (9%) 8 10	15, 25, 69, 126	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	197	ALA	9.9
1	B	200	SER	9.6
1	B	198	PRO	9.5
1	B	232	HIS	9.4
1	B	1	MET	8.6
1	B	19	GLY	7.8
1	A	225	LEU	6.5
1	B	196	TRP	6.3
1	A	233	HIS	6.1
1	A	232	HIS	5.8
1	B	20	SER	5.4
1	A	218	VAL	5.0
1	B	2	ARG	5.0
1	A	220	SER	5.0
1	A	147	ALA	4.9
1	B	201	ASP	4.9
1	A	227	ALA	4.7
1	B	199	GLY	4.6
1	B	233	HIS	4.3
1	A	196	TRP	4.3
1	A	223	LEU	4.3
1	A	197	ALA	4.2
1	B	17	ASN	4.0
1	B	18	VAL	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	228	ALA	3.9
1	A	200	SER	3.9
1	A	146	GLY	3.8
1	A	222	VAL	3.7
1	A	198	PRO	3.6
1	A	234	HIS	3.6
1	A	229	LEU	3.3
1	A	219	SER	3.2
1	A	221	ASP	3.1
1	A	148	ASP	3.1
1	A	199	GLY	3.0
1	B	90	TRP	3.0
1	A	226	ALA	3.0
1	B	117	PRO	3.0
1	B	231	HIS	2.9
1	A	115[A]	TRP	2.6
1	A	231	HIS	2.4
1	A	1	MET	2.3
1	A	224	LYS	2.2
1	B	107	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	GOL	A	301	6/6	0.81	0.12	37,49,55,58	0
2	GOL	A	302	6/6	0.88	0.13	54,57,61,62	0
3	CA	B	301	1/1	0.97	0.07	30,30,30,30	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	CA	B	303	1/1	0.97	0.10	30,30,30,30	1
3	CA	B	302	1/1	0.99	0.06	25,25,25,25	0

6.5 Other polymers (i)

There are no such residues in this entry.