



Full wwPDB X-ray Structure Validation Report i

Aug 14, 2023 – 11:43 PM EDT

PDB ID : 1S7S

Title : Crystal structures of the murine class I major histocompatibility complex H-2K_b in complex with LCMV-derived gp33 index peptide and three of its escape variants

Authors : Velloso, L.M.; Michaelsson, J.; Ljunggren, H.G.; Schneider, G.; Achour, A.

Deposited on : 2004-01-30

Resolution : 1.99 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467

Xtriage (Phenix) : 1.13

EDS : 2.35

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition [\(i\)](#)

There are 4 unique types of molecules in this entry. The entry contains 3399 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called H-2 class I histocompatibility antigen, K-B alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	276	Total	C 2247	N 1418	O 395	S 425	9	0	0

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	99	Total	C 820	N 524	O 138	S 151	7	0	0

- Molecule 3 is a protein called Glycoprotein 9-residue peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	8	Total	C 64	N 43	O 9	S 11	1	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	3	LEU	VAL	engineered mutation	UNP P07399
C	9	MET	CYS	SEE REMARK 999	UNP P07399


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	183	Total O 183 183	0	0
4	B	77	Total O 77 77	0	0
4	C	8	Total O 8 8	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: H-2 class I histocompatibility antigen, K-B alpha chain



- Molecule 2: Beta-2-microglobulin



- Molecule 3: Glycoprotein 9-residue peptide



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	88.26 Å 135.56 Å 45.13 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.88 – 1.99 19.87 – 1.99	Depositor EDS
% Data completeness (in resolution range)	92.9 (19.88-1.99) 92.9 (19.87-1.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	2.47 (at 1.99 Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R , R_{free}	0.190 , 0.232 0.203 , 0.244	Depositor DCC
R_{free} test set	1750 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	25.2	Xtriage
Anisotropy	0.329	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 49.9	EDS
L-test for twinning ²	$< L > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3399	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.76	0/2309	0.99	8/3137 (0.3%)
2	B	0.77	0/846	0.89	1/1148 (0.1%)
3	C	1.16	0/65	1.34	2/87 (2.3%)
All	All	0.78	0/3220	0.97	11/4372 (0.3%)

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	155	ARG	NE-CZ-NH2	-15.08	112.76	120.30
1	A	155	ARG	NE-CZ-NH1	12.32	126.46	120.30
3	C	3	LEU	CB-CG-CD1	7.39	123.56	111.00
1	A	272	LEU	CA-CB-CG	6.87	131.10	115.30
1	A	227	ASP	CB-CG-OD2	6.10	123.79	118.30
1	A	29	ASP	CB-CA-C	-5.98	98.44	110.40
1	A	39	ASP	CB-CG-OD2	5.80	123.52	118.30
3	C	3	LEU	CB-CG-CD2	-5.24	102.08	111.00
1	A	14	ARG	NE-CZ-NH2	-5.12	117.74	120.30
1	A	202	ARG	NE-CZ-NH1	5.09	122.84	120.30
2	B	76	ASP	CB-CG-OD2	5.02	122.82	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2247	0	2136	28	0
2	B	820	0	796	11	0
3	C	64	0	60	14	0
4	A	183	0	0	6	0
4	B	77	0	0	6	0
4	C	8	0	0	0	0
All	All	3399	0	2992	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:45:TYR:OH	3:C:3:LEU:CD2	1.85	1.24
1:A:45:TYR:OH	3:C:3:LEU:HD21	0.97	1.11
1:A:45:TYR:HH	3:C:3:LEU:HD21	1.08	0.86
1:A:45:TYR:CZ	3:C:3:LEU:HD21	2.15	0.80
3:C:3:LEU:H	3:C:3:LEU:HD12	1.49	0.78
1:A:50:ARG:HD2	4:A:521:HOH:O	1.85	0.77
1:A:24:GLU:OE1	3:C:3:LEU:HD22	1.85	0.74
4:A:351:HOH:O	3:C:3:LEU:HD13	1.92	0.68
1:A:189:VAL:HG23	1:A:272:LEU:HD22	1.78	0.65
1:A:126:LEU:HD22	1:A:156:LEU:HD13	1.77	0.65
2:B:97:ARG:NE	4:B:174:HOH:O	2.22	0.65
3:C:3:LEU:H	3:C:3:LEU:CD1	2.08	0.65
1:A:86:ASN:ND2	4:A:376:HOH:O	2.33	0.62
2:B:48:LYS:O	2:B:48:LYS:HG2	2.04	0.58
1:A:263:HIS:CD2	1:A:265:GLY:H	2.22	0.58
1:A:171:TYR:OH	3:C:2:ALA:HB2	2.04	0.57
2:B:73:THR:HG23	4:B:132:HOH:O	2.06	0.56
2:B:36:GLU:OE1	4:B:160:HOH:O	2.18	0.56
1:A:151:GLY:O	1:A:154:GLU:HG2	2.07	0.54
1:A:259:CYS:HB3	1:A:272:LEU:HD13	1.90	0.54
1:A:7:TYR:CD1	3:C:3:LEU:HD23	2.43	0.53
1:A:210:PRO:O	1:A:263:HIS:HE1	1.91	0.53
2:B:44:LYS:NZ	4:B:140:HOH:O	2.42	0.52
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.45	0.51
2:B:31:HIS:ND1	2:B:32:PRO:HA	2.24	0.51
1:A:72:GLN:HE22	1:A:75:ARG:CZ	2.23	0.51
1:A:195:PRO:O	1:A:196:GLU:CB	2.58	0.51
1:A:263:HIS:HD2	1:A:265:GLY:H	1.58	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:95:ILE:HD13	3:C:9:MET:HE3	1.94	0.49
1:A:7:TYR:CZ	3:C:3:LEU:HG	2.48	0.48
1:A:171:TYR:OH	3:C:2:ALA:CB	2.62	0.48
1:A:275:GLU:HG2	1:A:275:GLU:O	2.14	0.46
2:B:32:PRO:HB2	2:B:33:PRO:HD2	1.97	0.46
2:B:73:THR:HB	2:B:76:ASP:OD2	2.17	0.45
1:A:31:THR:HG23	1:A:209:TYR:OH	2.19	0.43
1:A:185:PRO:HD3	1:A:263:HIS:CD2	2.55	0.42
1:A:264:GLN:NE2	4:A:475:HOH:O	2.53	0.42
1:A:121:CYS:SG	4:A:522:HOH:O	2.13	0.41
2:B:50:GLU:HB3	4:B:136:HOH:O	2.20	0.41
1:A:8:PHE:CE1	1:A:98:ILE:HD12	2.56	0.41
2:B:72:PRO:HA	4:B:175:HOH:O	2.21	0.41
4:A:349:HOH:O	3:C:3:LEU:HD11	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	274/348 (79%)	267 (97%)	6 (2%)	1 (0%)	34 30
2	B	97/99 (98%)	97 (100%)	0	0	100 100
3	C	6/9 (67%)	6 (100%)	0	0	100 100
All	All	377/456 (83%)	370 (98%)	6 (2%)	1 (0%)	41 37

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	29	ASP

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	234/290 (81%)	216 (92%)	18 (8%)	13 8
2	B	94/94 (100%)	88 (94%)	6 (6%)	17 13
3	C	6/7 (86%)	5 (83%)	1 (17%)	2 1
All	All	334/391 (85%)	309 (92%)	25 (8%)	13 9

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	41	GLU
1	A	44	ARG
1	A	45	TYR
1	A	79	ARG
1	A	89	LYS
1	A	108	ARG
1	A	110	LEU
1	A	134	THR
1	A	155	ARG
1	A	156	LEU
1	A	165	VAL
1	A	222	GLU
1	A	225	ILE
1	A	227	ASP
1	A	230	LEU
1	A	255	GLN
1	A	272	LEU
1	A	275	GLU
2	B	1	ILE
2	B	48	LYS
2	B	56	PHE
2	B	58	LYS
2	B	73	THR
2	B	81	ARG
3	C	3	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	72	GLN
1	A	115	GLN
1	A	263	HIS
2	B	17	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	276/348 (79%)	0.36	22 (7%) 12 11	15, 22, 36, 48	0
2	B	99/99 (100%)	0.37	4 (4%) 38 37	17, 23, 32, 39	0
3	C	8/9 (88%)	0.53	1 (12%) 3 3	17, 20, 24, 25	0
All	All	383/456 (83%)	0.37	27 (7%) 16 15	15, 22, 34, 48	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	225	ILE	6.5
1	A	226	GLN	6.0
1	A	220	ASN	5.3
1	A	41	GLU	4.9
2	B	48	LYS	4.3
1	A	196	GLU	4.2
1	A	42	ASN	4.2
1	A	276	PRO	4.1
1	A	275	GLU	3.2
2	B	58	LYS	3.2
1	A	256	TYR	3.2
1	A	221	GLY	3.0
2	B	1	ILE	2.9
3	C	3	LEU	2.9
1	A	217	TRP	2.7
1	A	89	LYS	2.5
1	A	255	GLN	2.5
1	A	227	ASP	2.3
1	A	274	TRP	2.3
1	A	86	ASN	2.3
1	A	252	GLY	2.2
1	A	222	GLU	2.2
2	B	88	ALA	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	88	SER	2.1
1	A	79	ARG	2.0
1	A	58	GLU	2.0
1	A	195	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.