



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 15, 2023 – 04:08 AM EDT

PDB ID : 1SMS
Title : Structure of the Ribonucleotide Reductase Rnr4 Homodimer from *Saccharomyces cerevisiae*
Authors : Sommerhalter, M.; Voegtli, W.C.; Perlstein, D.L.; Ge, J.; Stubbe, J.; Rosenzweig, A.C.
Deposited on : 2004-03-09
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	345	 27% 52% 10% • 9%
1	B	345	 27% 52% 10% • 9%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 5182 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribonucleoside-diphosphate reductase small chain 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	314	2589	1679	416	480	14	0	0	0
1	B	314	2589	1679	416	480	14	0	0	0

- Molecule 2 is MERCURY (II) ION (three-letter code: HG) (formula: Hg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total 2	Hg 2	0	0
2	B	2	Total 2	Hg 2	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: Ribonucleoside-diphosphate reductase small chain 2

Chain A: 



- Molecule 1: Ribonucleoside-diphosphate reductase small chain 2

Chain B: 



F267	MET
G268	SER
M269	LYS
D270	SER
L271	ALA
K272	THR
S273	PRO
I274	SER
H275	LYS
T276	GLU
Y277	ILE
I278	ASN
E279	PHE
F280	ASP
V281	ASP
A282	ASP
D283	ASP
G284	THR
L285	THR
L286	GLY
Q287	LYS
G288	THR
F289	THR
G290	THR
N291	THR
E292	THR
K293	THR
Y294	THR
Y295	THR
N299	THR
F300	THR
F301	THR
E302	THR
F303	THR
M304	THR
E305	THR
ASP	THR
VAL	THR
ALA	THR
THR	THR
ALA	THR
ALA	THR
GLY	THR
LYS	THR
THR	THR
THR	THR
F315	THR
F316	THR
E317	THR
K318	THR
K319	THR
V320	THR
S321	THR
D322	THR
Y323	THR
Q324	THR
K325	THR
ALA	THR
SER	THR
ASP	THR

MET
SER
LYS
SER
ALA
THR
PRO
SER
LYS
GLU
ILE
ASN
PHE
ASP
ASP
ASP
PHE

4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	79.60Å 79.60Å 218.10Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	12.00 – 3.10	Depositor
% Data completeness (in resolution range)	(Not available) (12.00-3.10)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.266 , 0.305	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	5182	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/2652	0.66	2/3569 (0.1%)
1	B	0.43	0/2652	0.66	2/3569 (0.1%)
All	All	0.43	0/5304	0.66	4/7138 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	37	MET	N-CA-C	-6.42	93.67	111.00
1	A	37	MET	N-CA-C	-6.39	93.73	111.00
1	A	302	GLU	CB-CA-C	5.33	121.05	110.40
1	B	302	GLU	CB-CA-C	5.32	121.03	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2589	0	2530	232	0
1	B	2589	0	2530	230	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
All	All	5182	0	5060	456	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 45.

The worst 5 of 456 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:147:LEU:HD11	1:B:154:LEU:HD11	1.34	1.07
1:A:147:LEU:HD11	1:A:154:LEU:HD11	1.34	1.04
1:A:26:ILE:HA	1:A:29:MET:HB2	1.44	0.99
1:B:26:ILE:HA	1:B:29:MET:HB2	1.44	0.97
1:A:239:ASN:ND2	1:A:241:LYS:H	1.68	0.91

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	310/345 (90%)	225 (73%)	51 (16%)	34 (11%)	0	2
1	B	310/345 (90%)	224 (72%)	52 (17%)	34 (11%)	0	2
All	All	620/690 (90%)	449 (72%)	103 (17%)	68 (11%)	0	2

5 of 68 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	33	ARG
1	A	38	PHE
1	A	39	PRO
1	A	200	LEU
1	A	263	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	276/303 (91%)	248 (90%)	28 (10%)	7	28
1	B	276/303 (91%)	250 (91%)	26 (9%)	8	32
All	All	552/606 (91%)	498 (90%)	54 (10%)	8	29

5 of 54 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	16	HIS
1	B	63	GLU
1	B	275	HIS
1	B	20	GLU
1	B	33	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	6	GLN
1	B	31	ASN
1	B	239	ASN
1	B	120	GLN
1	B	190	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.