



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 19, 2023 – 07:16 PM EST

PDB ID : 1V26
Title : Crystal structure of tt0168 from *Thermus thermophilus* HB8
Authors : Hisanaga, Y.; Ago, H.; Nakatsu, T.; Hamada, K.; Ida, K.; Kanda, H.; Yamamoto, M.; Hori, T.; Arii, Y.; Sugahara, M.; Kuramitsu, S.; Yokoyama, S.; Miyano, M.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2003-10-07
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	541	 6% 79% 11% 10%
1	B	541	 6% 83% 11% 6%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8316 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called long-chain-fatty-acid-CoA synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	489	Total 3802	C 2436	N 661	O 695	S 10	0	0	0
1	B	510	Total 3964	C 2536	N 688	O 730	S 10	0	0	0

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Mg 1	0	0
2	B	1	Total 1	Mg 1	0	0

- Molecule 3 is MYRISTIC ACID (three-letter code: MYR) (formula: C₁₄H₂₈O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 15 14 1	0	0
3	B	1	Total C O 15 14 1	0	0

- Molecule 4 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula: C₁₀H₁₄N₅O₇P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O P 23 10 5 7 1	0	0
4	B	1	Total C N O P 23 10 5 7 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	251	Total O 251 251	0	0
5	B	221	Total O 221 221	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: long-chain-fatty-acid-CoA synthetase



- Molecule 1: long-chain-fatty-acid-CoA synthetase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	64.51Å 101.38Å 176.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.10 – 2.50 46.35 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (46.10-2.50) 99.7 (46.35-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.66 (at 2.51Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.204 , 0.240 0.203 , 0.237	Depositor DCC
R_{free} test set	4116 reflections (10.04%)	wwPDB-VP
Wilson B-factor (Å ²)	19.5	Xtrriage
Anisotropy	0.329	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 38.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	8316	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AMP, MYR, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/3890	0.63	8/5285 (0.2%)
1	B	0.29	0/4056	0.62	6/5511 (0.1%)
All	All	0.29	0/7946	0.62	14/10796 (0.1%)

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	473	TRP	C-N-CA	-6.90	104.45	121.70
1	A	473	TRP	C-N-CA	-6.55	105.31	121.70
1	B	432	ASP	CB-CG-OD2	5.94	123.65	118.30
1	A	432	ASP	CB-CG-OD2	5.92	123.62	118.30
1	A	126	ASP	CB-CG-OD2	5.50	123.25	118.30
1	B	15	ASP	CB-CG-OD2	5.42	123.17	118.30
1	B	126	ASP	CB-CG-OD2	5.39	123.15	118.30
1	A	212	ASP	CB-CG-OD2	5.33	123.10	118.30
1	A	120	ASP	CB-CG-OD2	5.24	123.02	118.30
1	A	449	ASP	CB-CG-OD2	5.23	123.01	118.30
1	B	120	ASP	CB-CG-OD2	5.19	122.97	118.30
1	A	15	ASP	CB-CG-OD2	5.06	122.86	118.30
1	A	255	ASP	CB-CG-OD2	5.05	122.84	118.30
1	B	509	ASP	CB-CG-OD2	5.04	122.83	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3802	0	3814	28	0
1	B	3964	0	3963	32	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	15	0	27	3	0
3	B	15	0	27	2	0
4	A	23	0	12	0	0
4	B	23	0	12	0	0
5	A	251	0	0	1	0
5	B	221	0	0	0	0
All	All	8316	0	7855	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:471:PRO:HD3	1:B:540:GLY:HA2	1.24	1.14
1:B:438:ILE:HG21	1:B:477:PRO:HD3	1.48	0.96
1:B:330:SER:N	1:B:331:PRO:HA	2.01	0.76
1:B:225:VAL:HG11	1:B:277:VAL:HG11	1.67	0.75
1:B:471:PRO:HD3	1:B:540:GLY:CA	2.12	0.74
1:A:330:SER:N	1:A:331:PRO:HA	2.04	0.72
1:A:310:ILE:O	1:A:314:GLU:HG2	1.92	0.70
1:B:372:GLU:HB2	1:B:374:ARG:HD3	1.79	0.63
1:A:308:SER:O	1:A:312:ARG:HB2	2.00	0.61
1:B:19:ASN:HD22	1:B:21:TRP:H	1.47	0.61
1:B:275:PRO:HB2	1:B:474:GLN:NE2	2.16	0.61
1:A:474:GLN:HG3	5:A:3091:HOH:O	2.01	0.59
1:A:446:SER:HB3	1:A:449:ASP:HB2	1.85	0.58
1:B:305:ALA:H	1:B:322:GLN:NE2	2.01	0.58
1:B:92:VAL:HB	1:B:93:PRO:HD3	1.85	0.57
1:B:329:THR:C	1:B:331:PRO:HA	2.26	0.56
1:B:335:GLN:HE21	1:B:361:ILE:HG21	1.70	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:82:HIS:HE1	1:B:126:ASP:OD1	1.89	0.56
1:B:471:PRO:CD	1:B:540:GLY:HA2	2.17	0.54
1:A:19:ASN:HD22	1:A:21:TRP:H	1.56	0.54
1:B:128:ASN:H	1:B:128:ASN:HD22	1.55	0.52
1:A:264:ASP:HB3	1:A:293:LYS:HE3	1.91	0.52
1:B:293:LYS:HG3	1:B:294:THR:HG23	1.92	0.52
1:A:92:VAL:HB	1:A:93:PRO:HD3	1.91	0.51
1:B:438:ILE:HD13	1:B:477:PRO:HG3	1.91	0.51
1:A:329:THR:C	1:A:331:PRO:HA	2.32	0.50
1:A:419:ILE:HD11	1:A:434:LEU:HG	1.94	0.49
1:A:438:ILE:HD13	1:A:477:PRO:HB3	1.94	0.49
1:B:305:ALA:H	1:B:322:GLN:HE22	1.61	0.48
1:B:275:PRO:HB2	1:B:474:GLN:HE22	1.79	0.48
1:B:125:PHE:CZ	1:B:148:VAL:HG22	2.48	0.47
1:B:214:THR:HG21	3:B:2001:MYR:H111	1.96	0.47
1:A:228:MET:HA	1:A:233:ALA:HB2	1.97	0.46
1:A:82:HIS:HE1	1:A:126:ASP:OD1	1.98	0.46
1:A:128:ASN:HD22	1:A:128:ASN:H	1.63	0.46
1:B:477:PRO:HB2	1:B:508:PRO:HA	1.98	0.46
1:A:23:PHE:CE1	1:A:242:LEU:HD13	2.52	0.45
1:A:477:PRO:HG2	1:A:508:PRO:HA	1.99	0.45
1:B:234:TRP:CD2	3:B:2001:MYR:H62	2.51	0.45
1:A:305:ALA:H	1:A:322:GLN:NE2	2.14	0.45
1:A:127:PRO:HG3	1:A:150:ASP:HB2	1.99	0.45
1:A:323:GLY:HA3	3:A:1001:MYR:H72	1.99	0.44
1:B:419:ILE:HB	1:B:432:ASP:HB3	1.99	0.44
1:B:372:GLU:HG3	1:B:374:ARG:NH1	2.33	0.44
1:A:234:TRP:CE2	3:A:1001:MYR:H51	2.53	0.43
1:B:128:ASN:H	1:B:128:ASN:ND2	2.16	0.43
1:A:225:VAL:HG11	1:A:277:VAL:HG11	1.99	0.43
1:B:498:LEU:HD21	1:B:507:LEU:HD11	2.00	0.43
1:A:77:THR:HA	1:A:124:LEU:O	2.18	0.43
1:A:370:ASP:OD2	1:A:374:ARG:HB2	2.19	0.42
1:B:77:THR:HA	1:B:124:LEU:O	2.20	0.42
1:A:128:ASN:H	1:A:128:ASN:ND2	2.18	0.41
1:A:153:ALA:HA	1:A:154:PRO:HD3	1.91	0.41
1:A:305:ALA:H	1:A:322:GLN:HE22	1.69	0.41
1:A:299:VAL:HG11	3:A:1001:MYR:H101	2.03	0.40
1:B:185:THR:HG22	1:B:191:PRO:HG3	2.03	0.40
1:B:225:VAL:HG13	1:B:254:LEU:HD22	2.03	0.40
1:B:419:ILE:HD11	1:B:434:LEU:HG	2.04	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:103:ASN:HA	1:B:104:PRO:HD3	1.89	0.40
1:A:225:VAL:HG13	1:A:254:LEU:HD22	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	481/541 (89%)	469 (98%)	11 (2%)	1 (0%)	47 68
1	B	502/541 (93%)	489 (97%)	11 (2%)	2 (0%)	34 54
All	All	983/1082 (91%)	958 (98%)	22 (2%)	3 (0%)	41 61

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	330	SER
1	B	330	SER
1	B	518	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	395/437 (90%)	380 (96%)	15 (4%)	33 58

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	411/437 (94%)	396 (96%)	15 (4%)	35	61
All	All	806/874 (92%)	776 (96%)	30 (4%)	34	60

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55	GLN
1	A	109	LYS
1	A	172	ARG
1	A	185	THR
1	A	187	THR
1	A	231	VAL
1	A	235	CYS
1	A	236	LEU
1	A	283	ASP
1	A	291	ARG
1	A	336	ASN
1	A	422	TRP
1	A	435	LYS
1	A	449	ASP
1	A	526	LEU
1	B	55	GLN
1	B	74	ARG
1	B	109	LYS
1	B	128	ASN
1	B	139	GLU
1	B	155	GLU
1	B	172	ARG
1	B	223	LEU
1	B	231	VAL
1	B	235	CYS
1	B	236	LEU
1	B	303	SER
1	B	422	TRP
1	B	435	LYS
1	B	448	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	19	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	55	GLN
1	A	82	HIS
1	A	128	ASN
1	A	144	GLN
1	A	322	GLN
1	A	335	GLN
1	A	336	ASN
1	B	19	ASN
1	B	55	GLN
1	B	82	HIS
1	B	128	ASN
1	B	145	HIS
1	B	232	ASN
1	B	322	GLN
1	B	335	GLN
1	B	336	ASN
1	B	474	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MYR	A	1001	4	14,14,15	0.29	0	13,13,15	0.73	0
3	MYR	B	2001	4	14,14,15	0.28	0	13,13,15	0.70	0
4	AMP	A	1002	2,3	22,25,25	1.18	2 (9%)	25,38,38	1.28	2 (8%)
4	AMP	B	2002	2,3	22,25,25	1.18	2 (9%)	25,38,38	1.34	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MYR	A	1001	4	-	6/11/12/13	-
3	MYR	B	2001	4	-	7/11/12/13	-
4	AMP	A	1002	2,3	-	0/6/26/26	0/3/3/3
4	AMP	B	2002	2,3	-	0/6/26/26	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	2002	AMP	O4'-C1'	3.89	1.46	1.41
4	A	1002	AMP	O4'-C1'	3.82	1.46	1.41
4	B	2002	AMP	P-O1P	2.71	1.59	1.50
4	A	1002	AMP	P-O1P	2.67	1.59	1.50

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	2002	AMP	N3-C2-N1	-4.76	121.23	128.68
4	A	1002	AMP	N3-C2-N1	-4.49	121.66	128.68
4	B	2002	AMP	C3'-C2'-C1'	2.05	104.06	100.98
4	B	2002	AMP	C4-C5-N7	-2.03	107.28	109.40
4	A	1002	AMP	C4-C5-N7	-2.02	107.30	109.40

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	2001	MYR	C5-C6-C7-C8
3	B	2001	MYR	C7-C8-C9-C10
3	A	1001	MYR	C6-C7-C8-C9

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	B	2001	MYR	C6-C7-C8-C9
3	A	1001	MYR	C9-C10-C11-C12
3	B	2001	MYR	C4-C5-C6-C7
3	A	1001	MYR	C5-C6-C7-C8
3	B	2001	MYR	C3-C4-C5-C6
3	A	1001	MYR	C2-C3-C4-C5
3	A	1001	MYR	C10-C11-C12-C13
3	B	2001	MYR	C11-C12-C13-C14
3	A	1001	MYR	C11-C12-C13-C14
3	B	2001	MYR	C2-C3-C4-C5

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1001	MYR	3	0
3	B	2001	MYR	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	489/541 (90%)	0.09	30 (6%) 21 22	10, 21, 70, 100	0
1	B	510/541 (94%)	0.06	33 (6%) 18 19	11, 21, 71, 102	0
All	All	999/1082 (92%)	0.07	63 (6%) 20 21	10, 21, 70, 102	0

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	493	LEU	8.5
1	A	481	VAL	8.2
1	B	514	ALA	5.9
1	B	541	ALA	5.6
1	B	490	PRO	5.2
1	A	526	LEU	5.1
1	A	478	LEU	5.0
1	B	515	GLU	4.9
1	A	502	PHE	4.7
1	B	463	ALA	4.6
1	B	497	LEU	4.6
1	B	516	GLU	4.6
1	B	517	ILE	4.2
1	B	528	ARG	4.1
1	A	531	ARG	4.1
1	A	468	ILE	3.9
1	B	533	GLN	3.9
1	A	501	GLY	3.9
1	B	8	ALA	3.7
1	B	480	VAL	3.7
1	B	464	ALA	3.7
1	A	528	ARG	3.6
1	A	464	ALA	3.5
1	B	489	THR	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	512	VAL	3.4
1	A	510	ALA	3.3
1	A	511	TYR	3.3
1	B	452	ASN	3.3
1	A	507	LEU	3.2
1	A	480	VAL	3.2
1	B	513	PHE	3.2
1	A	8	ALA	3.1
1	B	492	GLU	3.0
1	B	534	TYR	3.0
1	B	495	GLU	2.9
1	A	529	ALA	2.8
1	A	479	ALA	2.8
1	A	539	GLY	2.8
1	B	519	ARG	2.7
1	A	533	GLN	2.7
1	A	530	LEU	2.6
1	B	371	GLU	2.6
1	A	532	GLU	2.6
1	B	498	LEU	2.6
1	B	518	PRO	2.5
1	B	504	LYS	2.5
1	A	527	LYS	2.4
1	A	534	TYR	2.4
1	A	155	GLU	2.3
1	B	510	ALA	2.3
1	A	509	ASP	2.3
1	B	502	PHE	2.3
1	B	372	GLU	2.3
1	B	348	GLU	2.2
1	B	479	ALA	2.2
1	B	494	ASN	2.2
1	B	450	LEU	2.2
1	A	449	ASP	2.1
1	A	448	VAL	2.1
1	A	504	LYS	2.1
1	A	452	ASN	2.1
1	A	402	GLU	2.1
1	A	465	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	MYR	A	1001	15/16	0.87	0.19	6,7,9,9	0
3	MYR	B	2001	15/16	0.88	0.26	17,18,20,20	0
4	AMP	A	1002	23/23	0.95	0.11	7,13,13,14	0
2	MG	A	3001	1/1	0.96	0.12	35,35,35,35	0
4	AMP	B	2002	23/23	0.96	0.12	16,17,19,20	0
2	MG	B	3002	1/1	0.97	0.08	31,31,31,31	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around AMP A 1002:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around AMP B 2002:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.