wwPDB NMR Structure Validation Summary Report (i)

Feb 22, 2022 – 10:37 AM EST

PDB ID	:	1VAZ
Title	:	Solution structures of the p47 SEP domain
Authors	:	Yuan, X.; Simpson, P.; Mckeown, C.; Kondo, H.; Uchiyama, K.; Wallis, R.; Dreveny, I.; Keetch, C.; Zhang, X.; Robinson, C.; Freemont, P.; Matthews, S.
Deposited on	:	2004-02-20

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	2.26
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.26

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	88	

2 Ensemble composition and analysis (i)

This entry contains 10 models. Model 4 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:10-A:73 (64)	0.50	4

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 4, 7, 8, 9, 10
2	5, 6
3	2, 3

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1219 atoms, of which 596 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called NSFL1 cofactor p47.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	76	1219	381	596	122	119	1	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	expression tag	UNP O35987
A	-10	ARG	-	expression tag	UNP O35987
A	-9	GLY	-	expression tag	UNP O35987
A	-8	SER	-	expression tag	UNP O35987
A	-7	HIS	-	expression tag	UNP O35987
A	-6	HIS	-	expression tag	UNP O35987
A	-5	HIS	-	expression tag	UNP O35987
A	-4	HIS	-	expression tag	UNP O35987
A	-3	HIS	-	expression tag	UNP O35987
A	-2	HIS	-	expression tag	UNP O35987
A	-1	GLY	-	expression tag	UNP O35987
A	0	SER	-	expression tag	UNP O35987

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: NSFL1 cofactor p47

Chain A: • 40% 27% • 14% 14%

MET ARG GLY SER HIS HIS HIS HIS HIS HIS HIS GLY SER E1 R2 R3 R4 H5 S6 G7 Q8 D9 V10 H11 V12 V13 L14 K15 L16 W17 K18 T19 G20 F21 S22 L23 D24 N25 G26 D27 L28 R29 S30 Y31 Q32 D33 P34 S35 N36 A37 Q38 F39 L40 E41 S42 I43 R44 R45 G46 E47 V48

P49 A50 E51 L52 R53 R54 L55 A56 H57 G58 G59 Q60 V61 N62 L63 D64 M65 E66 D67 H68 R69 D70 E71 D72 F73 V74 K75 P76

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 4. Colouring as in section 4.1 above.

- Molecule 1: NSFL1 cofactor p47

Chain A: 7% 41% 22% • 14% 14%

MET ARG GLY SER HIS HIS HIS HIS HIS HIS HIS GLY SER E1 R2 R3 R4 H5 S6 G7 Q8 D9 V10 H11 V12 V13 L14 K15 L16 W17 K18 T19 G20 F21 S22 L23 D24 N25 L28 R29 S30 Y31 Q32 D33 P34 S35 N36 A37 Q38 F39 L40 E41 S42 I43 R44 R45 G46 E47 V48 P49

A50 E51 L52 R53 R54 L55 A56 H57 G58 G59 Q60 V61 N62 L63 D64 M65 E66 D67 H68 R69 D70 E71 D72 F73 V74 K75 P76

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *torsion angle dynamics, simulated annealing.*

Of the 100 calculated structures, 10 were deposited, based on the following criterion: *structures with acceptable covalent geometry, structures with favorable non-bond energy, structures with the least restraint violations, structures with the lowest energy.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	3.851

No chemical shift data was provided.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	5.8±0.4
All	All	0	58

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

5 of 6 unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	29	ARG	Sidechain	10
1	A	44	ARG	Sidechain	10
1	A	54	ARG	Sidechain	10
1	A	69	ARG	Sidechain	10
1	A	45	ARG	Sidechain	9

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	520	495	495	189±8
All	All	5200	4950	4950	1886

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 186.

5 of 960 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:52:LEU:HD13	1:A:53:ARG:N	1.18	1.53	7	1
1:A:14:LEU:HD13	1:A:63:LEU:HD13	1.09	1.23	10	2
1:A:14:LEU:HD22	1:A:43:ILE:HD11	1.02	1.29	9	2
1:A:52:LEU:HA	1:A:61:VAL:HG21	0.99	1.30	6	10
1:A:23:LEU:HD12	1:A:39:PHE:CE2	0.97	1.94	5	4

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	64/88 (73%)	44±1 (69±2%)	16±1 (25±2%)	4±1 (7±2%)	2 18
All	All	640/880 (73%)	439 (69%)	159 (25%)	42 (7%)	2 18

5 of 16 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	33	ASP	9
1	A	26	GLY	6
1	A	59	GLY	5
1	A	49	PRO	5
1	A	34	PRO	3

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	56/77 (73%)	31±3 (56±5%)	25±3 (44±5%)	0 3
All	All	560/770 (73%)	313 (56%)	247 (44%)	0 3

5 of 47 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	23	LEU	10
1	A	33	ASP	10
1	A	69	ARG	10
1	A	30	SER	9
1	A	52	LEU	9

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

No chemical shift data were provided