Full wwPDB X-ray Structure Validation Report (i)

Oct 5, 2023 – 05:22 AM EDT

PDB ID	:	6VC2
Title	:	LRH-1 bound to SS-RJW100 and a fragment of the Tif2 Coactivator
Authors	:	Mays, S.G.; Ortlund, E.A.
Deposited on	:	2019-12-20
Resolution	:	1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	FAILED
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	FAILED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 2207 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nuclear receptor subfamily 5 group A member 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	236	1941	1247	325	356	13	0	3	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	297	SER	-	expression tag	UNP O00482
A	298	ASN	-	expression tag	UNP O00482

- Molecule 2 is a protein called Nuclear receptor coactivator 2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	12	101	64	17	20	0	0	0

- Molecule 3 is (1S,3aS,6aS)-5-hexyl-4-phenyl-3a-(1-phenylethenyl)-1,2,3,3a,6,6a-hexahydropentalen-1-ol (three-letter code: QU7) (formula: $C_{28}H_{34}O$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 58 56 2	0	1

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	102	Total O 102 102	0	0
4	C	5	Total O 5 5	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.

3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α, β, γ	46.61Å 46.61Å 220.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.38 – 1.70	Depositor
% Data completeness (in resolution range)	97.4 (39.38-1.70)	Depositor
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.99 (at 1.70Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.210 , 0.228	Depositor
Wilson B-factor (Å^2)	28.1	Xtriage
Anisotropy	0.238	Xtriage
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2207	wwPDB-VP
Average B, all atoms (Å^2)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.17% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	QU7	A	601[B]	-	31,32,32	1.89	6 (19%)	29,45,45	1.99	10 (34%)
3	QU7	A	601[A]	-	31,32,32	1.94	6 (19%)	29,45,45	1.93	9 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	QU7	A	601[B]	-	-	4/19/50/50	0/4/4/4
3	QU7	A	601[A]	-	-	2/19/50/50	0/4/4/4

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601[A]	QU7	C08-C07	6.05	1.54	1.49
3	A	601[B]	QU7	C08-C07	5.74	1.53	1.49
3	A	601[A]	QU7	C06-C15	4.41	1.60	1.53
3	A	601[A]	QU7	C16-C15	3.78	1.55	1.48
3	A	601[B]	QU7	C01-C02	3.53	1.58	1.53
3	A	601[B]	QU7	C06-C15	3.46	1.59	1.53
3	A	601[B]	QU7	C16-C15	3.20	1.54	1.48
3	A	601[A]	QU7	C06-C01	2.89	1.60	1.56
3	A	601[B]	QU7	C22-C15	2.88	1.38	1.34
3	A	601[A]	QU7	C22-C15	2.71	1.38	1.34
3	A	601[B]	QU7	C06-C01	2.49	1.59	1.56
3	A	601[A]	QU7	C01-C02	2.24	1.56	1.53

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	A	601[B]	QU7	C06-C07-C08	5.33	126.93	116.83
3	A	601[A]	QU7	C06-C07-C08	5.03	126.35	116.83
3	A	601[B]	QU7	C20-C21-C16	3.54	124.54	120.34
3	A	601[A]	QU7	C20-C21-C16	3.54	124.53	120.34
3	A	601[A]	QU7	C21-C16-C17	-3.44	113.68	118.59
3	A	601[B]	QU7	C17-C16-C15	3.36	125.67	120.89

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	A	601[B]	QU7	C21-C16-C17	-3.32	113.86	118.59
3	A	601[A]	QU7	C17-C16-C15	3.31	125.59	120.89
3	A	601[B]	QU7	C24-C23-C22	3.02	120.21	113.74
3	A	601[A]	QU7	C24-C23-C22	2.73	119.58	113.74
3	A	601[B]	QU7	C13-C08-C09	-2.69	114.76	118.59
3	A	601[A]	QU7	C13-C08-C09	-2.66	114.80	118.59
3	A	601[A]	QU7	C18-C17-C16	2.63	123.45	120.34
3	A	601[B]	QU7	C10-C09-C08	2.59	123.41	120.34
3	A	601[A]	QU7	C10-C09-C08	2.55	123.36	120.34
3	A	601[B]	QU7	C18-C17-C16	2.50	123.30	120.34
3	A	601[A]	QU7	C12-C13-C08	2.46	123.26	120.34
3	A	601[B]	QU7	C12-C13-C08	2.45	123.24	120.34
3	A	601[B]	QU7	C29-C01-C06	-2.12	101.57	104.26

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	601[B]	QU7	C29-C22-C23-C24
3	A	601[A]	QU7	C24-C25-C26-C27
3	A	601[B]	QU7	C23-C24-C25-C26
3	A	601[B]	QU7	C25-C26-C27-C28
3	A	601[B]	QU7	C24-C25-C26-C27
3	A	601[A]	QU7	C25-C26-C27-C28

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand QU7 A 601 (B)



Bond lengths



Bond angles



Torsions



Rings

Ligand QU7 A 601 (A)



4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.