Full wwPDB X-ray Structure Validation Report (i)

May 14, 2020 – 02:50 am BST

PDB ID	:	3WIZ
Title	:	Crystal structure of Bcl-xL in complex with compound 10
Authors	:	Sogabe, S.; Igaki, S.; Hayano, Y.
Deposited on	:	2013-09-26
Resolution	:	2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1544 (2.48-2.44)
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)
RSRZ outliers	127900	1523 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	177	5% 69% 10% •• 20%
1	B	177	5% 71% 7% • 20%

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 2453 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bcl-2-like protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	141	1150	736	192	219	3	0	0	0
1	B	141	1150	736	192	219	3	0	0	0

- Molecule 2 is 7-(4-{[(4-{[(2R)-4-(dimethylamino)-1-(phenylsulfanyl)butan-2-yl]amino}-3-nitrophenyl)sulfonyl]carbamoyl}-2-methylphenyl)-3-[3-(naphthalen-1-yloxy)propyl]pyrazolo[1,5-a]pyridine-2-carboxylic acid (three-letter code: LC6) (formula: $C_{47}H_{46}N_6O_8S_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	A	1	63	47	6	8	2	0	0
2	B	1	63	47	6	8	2	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O P 5 4 1	0	0
3	B	1	Total O P 5 4 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	8	Total O 8 8	0	0
4	B	9	Total O 9 9	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bcl-2-like protein 1

Chain A:



- Molecule 1: Bcl-2-like protein 1

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 21 3	Depositor
Cell constants a, b, c, α, β, γ	152.66Å 152.66Å 152.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.45 48.28 – 2.45	Depositor EDS
% Data completeness (in resolution range)	99.7 (40.00-2.45) 99.7 (48.28-2.45)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$< I/\sigma(I) >$ [1]	2.08 (at 2.45Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.196 , 0.238 0.195 , 0.236	Depositor DCC
R_{free} test set	1117 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å^2)	56.8	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.36 , 46.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.52$, $< L^2 > = 0.35$	Xtriage
Estimated twinning fraction	0.480 for -l,-k,-h	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	2453	wwPDB-VP
Average B, all atoms (Å^2)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.23% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: LC6, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.45	0/1178	0.57	1/1594 (0.1%)
1	B	0.43	0/1178	0.56	0/1594
All	All	0.44	0/2356	0.57	1/3188 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	24	TRP	N-CA-C	-5.30	96.68	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1150	0	1084	25	0
1	B	1150	0	1084	19	0
2	A	63	0	45	2	0
2	B	63	0	45	1	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	8	0	0	0	0
4	B	9	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	2453	0	2258	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:159:MET:SD	1:B:159:MET:SD	2.52	1.07
1:A:159:MET:HA	1:A:159:MET:CE	2.06	0.83
1:A:159:MET:HE3	1:A:159:MET:HA	1.61	0.81
1:A:102:ARG:HA	1:A:102:ARG:HE	1.49	0.77
1:A:159:MET:HE1	1:B:155:VAL:HG22	1.69	0.74
1:B:159:MET:HA	1:B:159:MET:CE	2.20	0.72
1:A:23:SER:N	1:B:23:SER:O	2.26	0.68
1:A:24:TRP:O	1:A:25:SER:OG	2.11	0.68
1:A:159:MET:CE	1:B:155:VAL:HG22	2.23	0.67
1:A:102:ARG:HA	1:A:102:ARG:NE	2.14	0.61
1:B:159:MET:CE	1:B:162:LEU:HD23	2.31	0.61
1:B:102:ARG:O	1:B:106:SER:HB2	2.02	0.60
1:A:155:VAL:HA	1:B:159:MET:HE1	1.85	0.57
1:A:23:SER:O	1:B:23:SER:N	2.32	0.56
1:A:121:GLN:NE2	1:A:121:GLN:H	2.03	0.56
1:A:24:TRP:CE2	1:B:159:MET:HB3	2.41	0.55
1:A:159:MET:HB3	1:B:24:TRP:CE2	2.43	0.54
1:B:159:MET:HE1	1:B:162:LEU:HD23	1.90	0.54
1:A:98:GLU:O	1:A:102:ARG:HB2	2.08	0.53
1:A:159:MET:HG3	1:B:24:TRP:CD1	2.44	0.52
1:A:102:ARG:O	1:A:106:SER:HB2	2.10	0.52
1:A:155:VAL:HG22	1:B:159:MET:CE	2.41	0.51
1:A:159:MET:HE1	1:A:162:LEU:HD23	1.92	0.50
1:B:159:MET:HE3	1:B:162:LEU:HD23	1.96	0.47
1:A:24:TRP:NE1	1:B:159:MET:HB3	2.29	0.47
1:A:159:MET:HE3	1:A:162:LEU:HB3	1.96	0.47
1:B:102:ARG:HA	1:B:102:ARG:HH11	1.81	0.45
1:A:155:VAL:HG22	1:B:159:MET:HE2	1.98	0.45
2:A:501:LC6:H12	2:A:501:LC6:H41	1.99	0.45
1:A:24:TRP:O	1:A:25:SER:CB	2.65	0.43
1:A:23:SER:O	1:B:22:TYR:HA	2.19	0.42
1:A:16:LYS:NZ	1:A:98:GLU:OE2	2.49	0.42
2:A:501:LC6:N49	2:A:501:LC6:H22	2.36	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:501:LC6:H12	2:B:501:LC6:H41	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	137/177 (77%)	133 (97%)	4 (3%)	0	100	100
1	B	137/177 (77%)	131 (96%)	6 (4%)	0	100	100
All	All	274/354 (77%)	264 (96%)	10 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	121/151 (80%)	111 (92%)	10 (8%)	11	12
1	B	121/151 (80%)	110 (91%)	11 (9%)	9	10
All	All	242/302 (80%)	221 (91%)	21 (9%)	10	11

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	ARG
1	A	7	GLU
1	A	20	LYS
1	A	24	TRP
1	A	42	GLU
1	A	102	ARG
1	A	105	PHE
1	A	111	GLN
1	A	113	HIS
1	A	114	ILE
1	B	6	ARG
1	B	14	SER
1	B	24	TRP
1	B	88	GLN
1	B	102	ARG
1	B	105	PHE
1	B	111	GLN
1	B	113	HIS
1	B	114	ILE
1	B	132	ARG
1	B	159	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	111	GLN
1	A	121	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	PO4	B	502	-	4,4,4	0.82	0	6,6,6	0.45	0
3	PO4	A	502	-	4,4,4	0.84	0	6,6,6	0.40	0
2	LC6	A	501	-	61,69,69	1.25	8 (13%)	73,98,98	1.41	10 (13%)
2	LC6	B	501	-	61,69,69	1.26	10 (16%)	73,98,98	1.45	14 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LC6	A	501	-	-	10/42/48/48	0/7/7/7
2	LC6	B	501	-	-	13/42/48/48	0/7/7/7

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	LC6	S63-N51	3.45	1.71	1.64
2	B	501	LC6	S63-N51	3.26	1.71	1.64
2	B	501	LC6	C33-C34	2.85	1.44	1.38
2	A	501	LC6	C34-C35	-2.77	1.34	1.38
2	B	501	LC6	C36-N51	2.75	1.42	1.39
2	A	501	LC6	C33-C34	2.72	1.44	1.38
2	B	501	LC6	C34-C35	-2.72	1.34	1.38
2	A	501	LC6	C41-C24	-2.59	1.47	1.52
2	A	501	LC6	C36-N51	2.49	1.42	1.39
2	A	501	LC6	C20-C19	-2.44	1.38	1.43
2	B	501	LC6	C20-C19	-2.38	1.38	1.43
2	B	501	LC6	C41-C24	-2.37	1.48	1.52
2	B	501	LC6	C30-N49	-2.32	1.37	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	LC6	C30-N49	-2.26	1.37	1.40
2	B	501	LC6	C21-C35	-2.18	1.46	1.48
2	A	501	LC6	C21-C35	-2.09	1.46	1.48
2	B	501	LC6	O58-S63	2.04	1.45	1.43
2	B	501	LC6	C31-N48	-2.00	1.33	1.35

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	501	LC6	C43-C47-N50	-4.71	103.88	110.53
2	A	501	LC6	C25-N50-C47	-4.50	116.96	124.69
2	A	501	LC6	C43-C47-N50	-4.07	104.80	110.53
2	B	501	LC6	C25-N50-C47	-3.89	118.00	124.69
2	A	501	LC6	C45-O61-C27	-3.86	106.19	117.74
2	B	501	LC6	C45-O61-C27	-3.75	106.52	117.74
2	B	501	LC6	C26-C25-N50	-3.30	120.03	123.42
2	B	501	LC6	C36-N51-S63	-3.24	119.17	123.36
2	A	501	LC6	C26-C25-N50	-3.09	120.25	123.42
2	A	501	LC6	O59-S63-O58	2.97	123.20	119.55
2	A	501	LC6	C36-N51-S63	-2.94	119.57	123.36
2	B	501	LC6	O61-C27-C13	-2.85	118.18	124.46
2	B	501	LC6	O59-S63-O58	2.74	122.92	119.55
2	B	501	LC6	C22-C17-C23	-2.64	118.32	121.70
2	A	501	LC6	C22-C17-C23	-2.48	118.53	121.70
2	B	501	LC6	O55-C36-N51	2.45	124.09	121.08
2	A	501	LC6	O61-C27-C13	-2.41	119.14	124.46
2	A	501	LC6	O55-C36-N51	2.32	123.93	121.08
2	B	501	LC6	C43-C44-N52	-2.30	106.79	113.83
2	B	501	LC6	O59-S63-C29	-2.26	105.19	107.97
2	B	501	LC6	C29-S63-N51	-2.22	102.61	105.97
2	A	501	LC6	C43-C44-N52	-2.18	107.14	113.83
2	B	501	LC6	O61-C27-C20	2.16	121.04	115.01
2	B	501	LC6	C11-C22-C17	2.03	121.64	119.24

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	501	LC6	C30-C24-C41-C42
2	B	501	LC6	C31-C24-C41-C42
2	B	501	LC6	C24-C41-C42-C45
2	A	501	LC6	C13-C27-O61-C45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	501	LC6	C20-C27-O61-C45
2	A	501	LC6	C43-C44-N52-C39
2	A	501	LC6	C43-C44-N52-C40
2	B	501	LC6	C43-C44-N52-C39
2	B	501	LC6	C43-C44-N52-C40
2	B	501	LC6	C13-C27-O61-C45
2	B	501	LC6	C20-C27-O61-C45
2	B	501	LC6	C41-C42-C45-O61
2	B	501	LC6	C42-C45-O61-C27
2	B	501	LC6	C44-C43-C47-N50
2	A	501	LC6	C44-C43-C47-N50
2	A	501	LC6	C42-C45-O61-C27
2	B	501	LC6	C15-C28-S62-C46
2	A	501	LC6	C15-C28-S62-C46
2	A	501	LC6	C44-C43-C47-C46
2	B	501	LC6	C44-C43-C47-C46
2	A	501	LC6	C14-C28-S62-C46
2	B	501	LC6	C14-C28-S62-C46
2	A	501	LC6	C24-C41-C42-C45

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	LC6	2	0
2	B	501	LC6	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand LC6 A 501



Ligand LC6 B 501



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	141/177 (79%)	0.80	8 (5%)	23	20	40, 58, 109, 152	0
1	B	141/177 (79%)	0.76	9 (6%)	19	16	40, 59, 110, 161	0
All	All	282/354 (79%)	0.78	17 (6%)	21	18	40, 58, 112, 161	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	108	LEU	7.3
1	B	105	PHE	5.5
1	A	109	THR	5.3
1	B	108	LEU	5.3
1	A	105	PHE	4.0
1	B	24	TRP	3.8
1	B	104	ALA	2.7
1	A	114	ILE	2.6
1	A	110	SER	2.6
1	B	114	ILE	2.4
1	B	109	THR	2.3
1	B	150	LEU	2.3
1	A	22	TYR	2.2
1	A	24	TRP	2.2
1	A	162	LEU	2.2
1	B	103	ARG	2.1
1	B	162	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	LC6	A	501	63/63	0.95	0.29	44,74,187,229	0
2	LC6	B	501	63/63	0.95	0.26	48,70,191,223	0
3	PO4	B	502	5/5	0.97	0.16	64,89,96,96	5
3	PO4	A	502	5/5	0.98	0.15	60,94,97,97	5

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around LC6 A 501:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)



Electron density around LC6 B 501:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.