wwPDB X-ray Structure Validation Summary Report (i)

Nov 13, 2023 – 12:59 PM JST

PDB ID	:	5X66
Title	:	Human thymidylate synthase in complex with dUMP and methotrexate
Authors	:	Chen, D.; Jansson, A.; Larsson, A.; Nordlund, P.
Deposited on	:	2017-02-21
Resolution	:	1.99 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	290	93% 7% •
1	B	290	.% 92% 7% ••
1	C	290	2% 91% 7% ••
1	D	290	2% 91% 8% •
1	E	290	3% 90% 9% •
1	F	290	7% 91% 8% ••

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 15109 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thymidylate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	288	2314	1480	403	418	13	0	0	0
1	B	288	2310	1477	402	418	13	0	0	0
1	C	288	2309	1478	400	418	13	0	0	0
1	D	288	2312	1479	402	418	13	0	0	0
1	E	287	2300	1471	399	417	13	0	0	0
1	F	288	2312	1479	402	418	13	0	1	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	24	SER	-	expression tag	UNP P04818
A	25	MET	-	expression tag	UNP P04818
B	24	SER	-	expression tag	UNP P04818
B	25	MET	-	expression tag	UNP P04818
C	24	SER	-	expression tag	UNP P04818
C	25	MET	-	expression tag	UNP P04818
D	24	SER	-	expression tag	UNP P04818
D	25	MET	-	expression tag	UNP P04818
E	24	SER	-	expression tag	UNP P04818
E	25	MET	-	expression tag	UNP P04818
F	24	SER	-	expression tag	UNP P04818
F	25	MET	-	expression tag	UNP P04818

- Molecule 2 is 2'-DEOXYURIDINE 5'-MONOPHOSPHATE (three-letter code: UMP) (formula: $C_9H_{13}N_2O_8P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	B	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	C	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	D	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	E	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	F	1	Total	C	N	O	P	0	0
			20	9	2	8	1		

- Molecule 3 is METHOTREXATE (three-letter code: MTX) (formula: $C_{20}H_{22}N_8O_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	33	20	8	5	0	0
3	B	1	33	20	8	5	0	0
3	C	1	33	20	8	5	0	0
3	D	1	33	20	8	5	0	0
3	E	1	33	20	8	5	0	0
3	F	1	33	20	8	5	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	200	200	200	0	0
4	B	162	162	162	0	0
4	C	131	131	131	0	0
4	D	179	179	179	0	0
4	E	150	150	150	0	0
4	F	112	112	112	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Thymidylate synthase

Chain A:

- Molecule 1: Thymidylate synthase

Chain B:





- Molecule 1: Thymidylate synthase

Chain C:



- Molecule 1: Thymidylate synthase

Chain D:





- Molecule 1: Thymidylate synthase

Chain E:



• Molecule 1: Thymidylate synthase

Chain F:





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α, β, γ	109.20Å 109.20Å 317.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.18 – 1.99 25.18 – 1.99	Depositor EDS
% Data completeness (in resolution range)	91.3 (25.18-1.99) 91.4 (25.18-1.99)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.30 (at 1.99Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
R, R_{free}	0.201 , 0.238 0.211 , 0.247	Depositor DCC
R_{free} test set	6034 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å^2)	24.0	Xtriage
Anisotropy	0.277	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.37 , 47.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.58$, $< L^2 > = 0.43$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	15109	wwPDB-VP
Average B, all atoms (Å^2)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 60.94 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.4125e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MTX, UMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.91	2/2374 (0.1%)	0.99	9/3213 (0.3%)
1	B	0.85	0/2370	0.98	14/3209 (0.4%)
1	C	0.87	1/2369 (0.0%)	0.93	6/3207 (0.2%)
1	D	0.90	0/2372	0.94	10/3211 (0.3%)
1	E	0.86	4/2360 (0.2%)	0.94	8/3195 (0.3%)
1	F	0.85	1/2372 (0.0%)	0.98	12/3212 (0.4%)
All	All	0.87	8/14217 (0.1%)	0.96	59/19247 (0.3%)

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	175	ARG	CD-NE	-6.59	1.35	1.46
1	E	146	TYR	C-O	-6.07	1.11	1.23
1	A	175	ARG	CD-NE	-5.90	1.36	1.46
1	E	70	GLU	CD-OE2	-5.65	1.19	1.25
1	F	195	CYS	CB-SG	-5.63	1.72	1.81

The worst 5 of 59 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	E	115	ARG	NE-CZ-NH2	-9.27	115.67	120.30
1	E	175	ARG	NE-CZ-NH2	-9.14	115.73	120.30
1	E	115	ARG	NE-CZ-NH1	9.03	124.81	120.30
1	A	115	ARG	NE-CZ-NH2	-8.35	116.12	120.30
1	C	175	ARG	NE-CZ-NH2	-8.35	116.13	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2314	0	2275	5	0
1	B	2310	0	2264	5	0
1	C	2309	0	2266	11	0
1	D	2312	0	2268	7	0
1	E	2300	0	2253	11	0
1	F	2312	0	2261	11	0
2	A	20	0	10	0	0
2	B	20	0	11	0	0
2	C	20	0	10	0	0
2	D	20	0	10	0	0
2	E	20	0	10	0	0
2	F	20	0	10	3	0
3	A	33	0	19	0	0
3	B	33	0	21	0	0
3	C	33	0	20	2	0
3	D	33	0	20	1	0
3	E	33	0	20	2	0
3	F	33	0	21	0	0
4	A	200	0	0	1	0
4	B	162	0	0	0	0
4	C	131	0	0	2	0
4	D	179	0	0	0	0
4	E	150	0	0	4	0
4	F	112	0	0	3	0
All	All	15109	0	13769	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 48 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:51:THR:HG21	1:E:312:ALA:HB1	1.57	0.87
1:C:176:ARG:NH1	2:F:401:UMP:OP1	2.13	0.80
1:C:196:HIS:HB3	1:C:212:LEU:HD11	1.64	0.78

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:80:PHE:CD2	3:D:402:MTX:HA	2.30	0.67
1:F:99:LYS:HE2	4:F:512:HOH:O	1.95	0.65

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	286/290 (99%)	277 (97%)	9 (3%)	0	100	100
1	B	286/290 (99%)	277 (97%)	8 (3%)	1 (0%)	41	37
1	C	286/290 (99%)	277 (97%)	9 (3%)	0	100	100
1	D	286/290 (99%)	277 (97%)	9 (3%)	0	100	100
1	E	285/290 (98%)	276 (97%)	8 (3%)	1 (0%)	34	30
1	F	287/290 (99%)	278 (97%)	9 (3%)	0	100	100
All	All	1716/1740 (99%)	1662 (97%)	52 (3%)	2 (0%)	51	49

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	134	VAL
1	B	134	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	248/253 (98%)	243 (98%)	5 (2%)	55	58
1	B	247/253 (98%)	244 (99%)	3 (1%)	71	76
1	C	247/253 (98%)	241 (98%)	6 (2%)	49	51
1	D	247/253 (98%)	243 (98%)	4 (2%)	62	67
1	E	246/253 (97%)	243 (99%)	3 (1%)	71	76
1	F	246/253 (97%)	242 (98%)	4 (2%)	62	67
All	All	1481/1518 (98%)	1456 (98%)	25 (2%)	60	65

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	176	ARG
1	D	313	VAL
1	F	289	ASP
1	D	289	ASP
1	E	147	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	UMP	A	401	-	21,21,21	1.94	7 (33%)	31,31,31	1.86	6 (19%)
3	MTX	C	402	-	35,35,35	2.29	12 (34%)	46,49,49	2.57	19 (41%)
3	MTX	D	402	-	35,35,35	2.29	8 (22%)	46,49,49	4.13	19 (41%)
2	UMP	E	401	-	21,21,21	2.25	6 (28%)	31,31,31	2.41	13 (41%)
2	UMP	B	401	-	21,21,21	2.03	7 (33%)	31,31,31	1.68	6 (19%)
3	MTX	E	402	-	35,35,35	2.34	11 (31%)	46,49,49	2.54	17 (36%)
3	MTX	A	402	-	35,35,35	2.42	9 (25%)	46,49,49	2.11	10 (21%)
2	UMP	C	401	-	21,21,21	2.20	6 (28%)	31,31,31	2.11	4 (12%)
2	UMP	D	401	-	21,21,21	2.37	7 (33%)	31,31,31	2.42	11 (35%)
3	MTX	B	402	-	35,35,35	2.45	10 (28%)	46,49,49	1.93	13 (28%)
2	UMP	F	401	-	21,21,21	2.67	9 (42%)	31,31,31	2.20	8 (25%)
3	MTX	F	402	-	35,35,35	2.52	13 (37%)	46,49,49	2.02	14 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UMP	A	401	-	-	1/10/22/22	0/2/2/2
3	MTX	C	402	-	-	3/25/25/25	0/3/3/3
3	MTX	D	402	-	-	12/25/25/25	0/3/3/3
2	UMP	E	401	-	-	5/10/22/22	0/2/2/2
2	UMP	B	401	-	-	1/10/22/22	0/2/2/2
3	MTX	E	402	-	-	5/25/25/25	0/3/3/3
3	MTX	A	402	-	-	7/25/25/25	0/3/3/3
2	UMP	C	401	-	-	1/10/22/22	0/2/2/2
2	UMP	D	401	-	-	1/10/22/22	0/2/2/2
3	MTX	B	402	-	-	8/25/25/25	0/3/3/3
2	UMP	F	401	-	-	2/10/22/22	0/2/2/2
3	MTX	F	402	-	-	6/25/25/25	0/3/3/3

The worst 5 of 105 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	402	MTX	C-N	10.26	1.56	1.34
3	F	402	MTX	C-N	9.24	1.54	1.34
3	D	402	MTX	C-N	8.71	1.53	1.34
3	E	402	MTX	C-N	8.45	1.52	1.34
3	A	402	MTX	C-N	8.42	1.52	1.34

The worst 5 of 140 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	D	402	MTX	C11-C-N	16.96	149.58	117.06
3	D	402	MTX	O-C-C11	-9.81	103.45	120.94
2	C	401	UMP	C2'-C1'-N1	9.22	135.00	113.77
3	C	402	MTX	C2-N1-C8A	8.98	125.61	115.36
3	D	402	MTX	O-C-N	-8.77	106.32	122.45

There are no chirality outliers.

5 of 52 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	402	MTX	CT-CA-N-C
3	C	402	MTX	CA-CB-CG-CD
3	D	402	MTX	O-C-N-CA
3	D	402	MTX	C11-C-N-CA
3	B	402	MTX	CB-CA-N-C

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	402	MTX	2	0
3	D	402	MTX	1	0
3	E	402	MTX	2	0
2	F	401	UMP	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand UMP A 401



Ligand MTX C 402



Ligand MTX D 402



Bond lengths



Bond angles



Torsions



Rings

Ligand UMP E 401



Bond lengths



Bond angles



Torsions



Rings

Ligand UMP B 401



Bond lengths



Bond angles



Torsions



Rings

Ligand MTX E 402



Bond lengths



Bond angles



Torsions



Rings

Ligand MTX A 402



Bond lengths



Bond angles



Torsions



Rings

Ligand UMP C 401



Bond lengths



Bond angles



Torsions



Rings

Ligand UMP D 401



Bond lengths



Bond angles



Torsions



Rings

Ligand MTX B 402



Bond lengths



Bond angles



Torsions



Rings

Ligand UMP F 401



Ligand MTX F 402



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	288/290 (99%)	-0.16	1 (0%)	94	93	12, 21, 35, 59	0
1	B	288/290 (99%)	-0.07	2 (0%)	87	87	15, 25, 38, 56	0
1	C	288/290 (99%)	0.22	7 (2%)	59	57	17, 28, 49, 62	0
1	D	288/290 (99%)	-0.02	6 (2%)	63	62	13, 22, 42, 68	0
1	E	287/290 (98%)	0.08	9 (3%)	49	48	17, 26, 47, 60	0
1	F	288/290 (99%)	0.35	19 (6%)	18	17	19, 30, 53, 69	0
All	All	1727/1740 (99%)	0.07	44 (2%)	57	56	12, 26, 45, 69	0

The worst 5 of 44 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	313	VAL	10.4
1	D	313	VAL	8.2
1	A	313	VAL	6.5
1	F	313	VAL	5.9
1	F	27	PRO	4.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	MTX	E	402	33/33	0.81	0.19	33,38,69,72	0
3	MTX	C	402	33/33	0.82	0.19	32,38,73,83	0
3	MTX	F	402	33/33	0.89	0.17	36,43,93,99	0
3	MTX	A	402	33/33	0.91	0.14	22,29,58,63	0
3	MTX	D	402	33/33	0.91	0.14	27,31,65,67	0
3	MTX	B	402	33/33	0.92	0.14	20,29,92,96	0
2	UMP	E	401	20/20	0.96	0.08	27,32,35,36	0
2	UMP	F	401	20/20	0.96	0.08	25,34,37,37	0
2	UMP	D	401	20/20	0.98	0.07	22,26,29,29	0
2	UMP	B	401	20/20	0.98	0.08	18,20,22,22	0
2	UMP	C	401	20/20	0.98	0.07	22,27,29,31	0
2	UMP	A	401	20/20	0.99	0.08	17,18,21,21	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around MTX E 402:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MTX C 402:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MTX F 402:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around MTX A 402:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around MTX D 402:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MTX B 402:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around UMP E 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around UMP F 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around UMP D 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around UMP B 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around UMP C 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around UMP A 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.