



wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 04:59 pm BST

PDB ID : 2X7J
Title : Structure of the menaquinone biosynthesis protein MenD from *Bacillus subtilis*
Authors : Dawson, A.; Chen, M.; Fyfe, P.K.; Guo, Z.; Hunter, W.N.
Deposited on : 2010-03-01
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	604	
1	B	604	
1	C	604	
1	D	604	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EDO	A	603	-	-	X	-
4	EDO	B	603	-	-	X	-
4	EDO	C	603	-	-	X	-
4	EDO	D	603	-	-	X	-
6	SO4	A	1582	-	-	X	-
6	SO4	B	1582	-	-	X	-
6	SO4	C	1580	-	-	X	-
6	SO4	D	1582	-	-	X	-

2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 19356 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2-SUCCINYL-5-ENOLPYRUVYL-6-HYDROXY-3-CYCLOHEXENE -1-CARBOXYLATE SYNTHASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	579	4544	2885	789	851	19	0	6	0
1	B	579	4520	2868	784	849	19	0	2	0
1	C	577	4502	2857	782	844	19	0	2	0
1	D	579	4548	2887	793	849	19	0	6	0

There are 96 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-23	MET	-	expression tag	UNP P23970
A	-22	GLY	-	expression tag	UNP P23970
A	-21	SER	-	expression tag	UNP P23970
A	-20	SER	-	expression tag	UNP P23970
A	-19	HIS	-	expression tag	UNP P23970
A	-18	HIS	-	expression tag	UNP P23970
A	-17	HIS	-	expression tag	UNP P23970
A	-16	HIS	-	expression tag	UNP P23970
A	-15	HIS	-	expression tag	UNP P23970
A	-14	HIS	-	expression tag	UNP P23970
A	-13	SER	-	expression tag	UNP P23970
A	-12	SER	-	expression tag	UNP P23970
A	-11	GLY	-	expression tag	UNP P23970
A	-10	GLU	-	expression tag	UNP P23970
A	-9	ASN	-	expression tag	UNP P23970
A	-8	LEU	-	expression tag	UNP P23970
A	-7	TYR	-	expression tag	UNP P23970
A	-6	PHE	-	expression tag	UNP P23970
A	-5	GLN	-	expression tag	UNP P23970
A	-4	GLY	-	expression tag	UNP P23970

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	HIS	-	expression tag	UNP P23970
A	-2	MET	-	expression tag	UNP P23970
A	-1	LEU	-	expression tag	UNP P23970
A	0	GLU	-	expression tag	UNP P23970
B	-23	MET	-	expression tag	UNP P23970
B	-22	GLY	-	expression tag	UNP P23970
B	-21	SER	-	expression tag	UNP P23970
B	-20	SER	-	expression tag	UNP P23970
B	-19	HIS	-	expression tag	UNP P23970
B	-18	HIS	-	expression tag	UNP P23970
B	-17	HIS	-	expression tag	UNP P23970
B	-16	HIS	-	expression tag	UNP P23970
B	-15	HIS	-	expression tag	UNP P23970
B	-14	HIS	-	expression tag	UNP P23970
B	-13	SER	-	expression tag	UNP P23970
B	-12	SER	-	expression tag	UNP P23970
B	-11	GLY	-	expression tag	UNP P23970
B	-10	GLU	-	expression tag	UNP P23970
B	-9	ASN	-	expression tag	UNP P23970
B	-8	LEU	-	expression tag	UNP P23970
B	-7	TYR	-	expression tag	UNP P23970
B	-6	PHE	-	expression tag	UNP P23970
B	-5	GLN	-	expression tag	UNP P23970
B	-4	GLY	-	expression tag	UNP P23970
B	-3	HIS	-	expression tag	UNP P23970
B	-2	MET	-	expression tag	UNP P23970
B	-1	LEU	-	expression tag	UNP P23970
B	0	GLU	-	expression tag	UNP P23970
C	-23	MET	-	expression tag	UNP P23970
C	-22	GLY	-	expression tag	UNP P23970
C	-21	SER	-	expression tag	UNP P23970
C	-20	SER	-	expression tag	UNP P23970
C	-19	HIS	-	expression tag	UNP P23970
C	-18	HIS	-	expression tag	UNP P23970
C	-17	HIS	-	expression tag	UNP P23970
C	-16	HIS	-	expression tag	UNP P23970
C	-15	HIS	-	expression tag	UNP P23970
C	-14	HIS	-	expression tag	UNP P23970
C	-13	SER	-	expression tag	UNP P23970
C	-12	SER	-	expression tag	UNP P23970
C	-11	GLY	-	expression tag	UNP P23970
C	-10	GLU	-	expression tag	UNP P23970

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-9	ASN	-	expression tag	UNP P23970
C	-8	LEU	-	expression tag	UNP P23970
C	-7	TYR	-	expression tag	UNP P23970
C	-6	PHE	-	expression tag	UNP P23970
C	-5	GLN	-	expression tag	UNP P23970
C	-4	GLY	-	expression tag	UNP P23970
C	-3	HIS	-	expression tag	UNP P23970
C	-2	MET	-	expression tag	UNP P23970
C	-1	LEU	-	expression tag	UNP P23970
C	0	GLU	-	expression tag	UNP P23970
D	-23	MET	-	expression tag	UNP P23970
D	-22	GLY	-	expression tag	UNP P23970
D	-21	SER	-	expression tag	UNP P23970
D	-20	SER	-	expression tag	UNP P23970
D	-19	HIS	-	expression tag	UNP P23970
D	-18	HIS	-	expression tag	UNP P23970
D	-17	HIS	-	expression tag	UNP P23970
D	-16	HIS	-	expression tag	UNP P23970
D	-15	HIS	-	expression tag	UNP P23970
D	-14	HIS	-	expression tag	UNP P23970
D	-13	SER	-	expression tag	UNP P23970
D	-12	SER	-	expression tag	UNP P23970
D	-11	GLY	-	expression tag	UNP P23970
D	-10	GLU	-	expression tag	UNP P23970
D	-9	ASN	-	expression tag	UNP P23970
D	-8	LEU	-	expression tag	UNP P23970
D	-7	TYR	-	expression tag	UNP P23970
D	-6	PHE	-	expression tag	UNP P23970
D	-5	GLN	-	expression tag	UNP P23970
D	-4	GLY	-	expression tag	UNP P23970
D	-3	HIS	-	expression tag	UNP P23970
D	-2	MET	-	expression tag	UNP P23970
D	-1	LEU	-	expression tag	UNP P23970
D	0	GLU	-	expression tag	UNP P23970

- Molecule 2 is THIAMINE DIPHOSPHATE (three-letter code: TPP) (formula: C₁₂H₁₉N₄O₇P₂S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			S
2	A	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		
2	B	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		
2	C	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		
2	D	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mn		
3	B	1	Total	Mn	0	0
			1	1		
3	A	1	Total	Mn	0	0
			1	1		
3	D	1	Total	Mn	0	0
			1	1		
3	C	1	Total	Mn	0	0
			1	1		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Na 1 1	0	0
5	A	1	Total Na 1 1	0	0
5	D	1	Total Na 1 1	0	0
5	C	1	Total Na 1 1	0	0

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	B	1	Total	O	S	0	0
			5	4	1		
6	B	1	Total	O	S	0	0
			5	4	1		
6	C	1	Total	O	S	0	0
			5	4	1		
6	C	1	Total	O	S	0	0
			5	4	1		
6	D	1	Total	O	S	0	0
			5	4	1		
6	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	285	Total	O	0	0
			285	285		
7	B	239	Total	O	0	0
			239	239		
7	C	256	Total	O	0	0
			256	256		
7	D	294	Total	O	0	0
			294	294		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 2-SUCCINYL-5-ENOLPYRUVYL-6-HYDROXY-3-CYCLOHEXENE -1-CARBOXYLATE SYNTHASE





• Molecule 1: 2-SUCCINYL-5-ENOLPYRUVYL-6-HYDROXY-3-CYCLOHEXENE -1-CARBOXYLATE SYNTHASE







● Molecule 1: 2-SUCCINYL-5-ENOLPYRUVYL-6-HYDROXY-3-CYCLOHEXENE -1-CARBOXYLATE SYNTHASE



● Molecule 1: 2-SUCCINYL-5-ENOLPYRUVYL-6-HYDROXY-3-CYCLOHEXENE -1-CARBOXYLATE SYNTHASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	100.18Å 152.99Å 158.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	62.14 – 2.35 62.15 – 2.35	Depositor EDS
% Data completeness (in resolution range)	100.0 (62.14-2.35) 100.0 (62.15-2.35)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.39 (at 2.34Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.172 , 0.228 0.173 , 0.227	Depositor DCC
R_{free} test set	5118 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	30.3	Xtrriage
Anisotropy	0.082	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 43.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.010 for -h,l,k	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	19356	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 26.04 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.8341e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, TPP, MN, EDO, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/4671	0.64	3/6342 (0.0%)
1	B	0.45	0/4635	0.61	3/6294 (0.0%)
1	C	0.45	0/4617	0.60	2/6271 (0.0%)
1	D	0.48	0/4675	0.64	3/6346 (0.0%)
All	All	0.46	0/18598	0.62	11/25253 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
1	C	0	2
1	D	0	2
All	All	0	8

There are no bond length outliers.

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	175	GLU	C-N-CD	-11.48	95.34	120.60
1	C	162	ARG	C-N-CD	-11.36	95.60	120.60
1	A	175	GLU	C-N-CD	-11.26	95.83	120.60
1	A	162	ARG	C-N-CD	-10.23	98.08	120.60
1	B	162	ARG	C-N-CD	-10.04	98.52	120.60

There are no chirality outliers.

5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	162	ARG	Peptide
1	A	175	GLU	Peptide
1	B	162	ARG	Peptide
1	B	175	GLU	Peptide
1	C	162	ARG	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4544	0	4503	74	0
1	B	4520	0	4465	91	0
1	C	4502	0	4448	82	0
1	D	4548	0	4513	111	0
2	A	26	0	16	2	0
2	B	26	0	16	0	0
2	C	26	0	16	1	0
2	D	26	0	16	2	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	4	0	6	13	0
4	B	4	0	6	6	0
4	C	4	0	6	8	0
4	D	4	0	6	8	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
6	A	10	0	0	5	0
6	B	10	0	0	6	0
6	C	10	0	0	6	0
6	D	10	0	0	9	0
7	A	285	0	0	11	0
7	B	239	0	0	9	0
7	C	256	0	0	10	0
7	D	294	0	0	17	0
All	All	19356	0	18017	359	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 359 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:31:SER:H	4:C:603:EDO:H11	1.04	1.20
1:D:31:SER:H	4:D:603:EDO:H22	1.00	1.17
1:D:163:PRO:HD3	1:D:319:ASP:HB3	1.34	1.08
1:C:206:ARG:HH11	1:C:206:ARG:HG3	0.89	1.06
1:B:163:PRO:CD	1:B:319:ASP:HB3	1.85	1.06

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	583/604 (96%)	555 (95%)	24 (4%)	4 (1%)	22	23
1	B	579/604 (96%)	555 (96%)	17 (3%)	7 (1%)	13	11
1	C	577/604 (96%)	552 (96%)	22 (4%)	3 (0%)	29	32
1	D	583/604 (96%)	564 (97%)	17 (3%)	2 (0%)	41	47
All	All	2322/2416 (96%)	2226 (96%)	80 (3%)	16 (1%)	22	23

5 of 16 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	163	PRO
1	A	176	PRO
1	A	498	GLU
1	B	163	PRO
1	B	176	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	495/511 (97%)	467 (94%)	28 (6%)	20	22
1	B	491/511 (96%)	466 (95%)	25 (5%)	24	27
1	C	489/511 (96%)	468 (96%)	21 (4%)	29	35
1	D	495/511 (97%)	473 (96%)	22 (4%)	28	34
All	All	1970/2044 (96%)	1874 (95%)	96 (5%)	27	29

5 of 96 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	367	ARG
1	C	86[A]	TYR
1	D	448	LYS
1	B	386	LEU
1	B	498	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	431	ASN
1	C	117	GLN
1	D	127	ASN
1	B	334	HIS
1	D	117	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 8 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	C	603	-	3,3,3	0.53	0	2,2,2	0.33	0
4	EDO	A	603	-	3,3,3	0.46	0	2,2,2	0.35	0
6	SO4	B	1583	-	4,4,4	0.14	0	6,6,6	0.16	0
6	SO4	A	1583	-	4,4,4	0.15	0	6,6,6	0.14	0
6	SO4	D	1583	-	4,4,4	0.17	0	6,6,6	0.27	0
4	EDO	D	603	-	3,3,3	0.55	0	2,2,2	0.28	0
2	TPP	B	601	3	22,27,27	1.70	4 (18%)	29,40,40	1.97	10 (34%)
6	SO4	C	1580	-	4,4,4	0.31	0	6,6,6	0.50	0
4	EDO	B	603	-	3,3,3	0.54	0	2,2,2	0.45	0
2	TPP	D	601	3	22,27,27	2.19	5 (22%)	29,40,40	2.05	11 (37%)
6	SO4	C	1581	-	4,4,4	0.13	0	6,6,6	0.17	0
2	TPP	C	601	3	22,27,27	1.93	6 (27%)	29,40,40	2.12	10 (34%)
6	SO4	B	1582	-	4,4,4	0.23	0	6,6,6	0.58	0
2	TPP	A	601	3	22,27,27	1.99	3 (13%)	29,40,40	2.23	12 (41%)
6	SO4	A	1582	-	4,4,4	0.28	0	6,6,6	0.20	0
6	SO4	D	1582	-	4,4,4	0.24	0	6,6,6	0.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	C	603	-	-	0/1/1/1	-
4	EDO	A	603	-	-	0/1/1/1	-
4	EDO	D	603	-	-	1/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	B	603	-	-	0/1/1/1	-
2	TPP	D	601	3	-	1/16/17/17	0/2/2/2
2	TPP	B	601	3	-	1/16/17/17	0/2/2/2
2	TPP	C	601	3	-	0/16/17/17	0/2/2/2
2	TPP	A	601	3	-	1/16/17/17	0/2/2/2

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	601	TPP	C6-C5	7.20	1.54	1.50
2	C	601	TPP	C6-C5	6.44	1.53	1.50
2	A	601	TPP	C6-C5	6.35	1.53	1.50
2	A	601	TPP	C4-N3	-4.21	1.36	1.39
2	D	601	TPP	PB-O1B	4.02	1.63	1.50

The worst 5 of 43 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	601	TPP	CM2-C2'-N1'	4.41	121.99	117.14
2	C	601	TPP	C5-C4-N3	4.18	115.94	107.57
2	A	601	TPP	C6-C5-C4	4.15	130.76	127.43
2	A	601	TPP	C6'-N1'-C2'	4.10	122.93	115.96
2	C	601	TPP	C6-C5-C4	3.87	130.54	127.43

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	601	TPP	PB-O3A-PA-O7
2	B	601	TPP	PB-O3A-PA-O7
2	A	601	TPP	PB-O3A-PA-O7
4	D	603	EDO	O1-C1-C2-O2

There are no ring outliers.

12 monomers are involved in 64 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	603	EDO	8	0
4	A	603	EDO	13	0
4	D	603	EDO	8	0
6	C	1580	SO4	5	0



Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	603	EDO	6	0
2	D	601	TPP	2	0
6	C	1581	SO4	1	0
2	C	601	TPP	1	0
6	B	1582	SO4	6	0
2	A	601	TPP	2	0
6	A	1582	SO4	5	0
6	D	1582	SO4	9	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	579/604 (95%)	-0.11	4 (0%) 87 92	15, 29, 46, 61	1 (0%)
1	B	579/604 (95%)	0.16	23 (3%) 38 51	16, 34, 57, 67	3 (0%)
1	C	577/604 (95%)	0.07	13 (2%) 60 70	15, 33, 55, 68	6 (1%)
1	D	579/604 (95%)	-0.21	7 (1%) 79 86	17, 28, 44, 60	2 (0%)
All	All	2314/2416 (95%)	-0.02	47 (2%) 65 75	15, 30, 52, 68	12 (0%)

The worst 5 of 47 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	2	THR	6.4
1	B	352	THR	4.9
1	C	351	ALA	4.6
1	D	2	THR	4.6
1	B	2	THR	4.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	EDO	D	603	4/4	0.90	0.20	26,27,29,29	0
5	NA	A	1581	1/1	0.91	0.10	35,35,35,35	0
4	EDO	C	603	4/4	0.93	0.18	27,27,27,28	0
6	SO4	B	1583	5/5	0.93	0.13	31,31,31,31	5
4	EDO	A	603	4/4	0.94	0.20	25,27,28,29	0
6	SO4	B	1582	5/5	0.94	0.22	13,14,15,15	5
4	EDO	B	603	4/4	0.94	0.15	23,24,25,25	0
6	SO4	C	1581	5/5	0.94	0.15	26,27,27,27	5
6	SO4	A	1582	5/5	0.96	0.18	10,10,11,11	5
5	NA	C	1579	1/1	0.96	0.10	36,36,36,36	0
5	NA	B	1581	1/1	0.96	0.10	33,33,33,33	0
5	NA	D	1581	1/1	0.96	0.14	33,33,33,33	0
6	SO4	D	1583	5/5	0.97	0.24	21,22,22,22	5
6	SO4	A	1583	5/5	0.97	0.14	24,24,24,24	5
6	SO4	D	1582	5/5	0.98	0.28	16,16,17,17	5
2	TPP	C	601	26/26	0.98	0.10	22,23,24,24	0
2	TPP	B	601	26/26	0.98	0.10	21,24,26,27	0
6	SO4	C	1580	5/5	0.98	0.20	10,12,12,12	5
2	TPP	A	601	26/26	0.98	0.10	20,23,25,25	0
3	MN	B	602	1/1	0.99	0.08	29,29,29,29	0
2	TPP	D	601	26/26	0.99	0.09	19,21,23,24	0
3	MN	A	602	1/1	0.99	0.07	24,24,24,24	0
3	MN	C	602	1/1	1.00	0.04	24,24,24,24	0
3	MN	D	602	1/1	1.00	0.09	23,23,23,23	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around TPP C 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around TPP B 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around TPP A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around TPP D 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers

There are no such residues in this entry.