



# wwPDB X-ray Structure Validation Summary Report

May 23, 2020 – 02:07 pm BST

PDB ID : 1XEB  
Title : Crystal Structure of an Acyl-CoA N-acyltransferase from *Pseudomonas aeruginosa*  
Authors : Bertero, M.G.; Walker, J.R.; Skarina, T.; Gorodichtchenskaia, E.; Joachimiak, A.; Edwards, A.E.; Savchenko, A.; Strynadka, N.; Midwest Center for Structural Genomics (MCSG)  
Deposited on : 2004-09-09  
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	150	
1	B	150	
1	C	150	
1	D	150	
1	E	150	
1	F	150	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	G	150	
1	H	150	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9702 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called hypothetical protein PA0115.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	149	1196	760	214	217	2	3	0	0	0
1	B	148	1192	758	213	216	2	3	0	0	0
1	C	149	1196	760	214	217	2	3	0	0	0
1	D	148	1190	757	213	215	2	3	0	0	0
1	E	146	1172	746	211	210	2	3	0	0	0
1	F	145	1173	748	209	211	2	3	0	0	0
1	G	147	1184	754	212	213	2	3	0	0	0
1	H	146	1177	750	210	212	2	3	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	53	MSE	MET	MODIFIED RESIDUE	UNP Q9I717
A	97	MSE	MET	MODIFIED RESIDUE	UNP Q9I717
A	147	MSE	MET	MODIFIED RESIDUE	UNP Q9I717
B	53	MSE	MET	MODIFIED RESIDUE	UNP Q9I717
B	97	MSE	MET	MODIFIED RESIDUE	UNP Q9I717
B	147	MSE	MET	MODIFIED RESIDUE	UNP Q9I717
C	53	MSE	MET	MODIFIED RESIDUE	UNP Q9I717
C	97	MSE	MET	MODIFIED RESIDUE	UNP Q9I717
C	147	MSE	MET	MODIFIED RESIDUE	UNP Q9I717
D	53	MSE	MET	MODIFIED RESIDUE	UNP Q9I717
D	97	MSE	MET	MODIFIED RESIDUE	UNP Q9I717
D	147	MSE	MET	MODIFIED RESIDUE	UNP Q9I717
E	53	MSE	MET	MODIFIED RESIDUE	UNP Q9I717

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
E	97	MSE	MET	MODIFIED RESIDUE	UNP Q9I717
E	147	MSE	MET	MODIFIED RESIDUE	UNP Q9I717
F	53	MSE	MET	MODIFIED RESIDUE	UNP Q9I717
F	97	MSE	MET	MODIFIED RESIDUE	UNP Q9I717
F	147	MSE	MET	MODIFIED RESIDUE	UNP Q9I717
G	53	MSE	MET	MODIFIED RESIDUE	UNP Q9I717
G	97	MSE	MET	MODIFIED RESIDUE	UNP Q9I717
G	147	MSE	MET	MODIFIED RESIDUE	UNP Q9I717
H	53	MSE	MET	MODIFIED RESIDUE	UNP Q9I717
H	97	MSE	MET	MODIFIED RESIDUE	UNP Q9I717
H	147	MSE	MET	MODIFIED RESIDUE	UNP Q9I717


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	47	Total O 47 47	0	0
2	B	37	Total O 37 37	0	0
2	C	39	Total O 39 39	0	0
2	D	17	Total O 17 17	0	0
2	E	27	Total O 27 27	0	0
2	F	34	Total O 34 34	0	0
2	G	14	Total O 14 14	0	0
2	H	7	Total O 7 7	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: hypothetical protein PA0115



- Molecule 1: hypothetical protein PA0115




- Molecule 1: hypothetical protein PA0115




- Molecule 1: hypothetical protein PA0115






• Molecule 1: hypothetical protein PA0115




• Molecule 1: hypothetical protein PA0115



• Molecule 1: hypothetical protein PA0115



• Molecule 1: hypothetical protein PA0115



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	138.40Å 138.40Å 136.49Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.89 – 2.35 29.88 – 2.35	Depositor EDS
% Data completeness (in resolution range)	91.9 (29.89-2.35) 92.0 (29.88-2.35)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	93.19 (at 2.36Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.232 , 0.282 0.231 , 0.281	Depositor DCC
$R_{free}$ test set	4765 reflections (8.20%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.6	Xtriage
Anisotropy	0.049	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 38.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.022 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	9702	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.14% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.41	0/1220	0.69	0/1651
1	B	0.43	0/1215	0.73	1/1643 (0.1%)
1	C	0.42	0/1220	0.73	1/1651 (0.1%)
1	D	0.38	0/1214	0.70	1/1643 (0.1%)
1	E	0.40	0/1195	0.69	0/1614
1	F	0.42	0/1196	0.69	1/1618 (0.1%)
1	G	0.36	0/1208	0.66	0/1636
1	H	0.36	0/1200	0.68	0/1623
All	All	0.40	0/9668	0.70	4/13079 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	64	LEU	CA-CB-CG	5.67	128.34	115.30
1	B	115	LEU	CA-CB-CG	5.48	127.90	115.30
1	D	64	LEU	CA-CB-CG	5.36	127.62	115.30
1	F	115	LEU	CA-CB-CG	5.22	127.32	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1196	0	1178	31	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1192	0	1174	32	0
1	C	1196	0	1178	45	0
1	D	1190	0	1173	41	0
1	E	1172	0	1159	42	0
1	F	1173	0	1158	47	0
1	G	1184	0	1168	59	0
1	H	1177	0	1161	57	0
2	A	47	0	0	2	0
2	B	37	0	0	0	0
2	C	39	0	0	3	0
2	D	17	0	0	1	0
2	E	27	0	0	0	0
2	F	34	0	0	4	0
2	G	14	0	0	2	0
2	H	7	0	0	0	0
All	All	9702	0	9349	337	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 337 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:16:LYS:HD2	1:H:16:LYS:H	1.00	1.13
1:F:64:LEU:HD13	1:F:78:ILE:HG23	1.47	0.97
1:D:64:LEU:HD21	1:D:100:ALA:HB2	1.49	0.94
1:A:64:LEU:HD21	1:A:100:ALA:HB2	1.50	0.92
1:E:64:LEU:HD21	1:E:100:ALA:HB2	1.49	0.91

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	147/150 (98%)	137 (93%)	10 (7%)	0	100	100
1	B	144/150 (96%)	135 (94%)	9 (6%)	0	100	100
1	C	147/150 (98%)	138 (94%)	9 (6%)	0	100	100
1	D	146/150 (97%)	131 (90%)	13 (9%)	2 (1%)	11	9
1	E	142/150 (95%)	130 (92%)	11 (8%)	1 (1%)	22	23
1	F	141/150 (94%)	130 (92%)	10 (7%)	1 (1%)	22	23
1	G	145/150 (97%)	130 (90%)	14 (10%)	1 (1%)	22	23
1	H	142/150 (95%)	129 (91%)	13 (9%)	0	100	100
All	All	1154/1200 (96%)	1060 (92%)	89 (8%)	5 (0%)	34	38

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	120	HIS
1	D	31	VAL
1	E	10	HIS
1	F	10	HIS
1	G	119	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	124/122 (102%)	113 (91%)	11 (9%)	9	8
1	B	124/122 (102%)	113 (91%)	11 (9%)	9	8
1	C	124/122 (102%)	112 (90%)	12 (10%)	8	7
1	D	123/122 (101%)	108 (88%)	15 (12%)	5	4
1	E	121/122 (99%)	111 (92%)	10 (8%)	11	11
1	F	122/122 (100%)	114 (93%)	8 (7%)	16	17
1	G	123/122 (101%)	114 (93%)	9 (7%)	14	14
1	H	122/122 (100%)	104 (85%)	18 (15%)	3	3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	983/976 (101%)	889 (90%)	94 (10%)	<b>8</b> <b>7</b>

5 of 94 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	75	GLN
1	E	60	LEU
1	H	85	SER
1	D	101	LEU
1	D	141	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	23	GLN
1	H	59	GLN
1	G	59	GLN
1	C	122	GLN
1	F	94	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	146/150 (97%)	-0.01	5 (3%) 45 57	18, 29, 46, 52	0
1	B	145/150 (96%)	-0.07	2 (1%) 75 83	17, 30, 47, 56	0
1	C	146/150 (97%)	-0.11	1 (0%) 87 92	15, 29, 47, 61	0
1	D	145/150 (96%)	-0.01	4 (2%) 53 64	22, 36, 50, 63	0
1	E	143/150 (95%)	0.35	9 (6%) 20 29	21, 39, 58, 73	0
1	F	142/150 (94%)	0.16	5 (3%) 44 56	21, 34, 56, 64	0
1	G	144/150 (96%)	0.54	11 (7%) 13 21	25, 48, 67, 79	0
1	H	143/150 (95%)	0.48	9 (6%) 20 29	30, 45, 65, 72	0
All	All	1154/1200 (96%)	0.16	46 (3%) 38 51	15, 36, 59, 79	0

The worst 5 of 46 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	150	ALA	5.7
1	E	2	SER	5.1
1	F	57	ASP	4.3
1	F	87	ALA	3.9
1	F	3	LEU	3.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.