wwPDB X-ray Structure Validation Summary Report (i)

Aug 15, 2023 – 03:48 PM EDT

PDB ID	:	1XL8
Title	:	Crystal structure of mouse carnitine octanoyltransferase in complex with octanoylcarnitine
Authors	:	Jogl, G.; Hsiao, Y.S.; Tong, L.
Deposited on	:	2004-09-30
Resolution	:	2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	612	6% 69% 26%
1	B	612	% 77% 20%

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 10118 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peroxisomal carnitine O-octanoyltransferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	591	4776	3042	829	872	16	17	0	0	0
1	B	600	4857	3093	841	888	17	18	0	2	0

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	127	MSE	MET	modified residue	UNP Q9DC50
A	128	MSE	MET	modified residue	UNP Q9DC50
A	156	MSE	MET	modified residue	UNP Q9DC50
A	161	MSE	MET	modified residue	UNP Q9DC50
A	177	MSE	MET	modified residue	UNP Q9DC50
A	296	MSE	MET	modified residue	UNP Q9DC50
A	333	MSE	MET	modified residue	UNP Q9DC50
A	335	MSE	MET	modified residue	UNP Q9DC50
A	443	MSE	MET	modified residue	UNP Q9DC50
A	469	MSE	MET	modified residue	UNP Q9DC50
A	483	MSE	MET	modified residue	UNP Q9DC50
A	493	MSE	MET	modified residue	UNP Q9DC50
A	494	MSE	MET	modified residue	UNP Q9DC50
A	558	MSE	MET	modified residue	UNP Q9DC50
A	595	MSE	MET	modified residue	UNP Q9DC50
A	603	MSE	MET	modified residue	UNP Q9DC50
A	607	MSE	MET	modified residue	UNP Q9DC50
B	127	MSE	MET	modified residue	UNP Q9DC50
B	128	MSE	MET	modified residue	UNP Q9DC50
B	156	MSE	MET	modified residue	UNP Q9DC50
B	161	MSE	MET	modified residue	UNP Q9DC50
B	177	MSE	MET	modified residue	UNP Q9DC50
B	296	MSE	MET	modified residue	UNP Q9DC50
B	333	MSE	MET	modified residue	UNP Q9DC50
B	335	MSE	MET	modified residue	UNP Q9DC50

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	443	MSE	MET	modified residue	UNP Q9DC50
B	469	MSE	MET	modified residue	UNP Q9DC50
B	483	MSE	MET	modified residue	UNP Q9DC50
B	493	MSE	MET	modified residue	UNP Q9DC50
B	494	MSE	MET	modified residue	UNP Q9DC50
B	558	MSE	MET	modified residue	UNP Q9DC50
B	595	MSE	MET	modified residue	UNP Q9DC50
B	603	MSE	MET	modified residue	UNP Q9DC50
B	607	MSE	MET	modified residue	UNP Q9DC50

- Molecule 2 is CARNITINE (three-letter code: 152) (formula: $C_7H_{16}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	11	7	1	3	0	0

- Molecule 3 is OCTANOYLCARNITINE (three-letter code: OCB) (formula: $C_{15}H_{30}NO_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	B	1	20	15	1	4	0	0

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	B	1	8	6	2	0	0
4	B	1	8	6	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	B	1	8	6	2	0	0
			Total	C	O		
4	B	1	8	6	2	0	0
			Total	C	O		
4	B	1	8	6	2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	138	138	138	0	0
			Total	O		
5	B	276	276	276	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Peroxisomal carnitine O-octanoyltransferase

Chain A:



• Molecule 1: Peroxisomal carnitine O-octanoyltransferase

Chain B:




L512
I513
E516
E517
G518
L519
E523
L529
N538
S542
V546
G553
V554
V555
V559
H560
N561
G564
F565
F566
Y567
H568
I569
R570
D571
D572
R573
A577
S580
W581
R582
S583
C584
M595
I604
M607
N608
T609
A610
HIS
LEU

4 Data and refinement statistics (i)

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α, β, γ	163.61Å 163.61Å 159.48Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.09 – 2.20 29.08 – 2.20	Depositor EDS
% Data completeness (in resolution range)	95.2 (29.09-2.20) 95.4 (29.08-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.23 (at 2.20Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.199 , 0.230 0.197 , 0.228	Depositor DCC
R_{free} test set	5806 reflections (7.20%)	wwPDB-VP
Wilson B-factor (Å^2)	27.7	Xtriage
Anisotropy	0.059	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.35 , 57.3	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.019 for h,-h-k,-l	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	10118	wwPDB-VP
Average B, all atoms (Å^2)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.05% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 152, MPD, OCB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.31	0/4884	0.54	0/6585
1	B	0.36	0/4966	0.60	2/6694 (0.0%)
All	All	0.34	0/9850	0.57	2/13279 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	564	GLY	N-CA-C	-6.41	97.06	113.10
1	B	150	GLY	N-CA-C	-5.29	99.86	113.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4776	0	4676	123	0
1	B	4857	0	4762	105	0
2	A	11	0	15	3	0
3	B	20	0	29	8	0
4	B	40	0	70	2	0
5	A	138	0	0	4	0
5	B	276	0	0	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	10118	0	9552	231	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 231 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:568:HIS:HE1	1:A:570:ARG:HE	1.15	0.91
1:A:219:ARG:HH21	1:A:220:GLN:HE22	1.20	0.89
1:B:568:HIS:HE1	1:B:570:ARG:HE	1.17	0.89
1:B:252:ARG:HH21	1:B:270:GLN:HE22	1.23	0.86
1:B:480:GLN:HG3	1:B:607:MSE:HE1	1.58	0.84

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	587/612 (96%)	562 (96%)	23 (4%)	2 (0%)	41 46
1	B	600/612 (98%)	581 (97%)	19 (3%)	0	100 100
All	All	1187/1224 (97%)	1143 (96%)	42 (4%)	2 (0%)	47 55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	550	ARG
1	A	82	GLU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	522/524 (100%)	510 (98%)	12 (2%)	50	63
1	B	531/524 (101%)	520 (98%)	11 (2%)	53	67
All	All	1053/1048 (100%)	1030 (98%)	23 (2%)	52	65

5 of 23 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	139	LEU
1	B	297	LEU
1	B	265	LEU
1	B	312	ASN
1	A	312	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 36 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	270	GLN
1	B	594	GLN
1	B	292	GLN
1	B	470	GLN
1	A	490	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	MPD	B	616	-	7,7,7	0.48	0	9,10,10	0.47	0
2	152	A	613	-	10,10,10	1.17	1 (10%)	14,14,14	1.07	1 (7%)
4	MPD	B	618	-	7,7,7	0.49	0	9,10,10	0.48	0
3	OCB	B	613	-	19,19,19	0.93	2 (10%)	23,24,24	1.06	1 (4%)
4	MPD	B	617	-	7,7,7	0.46	0	9,10,10	0.42	0
4	MPD	B	614	-	7,7,7	0.48	0	9,10,10	0.47	0
4	MPD	B	615	-	7,7,7	0.44	0	9,10,10	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MPD	B	616	-	-	1/5/5/5	-
2	152	A	613	-	-	0/9/9/9	-
4	MPD	B	618	-	-	1/5/5/5	-
3	OCB	B	613	-	-	0/20/20/20	-
4	MPD	B	617	-	-	0/5/5/5	-
4	MPD	B	614	-	-	0/5/5/5	-
4	MPD	B	615	-	-	2/5/5/5	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	613	152	C4-N5	-2.24	1.47	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	613	OCB	CAO-NAN	-2.15	1.47	1.52
3	B	613	OCB	OAL-CAR	-2.09	1.23	1.30

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	613	152	C3-C4-N5	-3.41	111.00	116.83
3	B	613	OCB	OAK-CAH-CAG	3.19	118.39	111.50

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	616	MPD	C2-C3-C4-C5
4	B	615	MPD	C2-C3-C4-C5
4	B	618	MPD	C2-C3-C4-C5
4	B	615	MPD	C2-C3-C4-O4

There are no ring outliers.

4 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	616	MPD	1	0
2	A	613	152	3	0
3	B	613	OCB	8	0
4	B	615	MPD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand OCB B 613



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	574/612 (93%)	0.22	35 (6%)	21	20	23, 42, 64, 91	0
1	B	583/612 (95%)	-0.23	9 (1%)	73	72	14, 26, 48, 102	0
All	All	1157/1224 (94%)	-0.00	44 (3%)	40	38	14, 33, 63, 102	0

The worst 5 of 44 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	612	LEU	6.9
1	B	407	LYS	6.1
1	A	611	HIS	5.3
1	B	609	THR	4.6
1	A	610	ALA	4.3

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	MPD	B	617	8/8	0.70	0.34	62,64,65,66	0
4	MPD	B	616	8/8	0.74	0.33	68,69,70,70	0
4	MPD	B	618	8/8	0.78	0.31	75,76,77,77	0
4	MPD	B	615	8/8	0.83	0.33	68,68,72,72	0
2	152	A	613	11/11	0.86	0.19	28,33,35,36	0
4	MPD	B	614	8/8	0.91	0.21	48,50,52,53	0
3	OCB	B	613	20/20	0.95	0.17	20,24,36,37	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around OCB B 613:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.