wwPDB X-ray Structure Validation Summary Report (i)

May 24, 2020 – 01:27 am BST

PDB ID	:	4XNG
Title	:	Central Domain of Mycoplasma Genitalium Terminal Organelle protein MG491
Authors	:	Martinelli, L.; Fita, I.
Deposited on	:	2015-01-15
Resolution	:	3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	144	% 80% 16% • •
1	B	144	7% 82% 12% • 5%
1	C	144	% 82% 14% •
1	D	144	% 79% 19% ••

2 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 4588 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein MG218.1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	140	1155	738	196	219	1	1	0	0	0
1	B	137	1121	715	190	214	1	1	0	0	0
1	C	138	1135	723	193	217	1	1	0	0	0
1	D	144	1177	750	200	225	1	1	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	62	ALA	-	expression tag	UNP Q9ZB78
A	168	MSE	ILE	engineered mutation	UNP Q9ZB78
A	205	ALA	-	expression tag	UNP Q9ZB78
B	62	ALA	-	expression tag	UNP Q9ZB78
B	168	MSE	ILE	engineered mutation	UNP Q9ZB78
B	205	ALA	-	expression tag	UNP Q9ZB78
C	62	ALA	-	expression tag	UNP Q9ZB78
C	168	MSE	ILE	engineered mutation	UNP Q9ZB78
C	205	ALA	-	expression tag	UNP Q9ZB78
D	62	ALA	-	expression tag	UNP Q9ZB78
D	168	MSE	ILE	engineered mutation	UNP Q9ZB78
D	205	ALA	-	expression tag	UNP Q9ZB78

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Uncharacterized protein MG218.1

Chain A:



- Molecule 1: Uncharacterized protein MG218.1

Chain B:



- Molecule 1: Uncharacterized protein MG218.1

Chain C:



- Molecule 1: Uncharacterized protein MG218.1

Chain D:

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α, β, γ	98.27Å 108.42Å 62.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.00 47.47 – 2.90	Depositor EDS
% Data completeness (in resolution range)	95.3 (50.00-3.00) 89.7 (47.47-2.90)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.03 (at 2.91Å)	Xtriage
Refinement program	REFMAC 5.8.0069	Depositor
R, R_{free}	0.226 , 0.249 0.224 , 0.245	Depositor DCC
R_{free} test set	692 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å^2)	75.7	Xtriage
Anisotropy	0.067	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 57.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	4588	wwPDB-VP
Average B, all atoms (Å^2)	87.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.76 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.7535e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.60	0/1171	0.71	1/1568 (0.1%)
1	B	0.56	0/1135	0.66	0/1521
1	C	0.61	0/1150	0.73	0/1541
1	D	0.68	1/1193 (0.1%)	0.70	0/1598
All	All	0.61	1/4649 (0.0%)	0.70	1/6228 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	205	ALA	C-O	8.45	1.39	1.23

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	86	ARG	NE-CZ-NH1	5.18	122.89	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1155	0	1179	14	0
1	B	1121	0	1146	10	1
1	C	1135	0	1157	9	0
1	D	1177	0	1199	16	0
All	All	4588	0	4681	37	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 37 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:65:PHE:O	1:D:67:ASN:N	2.11	0.83
1:A:65:PHE:C	1:A:67:ASN:H	1.83	0.80
1:A:65:PHE:O	1:A:67:ASN:N	2.17	0.77
1:B:108:ASP:N	1:B:108:ASP:OD1	2.20	0.73
1:A:105:ARG:NH2	1:A:108:ASP:OD1	2.33	0.62

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:71:GLU:OE1	1:B:71:GLU:OE1[2_465]	2.15	0.05

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	138/144 (96%)	126 (91%)	10 (7%)	2 (1%)	11 43
1	B	135/144 (94%)	125 (93%)	7 (5%)	3 (2%)	6 31
1	C	136/144 (94%)	126 (93%)	7 (5%)	3 (2%)	6 31
1	D	142/144 (99%)	131 (92%)	8 (6%)	3 (2%)	7 33
All	All	551/576 (96%)	508 (92%)	32 (6%)	11 (2%)	7 34

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	66	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	107	PHE
1	B	108	ASP
1	D	66	HIS
1	D	108	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	132/133 (99%)	123 (93%)	9 (7%)	16	48
1	B	128/133 (96%)	120 (94%)	8 (6%)	18	51
1	C	130/133 (98%)	122 (94%)	8 (6%)	18	52
1	D	134/133 (101%)	125 (93%)	9 (7%)	16	49
All	All	524/532 (98%)	490 (94%)	34 (6%)	17	50

5 of 34 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	168	MSE
1	C	94	GLN
1	D	160	LYS
1	C	69	SER
1	A	168	MSE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	94	GLN
1	A	136	ASN
1	C	119	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	139/144 (96%)	-0.29	1 (0%)	87	69	43, 79, 132, 173	0
1	B	136/144 (94%)	0.23	10 (7%)	14	4	65, 100, 149, 189	0
1	C	137/144 (95%)	-0.12	2 (1%)	73	46	48, 78, 137, 156	0
1	D	143/144 (99%)	-0.18	1 (0%)	87	69	39, 70, 127, 161	0
All	All	555/576 (96%)	-0.09	14 (2%)	57	29	39, 83, 140, 189	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	65	PHE	3.5
1	C	106	SER	3.5
1	B	70	LYS	3.2
1	B	199	ILE	3.1
1	B	72	THR	2.8

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.