



# Full wwPDB X-ray Structure Validation Report i

Nov 2, 2020 – 10:35 PM GMT

PDB ID : 6YCS  
Title : Human Transcription Cofactor PC4 DNA-binding domain in complex with full phosphorothioate 5'-10-5' 2'-O-methyl DNA gapmer antisense oligonucleotide.  
Authors : Hyjek-Skladanowska, M.; Vickers, T.A.; Napiorkowska, A.; Anderson, B.; Tanowitz, M.; Crooke, S.T.; Liang, X.; Seth, P.P.; Nowotny, M.  
Deposited on : 2020-03-19  
Resolution : 3.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.1.3  
EDS : 2.14.6  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.14.6


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION

The reported resolution of this entry is 3.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1754 (3.10-3.02)
Clashscore	141614	1864 (3.10-3.02)
Ramachandran outliers	138981	1794 (3.10-3.02)
Sidechain outliers	138945	1793 (3.10-3.02)
RSRZ outliers	127900	1713 (3.10-3.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 3231 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PC4 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	66	Total	C	N	O	S	0	1	0
			525	333	91	98	3			
1	B	67	Total	C	N	O	S	0	0	0
			515	325	85	103	2			
1	C	68	Total	C	N	O	S	0	0	0
			525	333	91	97	4			
1	D	65	Total	C	N	O	S	0	0	0
			506	324	89	90	3			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	56	ARG	-	expression tag	UNP Q6IBA2
A	57	GLY	-	expression tag	UNP Q6IBA2
A	58	SER	-	expression tag	UNP Q6IBA2
A	59	HIS	-	expression tag	UNP Q6IBA2
A	60	SER	-	expression tag	UNP Q6IBA2
B	56	ARG	-	expression tag	UNP Q6IBA2
B	57	GLY	-	expression tag	UNP Q6IBA2
B	58	SER	-	expression tag	UNP Q6IBA2
B	59	HIS	-	expression tag	UNP Q6IBA2
B	60	SER	-	expression tag	UNP Q6IBA2
C	56	ARG	-	expression tag	UNP Q6IBA2
C	57	GLY	-	expression tag	UNP Q6IBA2
C	58	SER	-	expression tag	UNP Q6IBA2
C	59	HIS	-	expression tag	UNP Q6IBA2
C	60	SER	-	expression tag	UNP Q6IBA2
D	56	ARG	-	expression tag	UNP Q6IBA2
D	57	GLY	-	expression tag	UNP Q6IBA2
D	58	SER	-	expression tag	UNP Q6IBA2
D	59	HIS	-	expression tag	UNP Q6IBA2
D	60	SER	-	expression tag	UNP Q6IBA2


- Molecule 2 is a DNA chain called DNA (5'-D(P\*(OKQ))-D(P\*(OKT))-R(P\*(RFJ))-D(\*(OKQ)P\*(OKT)P\*(AS)P\*(GS)P\*(OKN)P\*(OKN)P\*(PST)P\*(OKN)P\*(PST)P\*(GS)P\*(GS)P\*(AS)P\*(OKT)P\*(OKT))-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace	
			Total	C	N	O	P	S				
2	E	16		660	314	106	176	32	32	0	16	0

- Molecule 3 is a DNA chain called DNA (5'-D(P\*(OKQ))-D(P\*(OKT))-R(P\*(RFJ))-D(\*(OKQ)P\*(OKT)P\*(AS)P\*(GS)P\*(OKN)P\*(OKN)P\*(PST)P\*(OKN)P\*(PST)P\*(GS)P\*(GS)P\*(AS)P\*(OKT)P\*(OKT))-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace	
			Total	C	N	O	P	S				
3	F	12		482	232	77	129	22	22	0	11	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	O	S			
4	A	1		5	4	1	0	0

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na<sup>+</sup>).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Na		
5	E	1		1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	2	Total O 2 2	0	0
6	B	3	Total O 3 3	0	0
6	C	3	Total O 3 3	0	0
6	D	2	Total O 2 2	0	0
6	E	2	Total O 2 2	0	0


### 3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PC4 protein




- Molecule 1: PC4 protein




- Molecule 1: PC4 protein



- Molecule 1: PC4 protein



- Molecule 2: DNA (5'-D(P\*(OKQ))-D(P\*(OKT))-R(P\*(RFJ))-D(\*(OKQ)P\*(OKT)P\*(AS)P\*(GS)P\*(OKN)P\*(OKN)P\*(PST)P\*(OKN)P\*(PST)P\*(GS)P\*(GS)P\*(AS)P\*(OKT)P\*(OKT))-3')



- Molecule 3: DNA (5'-D(P\*(OKQ))-D(P\*(OKT))-R(P\*(RFJ))-D(\*(OKQ)P\*(OKT)P\*(AS)P\*(GS)P\*(OKN)P\*(OKN)P\*(PST)P\*(OKN)P\*(PST)P\*(GS)P\*(GS)P\*(AS)P\*(OKT)P\*(OKT))-3')

Chain F: 



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	102.14Å    102.14Å    83.81Å 90.00°    90.00°    90.00°	Depositor
Resolution (Å)	45.68 – 3.05 45.68 – 3.05	Depositor EDS
% Data completeness (in resolution range)	99.8 (45.68-3.05) 99.9 (45.68-3.05)	Depositor EDS
$R_{merge}$	0.18	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	1.61 (at 3.06Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
$R$ , $R_{free}$	0.238 , 0.318 0.238 , 0.316	Depositor DCC
$R_{free}$ test set	445 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	90.8	Xtriage
Anisotropy	0.039	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 68.9	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.50$ , $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	3231	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	88.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.95% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: OKQ, GS, OKT, NA, PPS, SO4, RFJ, PST, OKN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.28	0/538	0.45	0/725
1	B	0.29	0/525	0.45	0/711
1	C	0.29	0/535	0.44	0/721
1	D	0.27	0/515	0.44	0/693
All	All	0.28	0/2113	0.45	0/2850

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	525	0	491	8	0
1	B	515	0	462	5	0
1	C	525	0	491	11	0
1	D	506	0	477	5	0
2	E	660	0	155	6	0
3	F	482	0	116	4	0
4	A	5	0	0	0	0
5	E	1	0	0	0	0
6	A	2	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	3	0	0	0	0
6	C	3	0	0	0	0
6	D	2	0	0	0	0
6	E	2	0	0	0	0
All	All	3231	0	2192	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:124:VAL:HG13	1:D:81:VAL:HG21	1.76	0.67
1:C:70:ARG:HH11	1:D:105:LEU:HD23	1.64	0.63
1:A:84:ASP:OD1	1:A:86:ARG:NH2	2.36	0.57
1:A:121:ASP:HA	1:A:124:VAL:HG12	1.88	0.56
3:F:6[A]:PPS:H5'2	3:F:6[A]:PPS:H8	1.89	0.54
1:A:114:LYS:HA	1:A:117:ILE:HG13	1.90	0.54
1:A:91:ASP:OD1	1:A:95:GLU:N	2.41	0.54
3:F:6[B]:PPS:H8	3:F:6[B]:PPS:H5'2	1.91	0.53
1:B:64:PHE:HB2	1:B:72:VAL:HG13	1.92	0.51
1:C:86:ARG:NH1	1:C:99:GLY:O	2.45	0.49
2:E:7[A]:GS:H8	2:E:7[A]:GS:H5"	1.96	0.48
1:C:65:GLN:O	1:D:116:GLN:NE2	2.47	0.47
2:E:14[A]:GS:S2P	2:E:14[A]:GS:H8	2.54	0.47
1:C:100:ARG:NH2	3:F:12[B]:PST:SP	2.88	0.46
1:A:123:ALA:HB1	1:B:74:VAL:HG21	1.98	0.46
1:C:86:ARG:HD2	1:C:98:PRO:CB	2.47	0.45
1:C:86:ARG:CB	1:C:102:GLY:HA3	2.46	0.45
1:B:101:LYS:NZ	2:E:16[A]:OKT:S01	2.84	0.44
2:E:10[A]:PST:O5'	2:E:10[A]:PST:H6	2.17	0.44
1:C:108:GLU:O	1:C:112:GLN:HG2	2.18	0.43
1:A:63:MET:HG2	1:A:73:SER:HB3	2.00	0.43
1:B:119:ASP:N	1:B:119:ASP:OD1	2.52	0.43
1:D:114:LYS:HA	1:D:117:ILE:HD11	2.00	0.42
1:A:84:ASP:HA	1:A:104:SER:HA	2.00	0.42
3:F:13[A]:GS:H8	3:F:13[A]:GS:O5'	2.19	0.42
1:C:86:ARG:HB2	1:C:102:GLY:HA3	2.02	0.42
1:A:100[A]:ARG:HG2	1:A:101:LYS:N	2.34	0.42
2:E:12[A]:PST:H6	2:E:12[A]:PST:H2"	1.74	0.42
1:C:110:TRP:CZ2	1:D:117:ILE:HG23	2.55	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:12[B]:PST:H2"	2:E:12[B]:PST:H6	1.81	0.40
1:B:110:TRP:O	1:B:114:LYS:HG3	2.22	0.40
1:C:86:ARG:HD2	1:C:98:PRO:HB2	2.03	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	65/72 (90%)	62 (95%)	3 (5%)	0	100 100
1	B	65/72 (90%)	63 (97%)	2 (3%)	0	100 100
1	C	66/72 (92%)	64 (97%)	2 (3%)	0	100 100
1	D	61/72 (85%)	59 (97%)	2 (3%)	0	100 100
All	All	257/288 (89%)	248 (96%)	9 (4%)	0	100 100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	52/65 (80%)	47 (90%)	5 (10%)	8 27
1	B	51/65 (78%)	51 (100%)	0	100 100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	C	52/65 (80%)	52 (100%)	0	100 100
1	D	49/65 (75%)	49 (100%)	0	100 100
All	All	204/260 (78%)	199 (98%)	5 (2%)	55 74

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	70	ARG
1	A	75	ARG
1	A	100[A]	ARG
1	A	100[B]	ARG
1	A	108	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

49 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	RFJ	E	3[A]	3,2	18,26,27	1.17	2 (11%)	20,38,41	2.73	6 (30%)
2	RFJ	E	3[B]	3,2	18,26,27	1.16	2 (11%)	20,38,41	2.69	6 (30%)
2	OKN	E	9[A]	3,2	15,21,22	0.70	1 (6%)	19,30,33	0.80	1 (5%)
2	OKN	E	8[B]	3,2	15,21,22	0.72	1 (6%)	19,30,33	1.01	1 (5%)
2	OKN	E	9[B]	3,2	15,21,22	0.71	1 (6%)	19,30,33	0.83	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	OKN	E	11[A]	2	15,21,22	0.74	1 (6%)	19,30,33	0.82	1 (5%)
2	OKN	E	11[B]	2	15,21,22	0.70	1 (6%)	19,30,33	0.81	1 (5%)
2	OKT	E	16[A]	2	10,14,23	0.40	0	11,19,34	1.07	1 (9%)
3	OKQ	F	4[A]	3,2	15,22,23	0.56	0	17,31,34	1.17	2 (11%)
2	OKT	E	16[B]	2	10,14,23	0.44	0	11,19,34	0.94	1 (9%)
3	GS	F	7[A]	3,2	18,24,25	1.19	2 (11%)	19,35,38	2.70	5 (26%)
3	OKQ	F	4[B]	3,2	15,22,23	0.57	0	17,31,34	1.10	2 (11%)
2	OKQ	E	4[A]	3,2	15,22,23	0.59	0	17,31,34	1.06	2 (11%)
2	OKQ	E	4[B]	3,2	15,22,23	0.59	0	17,31,34	1.03	2 (11%)
2	OKT	E	17[A]	2	14,22,23	1.22	1 (7%)	14,31,34	1.64	3 (21%)
2	OKT	E	17[B]	2	14,22,23	1.21	1 (7%)	14,31,34	1.70	2 (14%)
2	GS	E	14[A]	2	18,24,25	1.21	2 (11%)	19,35,38	2.65	5 (26%)
3	OKQ	F	1	3	15,19,23	0.55	0	17,27,34	1.15	2 (11%)
2	GS	E	14[B]	2	18,24,25	1.21	2 (11%)	19,35,38	2.65	5 (26%)
3	OKN	F	11[B]	3	15,21,22	0.71	1 (6%)	19,30,33	0.72	1 (5%)
3	OKT	F	5[A]	3	14,22,23	1.16	1 (7%)	14,31,34	1.61	3 (21%)
3	OKN	F	11[A]	3	15,21,22	0.72	1 (6%)	19,30,33	0.71	1 (5%)
3	OKN	F	8[B]	3,2	15,21,22	0.70	0	19,30,33	1.01	1 (5%)
3	OKT	F	5[B]	3	14,22,23	1.13	1 (7%)	14,31,34	1.37	1 (7%)
2	GS	E	7[B]	3,2	18,24,25	1.15	2 (11%)	19,35,38	2.75	4 (21%)
3	GS	F	13[A]	3	18,24,25	1.17	2 (11%)	19,35,38	2.69	4 (21%)
3	GS	F	13[B]	3	18,24,25	1.16	2 (11%)	19,35,38	2.69	4 (21%)
3	GS	F	7[B]	3,2	18,24,25	1.18	2 (11%)	19,35,38	2.68	5 (26%)
3	OKN	F	8[A]	3,2	15,21,22	0.71	0	19,30,33	1.04	1 (5%)
3	PST	F	10[A]	3	15,21,22	1.20	2 (13%)	16,30,33	3.64	1 (6%)
2	GS	E	7[A]	3,2	18,24,25	1.14	2 (11%)	19,35,38	2.77	5 (26%)
2	OKT	E	5[B]	2	14,22,23	1.12	1 (7%)	14,31,34	1.64	3 (21%)
3	PST	F	10[B]	3	15,21,22	1.19	2 (13%)	16,30,33	3.65	1 (6%)
2	OKT	E	5[A]	2	14,22,23	1.10	1 (7%)	14,31,34	1.40	2 (14%)
3	RFJ	F	3[B]	3,2	18,26,27	1.13	2 (11%)	20,38,41	2.70	6 (30%)
3	RFJ	F	3[A]	3,2	18,26,27	1.13	2 (11%)	20,38,41	2.70	6 (30%)
2	OKT	E	2[A]	2	14,22,23	1.08	1 (7%)	14,31,34	1.45	2 (14%)
3	OKT	F	2[B]	3,2	14,22,23	1.10	1 (7%)	14,31,34	1.32	2 (14%)
2	GS	E	13[B]	2	18,24,25	1.19	1 (5%)	19,35,38	2.65	4 (21%)
2	OKT	E	2[B]	2	14,22,23	1.08	1 (7%)	14,31,34	1.46	2 (14%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GS	E	13[A]	2	18,24,25	1.19	1 (5%)	19,35,38	2.65	4 (21%)
2	OKN	E	8[A]	3,2	15,21,22	0.73	1 (6%)	19,30,33	1.14	3 (15%)
3	OKT	F	2[A]	3,2	14,22,23	1.11	1 (7%)	14,31,34	1.38	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RFJ	E	3[A]	3,2	-	2/5/27/28	0/3/3/3
2	RFJ	E	3[B]	3,2	-	3/5/27/28	0/3/3/3
2	OKN	E	9[A]	3,2	-	0/3/21/22	0/2/2/2
2	OKN	E	8[B]	3,2	-	0/3/21/22	0/2/2/2
2	OKN	E	9[B]	3,2	-	2/3/21/22	0/2/2/2
2	OKN	E	11[A]	2	-	3/3/21/22	0/2/2/2
2	OKN	E	11[B]	2	-	2/3/21/22	0/2/2/2
2	OKT	E	16[A]	2	-	3/4/23/28	0/1/1/2
3	OKQ	F	4[A]	3,2	-	2/6/27/28	0/2/2/2
2	OKT	E	16[B]	2	-	2/4/23/28	0/1/1/2
3	GS	F	7[A]	3,2	-	1/2/21/22	0/3/3/3
3	OKQ	F	4[B]	3,2	-	0/6/27/28	0/2/2/2
2	OKQ	E	4[A]	3,2	-	0/6/27/28	0/2/2/2
2	OKQ	E	4[B]	3,2	-	0/6/27/28	0/2/2/2
2	OKT	E	17[A]	2	-	3/6/27/28	0/2/2/2
2	OKT	E	17[B]	2	-	2/6/27/28	0/2/2/2
2	GS	E	14[A]	2	-	1/2/21/22	0/3/3/3
3	OKQ	F	1	3	-	3/6/24/28	0/2/2/2
2	GS	E	14[B]	2	-	2/2/21/22	0/3/3/3
3	OKN	F	11[B]	3	-	3/3/21/22	0/2/2/2
3	OKT	F	5[A]	3	-	1/6/27/28	0/2/2/2
3	OKN	F	11[A]	3	-	2/3/21/22	0/2/2/2
3	OKN	F	8[B]	3,2	-	2/3/21/22	0/2/2/2
3	OKT	F	5[B]	3	-	1/6/27/28	0/2/2/2
2	GS	E	7[B]	3,2	-	2/2/21/22	0/3/3/3
3	GS	F	13[A]	3	-	2/2/21/22	0/3/3/3
3	GS	F	13[B]	3	-	2/2/21/22	0/3/3/3
3	GS	F	7[B]	3,2	-	1/2/21/22	0/3/3/3

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OKN	F	8[A]	3,2	-	2/3/21/22	0/2/2/2
3	PST	F	10[A]	3	-	0/4/21/22	0/2/2/2
2	GS	E	7[A]	3,2	-	0/2/21/22	0/3/3/3
2	OKT	E	5[B]	2	-	2/6/27/28	0/2/2/2
3	PST	F	10[B]	3	-	0/4/21/22	0/2/2/2
2	OKT	E	5[A]	2	-	2/6/27/28	0/2/2/2
3	RFJ	F	3[B]	3,2	-	0/5/27/28	0/3/3/3
3	RFJ	F	3[A]	3,2	-	2/5/27/28	0/3/3/3
2	OKT	E	2[A]	2	-	1/6/27/28	0/2/2/2
3	OKT	F	2[B]	3,2	-	1/6/27/28	0/2/2/2
2	GS	E	13[B]	2	-	2/2/21/22	0/3/3/3
2	OKT	E	2[B]	2	-	1/6/27/28	0/2/2/2
2	GS	E	13[A]	2	-	0/2/21/22	0/3/3/3
2	OKN	E	8[A]	3,2	-	0/3/21/22	0/2/2/2
3	OKT	F	2[A]	3,2	-	2/6/27/28	0/2/2/2

All (48) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	13[A]	GS	C6-N1	3.89	1.39	1.33
2	E	13[B]	GS	C6-N1	3.88	1.39	1.33
2	E	14[B]	GS	C6-N1	3.83	1.39	1.33
2	E	14[A]	GS	C6-N1	3.81	1.39	1.33
2	E	3[A]	RFJ	C6-N1	3.81	1.39	1.33
2	E	3[B]	RFJ	C6-N1	3.81	1.39	1.33
3	F	7[A]	GS	C6-N1	3.74	1.39	1.33
3	F	13[B]	GS	C6-N1	3.73	1.39	1.33
3	F	13[A]	GS	C6-N1	3.73	1.39	1.33
3	F	7[B]	GS	C6-N1	3.72	1.39	1.33
3	F	3[B]	RFJ	C6-N1	3.66	1.39	1.33
3	F	3[A]	RFJ	C6-N1	3.65	1.39	1.33
2	E	7[B]	GS	C6-N1	3.53	1.39	1.33
2	E	7[A]	GS	C6-N1	3.50	1.39	1.33
3	F	10[A]	PST	C4-N3	3.37	1.38	1.33
3	F	10[B]	PST	C4-N3	3.36	1.38	1.33
2	E	17[B]	OKT	C4-N3	3.26	1.38	1.33
2	E	17[A]	OKT	C4-N3	3.24	1.38	1.33
3	F	5[B]	OKT	C4-N3	3.09	1.38	1.33
2	E	2[A]	OKT	C4-N3	3.07	1.38	1.33
2	E	2[B]	OKT	C4-N3	3.06	1.38	1.33
3	F	2[B]	OKT	C4-N3	3.04	1.38	1.33

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	5[A]	OKT	C4-N3	3.04	1.38	1.33
3	F	2[A]	OKT	C4-N3	3.03	1.38	1.33
2	E	5[A]	OKT	C4-N3	3.02	1.38	1.33
2	E	5[B]	OKT	C4-N3	2.97	1.38	1.33
3	F	7[A]	GS	C8-N7	-2.21	1.30	1.34
3	F	7[B]	GS	C8-N7	-2.19	1.30	1.34
3	F	3[A]	RFJ	C8-N7	-2.18	1.30	1.34
2	E	7[A]	GS	C8-N7	-2.17	1.30	1.34
2	E	11[A]	OKN	C6-C5	-2.17	1.34	1.40
3	F	3[B]	RFJ	C8-N7	-2.14	1.30	1.34
3	F	10[A]	PST	C6-C5	-2.13	1.34	1.40
3	F	10[B]	PST	C6-C5	-2.13	1.34	1.40
2	E	14[A]	GS	C8-N7	-2.12	1.30	1.34
2	E	9[B]	OKN	C6-C5	-2.12	1.34	1.40
2	E	8[B]	OKN	C6-C5	-2.11	1.34	1.40
2	E	3[B]	RFJ	C8-N7	-2.11	1.30	1.34
2	E	7[B]	GS	C8-N7	-2.11	1.30	1.34
2	E	3[A]	RFJ	C8-N7	-2.10	1.31	1.34
2	E	14[B]	GS	C8-N7	-2.10	1.31	1.34
2	E	8[A]	OKN	C6-C5	-2.09	1.34	1.40
3	F	11[A]	OKN	C6-C5	-2.09	1.34	1.40
2	E	11[B]	OKN	C6-C5	-2.07	1.34	1.40
3	F	11[B]	OKN	C6-C5	-2.07	1.34	1.40
3	F	13[A]	GS	C8-N7	-2.04	1.31	1.34
2	E	9[A]	OKN	C6-C5	-2.03	1.34	1.40
3	F	13[B]	GS	C8-N7	-2.03	1.31	1.34

All (117) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	10[A]	PST	C4-N3-C2	14.26	127.18	115.14
3	F	10[B]	PST	C4-N3-C2	14.26	127.18	115.14
2	E	7[A]	GS	C5-C6-N1	-8.91	111.25	123.43
2	E	7[B]	GS	C5-C6-N1	-8.88	111.29	123.43
2	E	13[B]	GS	C5-C6-N1	-8.85	111.33	123.43
2	E	13[A]	GS	C5-C6-N1	-8.82	111.37	123.43
3	F	3[A]	RFJ	C5-C6-N1	-8.74	111.48	123.43
3	F	3[B]	RFJ	C5-C6-N1	-8.73	111.49	123.43
2	E	3[B]	RFJ	C5-C6-N1	-8.70	111.53	123.43
3	F	7[B]	GS	C5-C6-N1	-8.70	111.53	123.43
3	F	7[A]	GS	C5-C6-N1	-8.70	111.54	123.43
3	F	13[B]	GS	C5-C6-N1	-8.69	111.55	123.43

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	13[A]	GS	C5-C6-N1	-8.68	111.56	123.43
2	E	3[A]	RFJ	C5-C6-N1	-8.63	111.63	123.43
2	E	14[B]	GS	C5-C6-N1	-8.54	111.76	123.43
2	E	14[A]	GS	C5-C6-N1	-8.53	111.76	123.43
2	E	7[A]	GS	C6-N1-C2	5.99	125.45	115.93
3	F	3[B]	RFJ	C6-N1-C2	5.89	125.29	115.93
3	F	3[A]	RFJ	C6-N1-C2	5.86	125.24	115.93
2	E	7[B]	GS	C6-N1-C2	5.86	125.23	115.93
2	E	3[B]	RFJ	C6-N1-C2	5.85	125.23	115.93
2	E	3[A]	RFJ	C6-N1-C2	5.84	125.21	115.93
3	F	7[A]	GS	C6-N1-C2	5.81	125.17	115.93
3	F	7[B]	GS	C6-N1-C2	5.81	125.16	115.93
2	E	13[A]	GS	C6-N1-C2	5.79	125.14	115.93
2	E	13[B]	GS	C6-N1-C2	5.78	125.12	115.93
2	E	14[B]	GS	C6-N1-C2	5.75	125.07	115.93
2	E	14[A]	GS	C6-N1-C2	5.75	125.06	115.93
3	F	13[B]	GS	C6-N1-C2	5.73	125.03	115.93
3	F	13[A]	GS	C6-N1-C2	5.72	125.03	115.93
2	E	5[B]	OKT	C5-C4-N3	-4.00	114.50	123.31
2	E	2[B]	OKT	C5-C4-N3	-3.98	114.56	123.31
2	E	5[A]	OKT	C5-C4-N3	-3.98	114.56	123.31
2	E	2[A]	OKT	C5-C4-N3	-3.96	114.59	123.31
3	F	5[B]	OKT	C5-C4-N3	-3.92	114.68	123.31
3	F	2[A]	OKT	C5-C4-N3	-3.92	114.69	123.31
3	F	2[B]	OKT	C5-C4-N3	-3.90	114.73	123.31
2	E	17[B]	OKT	C5-C4-N3	-3.88	114.78	123.31
3	F	5[A]	OKT	C5-C4-N3	-3.80	114.95	123.31
2	E	17[A]	OKT	C5-C4-N3	-3.79	114.97	123.31
2	E	3[A]	RFJ	O2'-C2'-C1'	3.52	116.07	109.09
2	E	17[B]	OKT	C2'-C3'-C4'	-3.39	94.62	101.99
2	E	7[A]	GS	C2-N3-C4	-3.34	111.55	115.36
3	F	4[A]	OKQ	C2-N3-C4	3.21	119.59	116.34
3	F	1	OKQ	CM2-O2'-C2'	-3.20	106.14	114.52
2	E	7[B]	GS	C2-N3-C4	-3.19	111.72	115.36
3	F	3[B]	RFJ	C2-N3-C4	-3.16	111.75	115.36
3	F	3[A]	RFJ	C2-N3-C4	-3.12	111.80	115.36
3	F	7[A]	GS	C2-N3-C4	-3.10	111.82	115.36
2	E	14[A]	GS	C2-N3-C4	-3.08	111.84	115.36
3	F	7[B]	GS	C2-N3-C4	-3.08	111.84	115.36
2	E	14[B]	GS	C2-N3-C4	-3.06	111.86	115.36
3	F	3[A]	RFJ	C-O2'-C2'	-3.05	106.51	114.52
2	E	3[B]	RFJ	C2-N3-C4	-3.05	111.87	115.36

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	4[A]	OKQ	CM2-O2'-C2'	-3.04	106.55	114.52
3	F	4[B]	OKQ	C2-N3-C4	3.03	119.42	116.34
2	E	2[B]	OKT	C02-O01-C2'	-3.03	106.57	114.52
2	E	17[A]	OKT	C2'-C3'-C4'	-3.01	95.46	101.99
2	E	3[A]	RFJ	C2-N3-C4	-2.96	111.97	115.36
2	E	16[A]	OKT	C02-O01-C2'	-2.96	106.75	114.52
3	F	8[A]	OKN	C2-N3-C4	2.96	119.59	116.02
3	F	8[B]	OKN	C2-N3-C4	2.96	119.58	116.02
2	E	2[A]	OKT	C02-O01-C2'	-2.94	106.81	114.52
3	F	13[A]	GS	C2-N3-C4	-2.93	112.00	115.36
2	E	8[B]	OKN	C2-N3-C4	2.93	119.55	116.02
3	F	13[B]	GS	C2-N3-C4	-2.93	112.01	115.36
2	E	4[A]	OKQ	CM2-O2'-C2'	-2.92	106.86	114.52
2	E	13[B]	GS	C2-N3-C4	-2.91	112.03	115.36
3	F	3[B]	RFJ	C-O2'-C2'	-2.89	106.94	114.52
2	E	5[B]	OKT	C02-O01-C2'	-2.87	106.99	114.52
2	E	4[B]	OKQ	CM2-O2'-C2'	-2.87	107.00	114.52
2	E	13[A]	GS	C2-N3-C4	-2.87	112.08	115.36
3	F	5[A]	OKT	C2'-C3'-C4'	-2.87	95.77	101.99
3	F	4[B]	OKQ	CM2-O2'-C2'	-2.85	107.05	114.52
3	F	2[A]	OKT	C02-O01-C2'	-2.77	107.27	114.52
2	E	8[A]	OKN	C2-N3-C4	2.76	119.35	116.02
2	E	5[A]	OKT	C02-O01-C2'	-2.70	107.43	114.52
2	E	11[B]	OKN	C2-N3-C4	2.69	119.26	116.02
2	E	11[A]	OKN	C2-N3-C4	2.68	119.25	116.02
3	F	11[A]	OKN	C2-N3-C4	2.68	119.25	116.02
2	E	4[A]	OKQ	C2-N3-C4	2.67	119.04	116.34
3	F	11[B]	OKN	C2-N3-C4	2.66	119.23	116.02
2	E	9[A]	OKN	C2-N3-C4	2.66	119.23	116.02
2	E	9[B]	OKN	C2-N3-C4	2.66	119.22	116.02
2	E	4[B]	OKQ	C2-N3-C4	2.62	119.00	116.34
2	E	3[A]	RFJ	N3-C2-N1	-2.60	123.76	127.22
2	E	3[B]	RFJ	N3-C2-N1	-2.54	123.84	127.22
3	F	5[A]	OKT	C02-O01-C2'	-2.53	107.89	114.52
3	F	7[A]	GS	N3-C2-N1	-2.52	123.86	127.22
2	E	14[B]	GS	N3-C2-N1	-2.52	123.87	127.22
2	E	14[A]	GS	N3-C2-N1	-2.51	123.87	127.22
3	F	3[B]	RFJ	N3-C2-N1	-2.51	123.88	127.22
3	F	7[B]	GS	N3-C2-N1	-2.50	123.89	127.22
3	F	3[A]	RFJ	N3-C2-N1	-2.48	123.92	127.22
2	E	13[A]	GS	N3-C2-N1	-2.46	123.94	127.22
3	F	1	OKQ	C2-N3-C4	2.45	118.82	116.34

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	16[B]	OKT	C02-O01-C2'	-2.42	108.19	114.52
2	E	13[B]	GS	N3-C2-N1	-2.41	124.01	127.22
2	E	7[A]	GS	N3-C2-N1	-2.40	124.02	127.22
2	E	5[B]	OKT	O4'-C4'-C3'	-2.37	100.43	105.11
3	F	13[B]	GS	N3-C2-N1	-2.36	124.08	127.22
3	F	13[A]	GS	N3-C2-N1	-2.36	124.08	127.22
2	E	17[A]	OKT	O01-C2'-C1'	2.33	113.71	109.09
2	E	3[A]	RFJ	C6-C5-C4	-2.30	118.60	120.80
2	E	7[B]	GS	N3-C2-N1	-2.26	124.21	127.22
2	E	14[B]	GS	C6-C5-C4	-2.25	118.65	120.80
2	E	3[B]	RFJ	C-O2'-C2'	-2.22	108.69	114.52
2	E	3[B]	RFJ	C6-C5-C4	-2.22	118.68	120.80
2	E	14[A]	GS	C6-C5-C4	-2.21	118.68	120.80
3	F	2[B]	OKT	C02-O01-C2'	-2.20	108.74	114.52
2	E	8[A]	OKN	C2'-C3'-C4'	-2.18	98.23	102.76
3	F	7[A]	GS	C6-C5-C4	-2.14	118.75	120.80
3	F	3[B]	RFJ	C6-C5-C4	-2.12	118.77	120.80
3	F	7[B]	GS	C6-C5-C4	-2.09	118.80	120.80
2	E	7[A]	GS	C6-C5-C4	-2.06	118.83	120.80
2	E	8[A]	OKN	O4'-C4'-C3'	-2.05	100.89	105.67
3	F	3[A]	RFJ	C6-C5-C4	-2.01	118.88	120.80

There are no chirality outliers.

All (62) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	3[A]	RFJ	C1'-C2'-O2'-C
2	E	5[A]	OKT	C3'-C4'-C5'-O5'
2	E	5[A]	OKT	O4'-C4'-C5'-O5'
2	E	3[B]	RFJ	C1'-C2'-O2'-C
2	E	3[B]	RFJ	C3'-C4'-C5'-O5'
2	E	9[B]	OKN	C3'-C4'-C5'-O5'
2	E	11[A]	OKN	O4'-C1'-N1-C6
2	E	11[A]	OKN	O4'-C4'-C5'-O5'
2	E	11[B]	OKN	O4'-C1'-N1-C6
2	E	16[A]	OKT	C3'-C4'-C5'-O5'
2	E	16[A]	OKT	O4'-C4'-C5'-O5'
2	E	16[B]	OKT	O4'-C4'-C5'-O5'
2	E	17[A]	OKT	C1'-C2'-O01-C02
2	E	17[B]	OKT	C1'-C2'-O01-C02
3	F	1	OKQ	C2'-C1'-N1-C6
3	F	1	OKQ	O4'-C1'-N1-C6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	F	11[B]	OKN	O4'-C1'-N1-C6
3	F	5[A]	OKT	C1'-C2'-O01-C02
3	F	11[A]	OKN	O4'-C1'-N1-C6
3	F	5[B]	OKT	C1'-C2'-O01-C02
2	E	7[B]	GS	O4'-C4'-C5'-O5'
2	E	7[B]	GS	C3'-C4'-C5'-O5'
2	E	5[B]	OKT	C3'-C4'-C5'-O5'
2	E	5[B]	OKT	O4'-C4'-C5'-O5'
2	E	2[A]	OKT	O4'-C1'-N1-C6
3	F	2[B]	OKT	C1'-C2'-O01-C02
2	E	2[B]	OKT	O4'-C1'-N1-C6
3	F	2[A]	OKT	O4'-C4'-C5'-O5'
2	E	3[B]	RFJ	O4'-C4'-C5'-O5'
2	E	9[B]	OKN	O4'-C4'-C5'-O5'
2	E	11[A]	OKN	C3'-C4'-C5'-O5'
3	F	4[A]	OKQ	O4'-C4'-C5'-O5'
2	E	14[B]	GS	O4'-C4'-C5'-O5'
3	F	8[B]	OKN	C3'-C4'-C5'-O5'
3	F	13[B]	GS	O4'-C4'-C5'-O5'
3	F	3[A]	RFJ	O4'-C4'-C5'-O5'
3	F	2[A]	OKT	C3'-C4'-C5'-O5'
3	F	4[A]	OKQ	C3'-C4'-C5'-O5'
2	E	16[B]	OKT	C3'-C4'-C5'-O5'
3	F	8[B]	OKN	O4'-C4'-C5'-O5'
3	F	13[B]	GS	C3'-C4'-C5'-O5'
3	F	3[A]	RFJ	C3'-C4'-C5'-O5'
2	E	17[A]	OKT	O4'-C4'-C5'-O5'
3	F	8[A]	OKN	C3'-C4'-C5'-O5'
2	E	17[A]	OKT	C3'-C4'-C5'-O5'
3	F	8[A]	OKN	O4'-C4'-C5'-O5'
2	E	14[B]	GS	C3'-C4'-C5'-O5'
3	F	11[B]	OKN	O4'-C4'-C5'-O5'
2	E	13[B]	GS	C3'-C4'-C5'-O5'
2	E	3[A]	RFJ	C3'-C2'-O2'-C
2	E	13[B]	GS	O4'-C4'-C5'-O5'
2	E	17[B]	OKT	O4'-C4'-C5'-O5'
2	E	14[A]	GS	O4'-C4'-C5'-O5'
3	F	1	OKQ	O4'-C4'-C5'-O5'
3	F	13[A]	GS	O4'-C4'-C5'-O5'
3	F	7[B]	GS	C3'-C4'-C5'-O5'
2	E	11[B]	OKN	O4'-C4'-C5'-O5'
3	F	7[A]	GS	C3'-C4'-C5'-O5'

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	F	11[A]	OKN	O4'-C4'-C5'-O5'
3	F	13[A]	GS	C3'-C4'-C5'-O5'
2	E	16[A]	OKT	C3'-C2'-O01-C02
3	F	11[B]	OKN	C3'-C4'-C5'-O5'

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	16[A]	OKT	1	0
2	E	14[A]	GS	1	0
3	F	13[A]	GS	1	0
2	E	7[A]	GS	1	0

## 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [\(i\)](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	SO4	A	201	-	4,4,4	0.15	0	6,6,6	0.05	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
3	F	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	F	8[A]:OKN	O3'	10[A]:PST	P	9.73
1	F	8[B]:OKN	O3'	10[B]:PST	P	9.47

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	66/72 (91%)	0.24	1 (1%) 73   51	55, 82, 127, 149	0
1	B	67/72 (93%)	0.15	1 (1%) 73   51	51, 76, 133, 148	0
1	C	68/72 (94%)	1.02	16 (23%) 0   0	52, 95, 131, 151	0
1	D	65/72 (90%)	0.45	7 (10%) 5   2	62, 102, 137, 163	0
2	E	0/17	-	-	-	-
3	F	0/17	-	-	-	-
All	All	266/322 (82%)	0.47	25 (9%) 8   2	51, 89, 137, 163	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	113	LEU	5.2
1	C	110	TRP	4.3
1	C	109	GLN	3.7
1	A	127	LEU	3.7
1	C	105	LEU	3.6
1	C	106	ASN	3.6
1	D	65	GLN	3.5
1	C	111	SER	3.3
1	C	77	PHE	3.3
1	C	112	GLN	3.1
1	D	66	ILE	3.0
1	C	72	VAL	3.0
1	C	107	PRO	2.9
1	C	120	ILE	2.8
1	C	104	SER	2.8
1	D	64	PHE	2.8
1	D	71	TYR	2.4
1	B	83	ILE	2.4
1	C	83	ILE	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	81	VAL	2.4
1	C	108	GLU	2.3
1	D	83	ILE	2.2
1	C	82	LEU	2.1
1	D	63	MET	2.0
1	D	113	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	OKT	E	17[A]	21/22	0.70	0.32	107,108,108,110	21
2	OKT	E	17[B]	21/22	0.70	0.32	107,108,108,109	21
3	OKN	F	8[B]	20/21	0.79	0.21	101,102,103,103	20
3	OKN	F	8[A]	20/21	0.79	0.21	101,102,103,103	20
2	OKT	E	16[A]	14/22	0.83	0.28	96,104,105,107	14
2	OKT	E	16[B]	14/22	0.83	0.28	96,104,105,106	14
3	GS	F	13[A]	22/23	0.85	0.29	99,107,109,110	22
3	GS	F	13[B]	22/23	0.85	0.29	99,107,109,110	22
3	GS	F	7[B]	22/23	0.88	0.15	97,98,99,102	22
3	GS	F	7[A]	22/23	0.88	0.15	97,98,99,101	22
3	PST	F	12[B]	20/21	0.92	0.25	90,94,96,98	20
3	PST	F	12[A]	20/21	0.92	0.25	90,94,96,98	20
2	GS	E	14[B]	22/23	0.92	0.20	85,86,88,88	22
3	OKT	F	5[A]	21/22	0.92	0.20	78,83,85,87	21
2	GS	E	14[A]	22/23	0.92	0.20	84,86,88,88	22
3	OKT	F	5[B]	21/22	0.92	0.20	78,83,85,87	21
2	PST	E	10[B]	20/21	0.93	0.26	76,80,82,82	20
2	PST	E	10[A]	20/21	0.93	0.26	76,80,82,82	20
3	PST	F	10[A]	20/21	0.93	0.27	87,88,92,92	20
3	PST	F	10[B]	20/21	0.93	0.27	87,88,92,92	20
3	RFJ	F	3[B]	24/25	0.93	0.25	66,69,71,71	24
3	RFJ	F	3[A]	24/25	0.93	0.25	66,69,71,71	24
2	OKT	E	2[A]	21/22	0.93	0.28	76,78,84,84	21
2	GS	E	13[B]	22/23	0.93	0.21	77,84,86,87	22
2	OKT	E	2[B]	21/22	0.93	0.28	76,78,84,84	21
2	GS	E	13[A]	22/23	0.93	0.21	77,84,86,87	22
3	OKN	F	11[A]	20/21	0.94	0.24	88,89,90,90	20

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GS	E	7[B]	22/23	0.94	0.18	67,70,72,73	22
2	GS	E	7[A]	22/23	0.94	0.18	67,69,72,73	22
2	PST	E	12[B]	20/21	0.94	0.22	76,77,78,78	20
3	OKN	F	11[B]	20/21	0.94	0.24	87,89,90,90	20
2	PST	E	12[A]	20/21	0.94	0.22	76,77,78,78	20
2	OKN	E	9[B]	20/21	0.95	0.23	69,70,73,74	20
2	OKN	E	9[A]	20/21	0.95	0.23	69,70,73,75	20
3	OKQ	F	4[B]	21/22	0.95	0.18	71,74,74,76	21
3	OKQ	F	4[A]	21/22	0.95	0.18	70,74,74,77	21
2	OKN	E	11[A]	20/21	0.96	0.22	77,79,79,80	20
2	OKN	E	11[B]	20/21	0.96	0.22	77,79,79,80	20
2	OKT	E	5[B]	21/22	0.96	0.18	66,70,70,71	21
2	RFJ	E	3[B]	24/25	0.96	0.21	68,69,75,75	24
2	RFJ	E	3[A]	24/25	0.96	0.21	68,69,75,75	24
2	OKT	E	5[A]	21/22	0.96	0.18	66,70,70,71	21
2	OKN	E	8[B]	20/21	0.97	0.20	61,65,67,69	20
3	OKT	F	2[B]	21/22	0.97	0.20	64,65,66,66	21
3	OKQ	F	1	18/22	0.97	0.17	65,68,72,74	0
2	OKQ	E	4[A]	21/22	0.97	0.20	66,67,68,68	21
2	OKQ	E	4[B]	21/22	0.97	0.20	66,67,68,68	21
2	OKN	E	8[A]	20/21	0.97	0.20	61,64,67,68	20
3	OKT	F	2[A]	21/22	0.97	0.20	64,65,66,66	21

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	SO4	A	201	5/5	0.91	0.25	144,144,144,144	0
5	NA	E	101	1/1	0.95	0.47	57,57,57,57	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.