



wwPDB X-ray Structure Validation Summary Report ⓘ

Jul 24, 2023 – 12:12 PM JST

PDB ID : 7YN1
Title : Crystal structure of selenomethionine labeled CcbD
Authors : Mori, T.; Kadlcik, S.; Abe, I.
Deposited on : 2022-07-29
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.34
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.34


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	370	 79% 15% 6% 2%
1	B	370	 81% 13% 6% 2%
1	C	370	 79% 15% 6%
1	D	370	 82% 12% 6%

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 10926 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CcbD.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	347	Total 2673	C 1679	N 498	O 486	S 5	Se 5	0	0	0
1	B	348	Total 2695	C 1693	N 504	O 488	S 5	Se 5	0	0	0
1	C	347	Total 2691	C 1690	N 504	O 487	S 5	Se 5	0	0	0
1	D	347	Total 2683	C 1686	N 503	O 484	S 5	Se 5	0	0	0

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	356	LYS	-	expression tag	UNP E9JES9
A	357	LEU	-	expression tag	UNP E9JES9
A	358	ALA	-	expression tag	UNP E9JES9
A	359	ALA	-	expression tag	UNP E9JES9
A	360	ALA	-	expression tag	UNP E9JES9
A	361	LEU	-	expression tag	UNP E9JES9
A	362	GLU	-	expression tag	UNP E9JES9
A	363	HIS	-	expression tag	UNP E9JES9
A	364	HIS	-	expression tag	UNP E9JES9
A	365	HIS	-	expression tag	UNP E9JES9
A	366	HIS	-	expression tag	UNP E9JES9
A	367	HIS	-	expression tag	UNP E9JES9
A	368	HIS	-	expression tag	UNP E9JES9
A	369	HIS	-	expression tag	UNP E9JES9
A	370	HIS	-	expression tag	UNP E9JES9
B	356	LYS	-	expression tag	UNP E9JES9
B	357	LEU	-	expression tag	UNP E9JES9
B	358	ALA	-	expression tag	UNP E9JES9
B	359	ALA	-	expression tag	UNP E9JES9
B	360	ALA	-	expression tag	UNP E9JES9
B	361	LEU	-	expression tag	UNP E9JES9

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	362	GLU	-	expression tag	UNP E9JES9
B	363	HIS	-	expression tag	UNP E9JES9
B	364	HIS	-	expression tag	UNP E9JES9
B	365	HIS	-	expression tag	UNP E9JES9
B	366	HIS	-	expression tag	UNP E9JES9
B	367	HIS	-	expression tag	UNP E9JES9
B	368	HIS	-	expression tag	UNP E9JES9
B	369	HIS	-	expression tag	UNP E9JES9
B	370	HIS	-	expression tag	UNP E9JES9
C	356	LYS	-	expression tag	UNP E9JES9
C	357	LEU	-	expression tag	UNP E9JES9
C	358	ALA	-	expression tag	UNP E9JES9
C	359	ALA	-	expression tag	UNP E9JES9
C	360	ALA	-	expression tag	UNP E9JES9
C	361	LEU	-	expression tag	UNP E9JES9
C	362	GLU	-	expression tag	UNP E9JES9
C	363	HIS	-	expression tag	UNP E9JES9
C	364	HIS	-	expression tag	UNP E9JES9
C	365	HIS	-	expression tag	UNP E9JES9
C	366	HIS	-	expression tag	UNP E9JES9
C	367	HIS	-	expression tag	UNP E9JES9
C	368	HIS	-	expression tag	UNP E9JES9
C	369	HIS	-	expression tag	UNP E9JES9
C	370	HIS	-	expression tag	UNP E9JES9
D	356	LYS	-	expression tag	UNP E9JES9
D	357	LEU	-	expression tag	UNP E9JES9
D	358	ALA	-	expression tag	UNP E9JES9
D	359	ALA	-	expression tag	UNP E9JES9
D	360	ALA	-	expression tag	UNP E9JES9
D	361	LEU	-	expression tag	UNP E9JES9
D	362	GLU	-	expression tag	UNP E9JES9
D	363	HIS	-	expression tag	UNP E9JES9
D	364	HIS	-	expression tag	UNP E9JES9
D	365	HIS	-	expression tag	UNP E9JES9
D	366	HIS	-	expression tag	UNP E9JES9
D	367	HIS	-	expression tag	UNP E9JES9
D	368	HIS	-	expression tag	UNP E9JES9
D	369	HIS	-	expression tag	UNP E9JES9
D	370	HIS	-	expression tag	UNP E9JES9

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	36	Total O 36 36	0	0
2	B	43	Total O 43 43	0	0
2	C	57	Total O 57 57	0	0
2	D	48	Total O 48 48	0	0

- Molecule 1: CcbD

Chain D:  %



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	52.68Å 90.93Å 98.56Å 90.27° 103.67° 90.19°	Depositor
Resolution (Å)	26.13 – 2.50 47.88 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.7 (26.13-2.50) 98.7 (47.88-2.50)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.01 (at 2.51Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.203 , 0.240 0.203 , 0.240	Depositor DCC
R_{free} test set	1997 reflections (3.29%)	wwPDB-VP
Wilson B-factor (Å ²)	43.4	Xtrriage
Anisotropy	0.289	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 26.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.145 for h,-k,-h-l 0.013 for -h,k,-l 0.002 for -h,-k,h+l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	10926	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/2731	0.51	0/3713
1	B	0.24	0/2753	0.52	0/3740
1	C	0.23	0/2749	0.51	0/3733
1	D	0.24	0/2741	0.52	0/3723
All	All	0.24	0/10974	0.52	0/14909

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2673	0	2625	32	0
1	B	2695	0	2664	26	0
1	C	2691	0	2664	30	0
1	D	2683	0	2654	26	0
2	A	36	0	0	0	0
2	B	43	0	0	0	0
2	C	57	0	0	0	0
2	D	48	0	0	0	0
All	All	10926	0	10607	106	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 106 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:136:ASP:OD2	1:B:156:ARG:NH2	2.30	0.65
1:C:151:SER:O	1:C:154:ARG:NH1	2.32	0.63
1:B:99:LEU:HD22	1:B:135:PRO:HG2	1.81	0.63
1:B:116:ALA:HB1	1:B:167:ALA:HB1	1.80	0.62
1:D:37:VAL:HA	1:D:80:MSE:HE1	1.81	0.62

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	345/370 (93%)	340 (99%)	5 (1%)	0	100	100
1	B	346/370 (94%)	338 (98%)	8 (2%)	0	100	100
1	C	345/370 (93%)	337 (98%)	8 (2%)	0	100	100
1	D	345/370 (93%)	334 (97%)	11 (3%)	0	100	100
All	All	1381/1480 (93%)	1349 (98%)	32 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	266/282 (94%)	262 (98%)	4 (2%)	65 85
1	B	270/282 (96%)	262 (97%)	8 (3%)	41 68
1	C	270/282 (96%)	261 (97%)	9 (3%)	38 64
1	D	268/282 (95%)	265 (99%)	3 (1%)	73 89
All	All	1074/1128 (95%)	1050 (98%)	24 (2%)	52 77

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	192	ARG
1	C	263	ARG
1	C	258	SER
1	C	344	ASP
1	B	46	MSE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	342/370 (92%)	0.21	4 (1%) 79 80	29, 49, 72, 84	0
1	B	343/370 (92%)	0.28	6 (1%) 70 72	27, 50, 74, 94	0
1	C	342/370 (92%)	0.20	1 (0%) 94 94	28, 45, 68, 89	0
1	D	342/370 (92%)	0.22	3 (0%) 84 86	27, 45, 68, 90	0
All	All	1369/1480 (92%)	0.23	14 (1%) 82 84	27, 47, 71, 94	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	302	VAL	3.5
1	D	256	ASP	3.4
1	B	301	ALA	3.4
1	A	251	LEU	3.3
1	B	251	LEU	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.