Full wwPDB X-ray Structure Validation Report (i)

May 25, 2020 – 08:02 am BST

PDB ID	:	5YQJ
Title	:	Crystal structure of the first StARkin domain of Lam4
Authors	:	Im, Y.J.; Tong, J.S.
Deposited on	:	2017-11-06
Resolution	:	1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	187	6% 90% • 5%
1	B	187	4% 86% 9% • 5%
1	C	187	4% 91% • 6%

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 4968 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Membrane-anchored lipid-binding protein LAM4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	177	1376	883	223	267	3	0	0	0
			Total	C	N	O	S			
1	B	178	1386	888	227	268	3	0	0	0
			Total	C	N	O	S			
1	C	176	1369	879	222	265	3	0	0	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	743	GLY	-	expression tag	UNP P38800
A	744	SER	-	expression tag	UNP P38800
A	745	ALA	-	expression tag	UNP P38800
A	746	MET	-	expression tag	UNP P38800
A	747	GLY	-	expression tag	UNP P38800
A	748	SER	-	expression tag	UNP P38800
B	743	GLY	-	expression tag	UNP P38800
B	744	SER	-	expression tag	UNP P38800
B	745	ALA	-	expression tag	UNP P38800
B	746	MET	-	expression tag	UNP P38800
B	747	GLY	-	expression tag	UNP P38800
B	748	SER	-	expression tag	UNP P38800
C	743	GLY	-	expression tag	UNP P38800
C	744	SER	-	expression tag	UNP P38800
C	745	ALA	-	expression tag	UNP P38800
C	746	MET	-	expression tag	UNP P38800
C	747	GLY	-	expression tag	UNP P38800
C	748	SER	-	expression tag	UNP P38800

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	286	286	286	0	0
			Total	O		
2	B	286	286	286	0	0
			Total	O		
2	C	265	265	265	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Membrane-anchored lipid-binding protein LAM4

Chain A:





- Molecule 1: Membrane-anchored lipid-binding protein LAM4

Chain B:



- Molecule 1: Membrane-anchored lipid-binding protein LAM4

Chain C:

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	38.81Å 54.95Å 119.48Å 90.00° 95.08° 90.00°	Depositor
Resolution (Å)	29.75 – 1.50 29.75 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.5 (29.75-1.50) 99.5 (29.75-1.50)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	4.44 (at 1.50Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.158 , 0.211 0.158 , 0.211	Depositor DCC
R_{free} test set	4019 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å^2)	14.3	Xtriage
Anisotropy	0.455	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.38 , 56.1	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	4968	wwPDB-VP
Average B, all atoms (Å^2)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.37% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.32	0/1403	0.52	0/1909
1	B	0.32	0/1413	0.54	0/1921
1	C	0.31	0/1396	0.54	0/1899
All	All	0.32	0/4212	0.53	0/5729

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1376	0	1414	7	0
1	B	1386	0	1425	11	0
1	C	1369	0	1407	3	0
2	A	286	0	0	0	0
2	B	286	0	0	3	0
2	C	265	0	0	0	1
All	All	4968	0	4246	20	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:845:ASN:ND2	1:A:848:GLU:OE2	2.19	0.75
1:A:845:ASN:HB3	1:A:848:GLU:HG2	1.73	0.69
1:B:822:LYS:HE3	1:B:824:THR:HG22	1.75	0.68
1:B:753:TYR:OH	1:B:759:GLU:OE1	2.08	0.68
1:B:870:ARG:NH2	2:B:1005:HOH:O	2.27	0.67
1:B:921:GLU:OE1	2:B:1001:HOH:O	2.15	0.63
1:B:754:LYS:HD3	1:B:754:LYS:H	1.69	0.57
1:C:781:LYS:HE3	1:C:806:ASP:O	2.05	0.56
1:B:768:HIS:ND1	1:B:882:GLU:OE2	2.39	0.53
1:B:857:LYS:NZ	2:B:1009:HOH:O	2.38	0.52
1:C:834:LYS:HE2	1:C:836:MET:SD	2.50	0.52
1:A:923:GLY:O	1:A:927:THR:HG23	2.11	0.51
1:B:884:LYS:HZ1	1:B:886:LYS:NZ	2.12	0.48
1:A:848:GLU:CG	1:A:849:TYR:H	2.27	0.47
1:A:857:LYS:HG2	1:A:859:PRO:HD3	1.98	0.45
1:B:889:VAL:HG21	1:B:911:VAL:HG11	1.98	0.45
1:B:884:LYS:NZ	1:B:886:LYS:HD2	2.33	0.42
1:A:870:ARG:CZ	1:C:800:PRO:HD3	2.50	0.41
1:A:848:GLU:HG3	1:A:849:TYR:H	1.85	0.41
1:B:794:LYS:HD2	1:B:794:LYS:HA	1.93	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:1220:HOH:O	2:C:1235:HOH:O[1_655]	2.18	0.02

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	175/187 (94%)	170 (97%)	5 (3%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	176/187 (94%)	170 (97%)	6 (3%)	0	100	100
1	C	174/187 (93%)	169 (97%)	5 (3%)	0	100	100
All	All	525/561 (94%)	509 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	159/166 (96%)	159 (100%)	0	100	100
1	B	160/166 (96%)	157 (98%)	3 (2%)	57	27
1	C	158/166 (95%)	157 (99%)	1 (1%)	86	74
All	All	477/498 (96%)	473 (99%)	4 (1%)	81	66

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	754	LYS
1	B	807	SER
1	B	833	THR
1	C	833	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	177/187 (94%)	0.25	12 (6%)	17	18	10, 18, 48, 76	0
1	B	178/187 (95%)	0.21	8 (4%)	33	36	11, 17, 41, 97	0
1	C	176/187 (94%)	0.21	8 (4%)	33	36	10, 20, 35, 48	0
All	All	531/561 (94%)	0.22	28 (5%)	26	29	10, 18, 41, 97	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	752	VAL	10.6
1	A	751	THR	6.8
1	B	828	ILE	6.5
1	A	825	PRO	5.5
1	A	827	ALA	5.3
1	B	929	SER	5.3
1	A	752	VAL	4.6
1	C	927	THR	4.2
1	A	826	GLY	4.1
1	C	752	VAL	4.1
1	A	824	THR	4.0
1	A	927	THR	3.8
1	B	825	PRO	3.8
1	C	908	PHE	3.5
1	A	830	PRO	3.4
1	B	753	TYR	3.4
1	C	753	TYR	3.4
1	A	828	ILE	3.3
1	B	754	LYS	3.2
1	B	880	ASN	3.1
1	C	894	THR	2.8
1	B	755	PRO	2.8
1	C	768	HIS	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	901	SER	2.6
1	A	831	GLY	2.4
1	A	823	THR	2.3
1	A	829	GLY	2.2
1	C	757	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.