



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 14, 2023 – 01:36 AM JST

PDB ID : 5YSL
Title : Crystal structure of antibody 1H1 Fab
Authors : Hu, X.L.; Yang, F.L.
Deposited on : 2017-11-14
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	221	 79% 14% .. 5%
1	C	221	 84% 9% .. 5%
1	E	221	 80% 11% • 5%
1	G	221	 79% 13% • 5%
2	B	215	 82% 15% ..
2	D	215	 86% 10% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	215	 80% 16% ..
2	H	215	%  77% 20% ..

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 12923 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 1H1 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	211	Total 1589	C 1005	N 260	O 319	S 5	0	0	0
1	C	210	Total 1582	C 1000	N 259	O 318	S 5	0	0	0
1	E	210	Total 1582	C 1000	N 259	O 318	S 5	0	0	0
1	G	210	Total 1582	C 1000	N 259	O 318	S 5	0	0	0


- Molecule 2 is a protein called 1H1 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	213	Total 1647	C 1029	N 276	O 335	S 7	0	0	0
2	D	213	Total 1647	C 1029	N 276	O 335	S 7	0	0	0
2	F	213	Total 1647	C 1029	N 276	O 335	S 7	0	0	0
2	H	213	Total 1647	C 1029	N 276	O 335	S 7	0	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 1H1 heavy chain





- Molecule 1: 1H1 heavy chain




- Molecule 1: 1H1 heavy chain




- Molecule 1: 1H1 heavy chain




- Molecule 2: 1H1 light chain




- Molecule 2: 1H1 light chain



- Molecule 2: 1H1 light chain



- Molecule 2: 1H1 light chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	64.18Å 81.09Å 98.47Å 85.68° 82.85° 71.45°	Depositor
Resolution (Å)	50.00 – 2.50 46.66 – 2.48	Depositor EDS
% Data completeness (in resolution range)	97.2 (50.00-2.50) 97.3 (46.66-2.48)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.58 (at 2.48Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.212 , 0.273 0.217 , 0.274	Depositor DCC
R_{free} test set	3251 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	46.8	Xtrriage
Anisotropy	0.291	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 24.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12923	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 73.82 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.7141e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.77	0/1630	0.86	1/2228 (0.0%)
1	C	0.79	0/1623	0.86	2/2218 (0.1%)
1	E	0.74	0/1623	0.86	3/2218 (0.1%)
1	G	0.74	0/1623	0.86	2/2218 (0.1%)
2	B	0.74	0/1689	0.84	0/2299
2	D	0.74	0/1689	0.86	2/2299 (0.1%)
2	F	0.72	0/1689	0.86	3/2299 (0.1%)
2	H	0.74	0/1689	0.85	1/2299 (0.0%)
All	All	0.75	0/13255	0.86	14/18078 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	H	0	1

There are no bond length outliers.

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	183	LEU	CA-CB-CG	6.76	130.86	115.30
1	C	183	LEU	CA-CB-CG	6.57	130.41	115.30
2	H	118	ARG	NE-CZ-NH1	5.86	123.23	120.30
1	A	183	LEU	CA-CB-CG	5.78	128.60	115.30
2	D	103	ARG	NE-CZ-NH2	5.53	123.07	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	H	221	ARG	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1589	0	1543	13	0
1	C	1582	0	1534	6	0
1	E	1582	0	1534	11	0
1	G	1582	0	1534	13	0
2	B	1647	0	1567	14	0
2	D	1647	0	1567	11	0
2	F	1647	0	1567	15	0
2	H	1647	0	1567	17	0
All	All	12923	0	12413	89	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 89 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:122:THR:O	1:C:152:PHE:O	1.88	0.92
1:A:123:THR:HG23	2:H:136:THR:HG21	1.51	0.92
1:G:122:THR:O	1:G:152:PHE:O	1.92	0.87
2:H:201:SER:O	2:H:219:PHE:O	2.01	0.77
1:A:122:THR:O	1:A:152:PHE:O	2.02	0.77

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	207/221 (94%)	193 (93%)	9 (4%)	5 (2%)	6	9
1	C	206/221 (93%)	192 (93%)	12 (6%)	2 (1%)	15	28
1	E	206/221 (93%)	191 (93%)	12 (6%)	3 (2%)	10	18
1	G	206/221 (93%)	188 (91%)	14 (7%)	4 (2%)	8	13
2	B	211/215 (98%)	203 (96%)	6 (3%)	2 (1%)	17	31
2	D	211/215 (98%)	201 (95%)	7 (3%)	3 (1%)	11	20
2	F	211/215 (98%)	194 (92%)	12 (6%)	5 (2%)	6	9
2	H	211/215 (98%)	197 (93%)	9 (4%)	5 (2%)	6	9
All	All	1669/1744 (96%)	1559 (93%)	81 (5%)	29 (2%)	9	16

5 of 29 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	101	PHE
1	E	102	GLY
1	G	217	VAL
1	A	103	ASN
2	B	91	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	180/188 (96%)	160 (89%)	20 (11%)	6	11
1	C	179/188 (95%)	160 (89%)	19 (11%)	6	13
1	E	179/188 (95%)	158 (88%)	21 (12%)	5	10
1	G	179/188 (95%)	161 (90%)	18 (10%)	7	14
2	B	187/189 (99%)	167 (89%)	20 (11%)	6	13
2	D	187/189 (99%)	171 (91%)	16 (9%)	10	20

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	F	187/189 (99%)	168 (90%)	19 (10%)	7	14
2	H	187/189 (99%)	170 (91%)	17 (9%)	9	18
All	All	1465/1508 (97%)	1315 (90%)	150 (10%)	7	14

5 of 150 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	3	LYS
2	H	153	ASP
1	G	63	ARG
1	G	198	THR
1	C	111	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
2	F	100	GLN
2	F	220	ASN
2	F	171	ASN
2	F	222	ASN
2	D	99	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	211/221 (95%)	-0.09	3 (1%) 75 77	27, 46, 68, 82	0
1	C	210/221 (95%)	-0.12	0 100 100	27, 44, 68, 90	0
1	E	210/221 (95%)	0.01	1 (0%) 91 91	26, 51, 72, 95	0
1	G	210/221 (95%)	-0.03	2 (0%) 82 84	27, 49, 69, 84	0
2	B	213/215 (99%)	-0.22	1 (0%) 91 91	28, 45, 64, 89	0
2	D	213/215 (99%)	-0.22	1 (0%) 91 91	29, 46, 65, 89	0
2	F	213/215 (99%)	-0.14	0 100 100	31, 49, 66, 84	0
2	H	213/215 (99%)	-0.11	2 (0%) 84 86	29, 44, 64, 84	0
All	All	1693/1744 (97%)	-0.12	10 (0%) 89 90	26, 46, 68, 95	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	212	THR	3.8
1	E	75	THR	3.6
1	A	2	VAL	3.3
1	A	3	LYS	2.7
2	B	212	THR	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.